


RESEARCH ARTICLE

# Combination interventions for Hepatitis C and Cirrhosis reduction among people who inject drugs: An agent-based, networked population simulation experiment

Bilal Khan<sup>1</sup>, Ian Duncan<sup>1</sup>, Mohamad Saad<sup>1</sup>, Daniel Schaefer<sup>1</sup>, Ashly Jordan<sup>2,3</sup>, Daniel Smith<sup>2</sup>, Alan Neaigus<sup>4</sup>, Don Des Jarlais<sup>5</sup>, Holly Hagan<sup>2,3</sup>, Kirk Dombrowski<sup>1\*</sup>

**1** Department of Sociology, University of Nebraska, Lincoln NE, United States of America, **2** Rory Meyers College of Nursing, New York University, New York, NY, United States of America, **3** Center for Drug Use and HIV Research, New York University, New York, NY, United States of America, **4** Department of Epidemiology, Mailman School of Public Health, Columbia University, New York, NY, United States of America, **5** Icahn School of Medicine at Mount Sinai, New York, NY, United States of America

\* [kdombrowski2@unl.edu](mailto:kdombrowski2@unl.edu)



## OPEN ACCESS

**Citation:** Khan B, Duncan I, Saad M, Schaefer D, Jordan A, Smith D, et al. (2018) Combination interventions for Hepatitis C and Cirrhosis reduction among people who inject drugs: An agent-based, networked population simulation experiment. PLoS ONE 13(11): e0206356. <https://doi.org/10.1371/journal.pone.0206356>

**Editor:** Jason Blackard, University of Cincinnati College of Medicine, UNITED STATES

**Received:** January 24, 2018

**Accepted:** October 11, 2018

**Published:** November 29, 2018

**Copyright:** © 2018 Khan et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** The data presented here was produced via simulation. The field data used to parameterize the simulation was obtained via permission from the New York Department of Health and the US CDC and remains restricted. We do not have permission to make the original field data available to the public. Request for data access can be made to the New York Department of Health via Sarah Braunstein, Director, HIV Epidemiology and Field Services Program, NYC

## Abstract

Hepatitis C virus (HCV) infection is endemic in people who inject drugs (PWID), with prevalence estimates above 60% for PWID in the United States. Previous modeling studies suggest that direct acting antiviral (DAA) treatment can lower overall prevalence in this population, but treatment is often delayed until the onset of advanced liver disease (fibrosis stage 3 or later) due to cost. Lower cost interventions featuring syringe access (SA) and medically assisted treatment (MAT) have shown mixed results in lowering HCV rates below current levels. However, little is known about the potential cumulative effects of combining DAA and MAT treatment. While simulation experiments can reveal likely long-term effects, most prior simulations have been performed on closed populations of model agents—a scenario quite different from the open, mobile populations known to most health agencies. This paper uses data from the Centers for Disease Control's National HIV Behavioral Surveillance project, IDU round 3, collected in New York City in 2012 to parameterize simulations of open populations. To test the effect of combining DAA treatment with SA/MAT participation, multiple, scaled implementations of the two intervention strategies were simulated. Our results show that, in an open population, SA/MAT by itself has only small effects on HCV prevalence, while DAA treatment by itself can lower both HCV and HCV-related advanced liver disease prevalence. More importantly, the simulation experiments suggest that combinations of the two strategies can, when implemented together and at sufficient levels, dramatically reduce HCV incidence. We conclude that adopting SA/MAT implementations alongside DAA interventions can play a critical role in reducing the long-term consequences of ongoing HCV infection.

57. Turner KM, Hutchinson S, Vickerman P, Hope V, Craine N, Palmateer N, et al. The impact of needle and syringe provision and opiate substitution therapy on the incidence of hepatitis C virus in injecting drug users: pooling of UK evidence. *Addiction*. 2011; 106(11):1978–1988. <https://doi.org/10.1111/j.1360-0443.2011.03515.x> PMID: 21615585
58. Holtzman D, Barry V, Ouellet LJ, Des Jarlais DC, Vlahov D, Golub ET, et al. The influence of needle exchange programs on injection risk behaviors and infection with hepatitis C virus among young injection drug users in select cities in the United States, 1994–2004. *Preventive medicine*. 2009; 49(1):68–73. <https://doi.org/10.1016/j.ypmed.2009.04.014> PMID: 19410600
59. Tsui JI, Evans JL, Lum PJ, Hahn JA, Page K. Association of opioid agonist therapy with lower incidence of hepatitis C virus infection in young adult injection drug users. *JAMA internal medicine*. 2014; 174(12):1974–1981. <https://doi.org/10.1001/jamainternmed.2014.5416> PMID: 25347412
60. Neaigus A, Gyarmathy VA, Miller M, Frajzyngier VM, Friedman SR, Des Jarlais DC. Transitions to injecting drug use among noninjecting heroin users: social network influence and individual susceptibility. *JAIDS Journal of Acquired Immune Deficiency Syndromes*. 2006; 41(4):493–503. <https://doi.org/10.1097/01.qai.0000186391.49205.3b> PMID: 16652059
61. Czitrom V. One-factor-at-a-time versus designed experiments. *The American Statistician*. 1999; 53(2):126–131. <https://doi.org/10.2307/2685731>
62. Briggs AH, Weinstein MC, Fenwick EAL, Karnon J, Sculpher MJ, Paltiel AD. Model Parameter Estimation and Uncertainty Analysis: A Report of the ISPOR-SMDM Modeling Good Research Practices Task Force Working Group–6. *Medical Decision Making*. 2012; 32(5):722–732. <https://doi.org/10.1177/0272989X12458348> PMID: 22990087
63. Benhamou Y, Bochet M, Di Martino V, Charlotte F, Azria F, Coutellier A, et al. Liver fibrosis progression in human immunodeficiency virus and hepatitis C virus coinfecting patients. *Hepatology*. 1999; 30(4):1054–1058. <https://doi.org/10.1002/hep.510300409> PMID: 10498659
64. Bräu N, Salvatore M, Ríos-Bedoya CF, Fernández-Carbia A, Paronetto F, Rodríguez-Orengo JF, et al. Slower fibrosis progression in HIV/HCV-coinfecting patients with successful HIV suppression using anti-retroviral therapy. *Journal of hepatology*. 2006; 44(1):47–55. <https://doi.org/10.1016/j.jhep.2005.07.006> PMID: 16182404
65. Hu KQ, Cui W. Undetectable Serum HCV Antigens After 12 Weeks of HCV DAA Treatment Predicts Sustained Virologic Response. *Gastroenterology*. 2017; 152(5):S1092. [https://doi.org/10.1016/S0016-5085\(17\)33680-6](https://doi.org/10.1016/S0016-5085(17)33680-6)