

Lerdahl's tonal pitch space model and associated metric spaces

Richard R. Randall^{a*} and Bilal Khan^b

^a*School of Music, Carnegie Mellon University, Pittsburgh, PA, USA;* ^b*Department of Math & Computer Science, John Jay College of Criminal Justice, City University of New York, New York, NY, USA*

(Received 3 May 2007; final version received 7 July 2010)

This paper explores the boundary separating a theoretically derived model of chord relations from an empirically derived model. Fred Lerdahl's tonal pitch space (TPS) model approximates cognitive perceptual relations by providing a combinatorial procedure for computing the distance value between any two chords in a key. If TPS posits a hypothesized model of perception, then we would like to know if, and the extent to which, it differs from experimental data it claims to approximate. To achieve such a comparison, we develop three conceptual tools. First, we develop normalized canonical representations of each model, thereby avoiding comparisons affected by design choices. Second, we develop a distance measure that allows us to accurately compare the TPS model with another model derived from perceived chord relations described by Bharucha and Krumhansl. Finally, we use the distance measure to inform the design of a third model. These three models are shown to create a metric space of metric tonal models. The proposed distance measure and the method of normalization are applicable to any model with formal properties described herein and have the potential to focus experimental design and strengthen the relationship between experimental data and analytic systems.

Keywords: Lerdahl; tonal pitch space; music cognition; perception; metric space

1. Introduction

The search for compelling representations of tonal hierarchy and its constituent harmonic relations has a long-standing history. Lerdahl [1] describes geometric approaches to this problem that involve the collection and modelling of data from experiments in music cognition. The multidimensional-scaling models of Bharucha and Krumhansl [2], and Deutsch [3], for example, seek to encode cognitive relationships between chords *within a single key area* (region) as Euclidean distances. The work of Heinichen [4], Kellner [5], and Weber [6], is more speculative and when formalized, develops geometric representations of relationships between different key areas (regions) through their placement within a multidimensional space.

Fred Lerdahl's tonal pitch space (TPS) model [1] approximates the cognitive perceptual relation between chords by providing a combinatorial procedure for computing a distance value between two triads. The procedure employed by the TPS model is informed by experimental data and plausible hypotheses about how we perceive tonal relations. The model is extraordinarily powerful

*Corresponding author. Email: randall@cmu.edu

models. This served as a similarity measure allowing us to articulate specific ways the two models differed. Third, we showed how to derive canonical representatives. Comparing canonical representatives allowed us to critique analytic claims made by L against BK without the interference of arbitrary design choices. Fourth, we showed how the distance between canonical representatives provided the point of reference used to inform our design of a third model, F . Finally, we showed how these three tonal models are members of three different equivalence classes whose representatives each come from a metric space of tonal models. Since every subset of a metric space is a metric space, the three models form a metric space. In closing, it is important to mention that the choice of canonical representatives clearly influences the resulting distance judgement. In principle, it could be the case that where one choice would show $[\mathcal{M}_R^1]$ as closer to $[\mathcal{M}_R^2]$, than to $[\mathcal{M}_R^3]$, another choice might show $[\mathcal{M}_R^1]$ as closer to $[\mathcal{M}_R^3]$. The strategy we present above was designed to facilitate intuitive comparisons by choosing canonical representatives with the same minimal and maximal separations. Nevertheless, the choice of canonical representatives is an important issue that we hope to explore more fully in a future paper.

Acknowledgement

The authors wish to thank the reviewers for their comments and suggestions, many of which were incorporated into the final submission.

Notes

1. Noll and Garbers [7] provide an excellent critique of Lerdahl's TPS model in the context of theoretical problems associated with his attempt to combine a principle of hierarchy with a principle of shortest path. Their discussion is comprehensive and addresses issues outside the scope of what we present here.
2. An earlier form of this methodology was introduced in [8] in a broader context and in [9].
3. We identify the basic model as L to distinguish it from the larger TPS framework.
4. The data in Table 2 is shown in [10, Table 8.2]. However, [2] is given as the source.

References

- [1] F. Lerdahl, *Tonal Pitch Space*, Oxford University Press, New York, 2001.
- [2] J. Bharucha and C. Krumhansl, *The representation of harmonic structures in music: Hierarchies of stability as a function of context*, *Cognition* 13(1) (1983), pp. 63–103.
- [3] D. Deutsch, *The processing of pitch combinations*, in *The Psychology of Music*, Chapter 10, D. Deutsch, ed., Academic Press, New York, 1999.
- [4] J. D. Heinichen, *Der Generalbass in Der Composition*, Published by the author, Dresden, 1728.
- [5] D. Kellner, *Treulich Unterricht im General-Bass*, Christian Herold, Hamburg, 1737.
- [6] G. Weber, *Versuch Einer Geordenten Theorie Der Tonsetzkunst*, B. Schotts Sohne, Mainz, 1821–1824.
- [7] T. Noll and J. Garbers, *harmonic path analysis*, in *Perspectives of Mathematical and Computational Music Theory*, G. Mazzola, T. Noll, and E. Lluis-Puebla, eds., epOs-Music, Osnabrueck, 2004.
- [8] R.R. Randall, *A general theory of comparative music analysis*, Ph.D. thesis, Eastman School of Music, Rochester, 2006.
- [9] R. Randall and B. Khan, *Similarity measures for tonal models*, in *Proceedings of the International Conference on Music Perception and Cognition*, M. Baroni, A.R. Addessi, R. Casterina, and M. Costa, eds., Bologna, Italy, Bononia University Press, 2006.
- [10] C. Krumhansl, *Cognitive Foundations of Musical Pitch*, Oxford University Press, Oxford, 1990.
- [11] S. Shiralil and H.L. Vasudeva, *Metric Spaces*, Springer-Verlag, London, 2006.
- [12] H. Riemann, *Simplified Harmony*, Augener, London, 1896.
- [13] H. Riemann, *Systematische Modulationslehre als Grundlage der musikalischen Formenlehre*, J.F. Richter, Hamburg, 1887.
- [14] D. Harrison, *Harmonic Function in Chromatic Music*, University of Chicago Press, Chicago, IL, 1994.