

PROTOCOL Open Access

## Hepatitis C virus infection among HIV-positive men who have sex with men: protocol for a systematic review and meta-analysis

Holly Hagan<sup>1\*</sup>, Joshua Neurer<sup>2</sup>, Ashly E Jordan<sup>1</sup>, Don C Des Jarlais<sup>3</sup>, Jennifer Wu<sup>4</sup>, Kirk Dombrowski<sup>5</sup>, Bilal Khan<sup>6</sup>, Ronald Scott Braithwaite<sup>4</sup> and Jason Kessler<sup>4</sup>

## **Abstract**

**Background:** Outbreaks of hepatitis C virus (HCV) infection have been reported in HIV-positive men who have sex with men (MSM) in North America, Europe and Asia. Transmission is believed to be the result of exposure to blood during sexual contact. In those infected with HIV, acute HCV infection is more likely to become chronic, treatment for both HIV and HCV is more complicated and HCV disease progression may be accelerated. There is a need for systematic reviews and meta-analyses to synthesize the epidemiology, prevention and methods to control HCV infection in this population.

**Methods/design:** Eligible studies will include quantitative empirical data related to sexual transmission of HCV in HIV-positive MSM, including data describing incidence or prevalence, and associations between risk factors or interventions and the occurrence or progression of HCV disease. Care will be taken to ensure that HCV transmission related to injection drug use is excluded from the incidence estimates. Scientific databases will be searched using a comprehensive search strategy. Proceedings of scientific conferences, reference lists and personal files will also be searched. Quality ratings will be assigned to each eligible report using the Newcastle–Ottawa scale. Pooled estimates of incidence rates and measures of association will be calculated using random effects models. Heterogeneity will be assessed at each stage of data synthesis.

**Discussion:** HIV-positive MSM are a key HCV-affected population in the US and other high-income countries. This review seeks to identify modifiable risk factors and settings that will be the target of interventions, and will consider how to constitute a portfolio of interventions to deliver the greatest health benefit. This question must be considered in relation to the magnitude of HCV infection and its consequences in other key affected populations, namely, young prescription opioid users who have transitioned to illicit opiate injection, and older injection drug users among whom HCV prevalence and incidence are extremely high. This review is part of a series of systematic reviews and meta-analyses that will synthesize the evidence across all these population groups and develop recommendations and decision tools to guide public health resource allocation.

Trial registration: PROSPERO registration number: CRD42013006462

Keywords: Hepatitis C virus, Men who have sex with men, HIV infection, Epidemiology, Systematic review

<sup>\*</sup> Correspondence: hh50@nyu.edu <sup>1</sup>College of Nursing, New York University, New York, USA Full list of author information is available at the end of the article



- Cooper H, Hedges LV: The Handbook of Research Synthesis. New York: Russell Sage Foundation; 1994.
- 41. Raudenbush SW: Analyzing effect sizes: Random effects models. In *The Handbook of Research Synthesis*. Edited by Cooper H, Hedges LV. New York: Russell Sage Foundation; 1994:301–321.
- 42. Morton S, Adams J, Suttorp M, Shekelle P, Rockville MD:

  Meta-regression Approaches: What, Why, When, and How?. Rockville, MD:

  Agency for Healthcare Research and Quality; 2004. Report No.: 04-0033.
- 43. Thompson SG, Higgins JP: How should meta-regression analyses be undertaken and interpreted? *Stat Med* 2002, **21**(11):1559–1573.
- 44. Centers for Disease Control and Prevention (CDC): Hepatitis C virus infection among adolescents and young adults Massachusetts, 2002–2009. *Am J Transplant* 2011, 11(7):1535–1538.
- Centers for Disease Control and Prevention (CDC): Use of enhanced surveillance for hepatitis C virus infection to detect a cluster among young injection-drug users – New York, November 2004–April 2007. MMWR Morb Mortal Wkly Rep 2008, 57(19):517–521.
- Catalano RF, White HR, Fleming CB, Haggerty KP: Is nonmedical prescription opiate use a unique form of illicit drug use? Addict Behav 2011. 36(1–2):79–86.
- Smyth BP, O'Connor JJ, Barry J, Keenan E: Retrospective cohort study examining incidence of HIV and hepatitis C infection among injecting drug users in Dublin. J Epidemiol Community Health 2003, 57(4):310–311.
- Maher L, Jalaludin B, Chant KG, Jayasuriya R, Sladden T, Kaldor JM, Sargeant PL: Incidence and risk factors for hepatitis C seroconversion in injecting drug users in Australia. Addiction 2006, 101(10):1499–1508.

## doi:10.1186/2046-4053-3-31

Cite this article as: Hagan *et al.*: Hepatitis C virus infection among HIV-positive men who have sex with men: protocol for a systematic review and meta-analysis. *Systematic Reviews* 2014 3:31.

## Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit

