



← Back

Q1 Integer Variable Creation

2 marks

Which statement correctly creates an integer variable x with value 25?

A. `x='25'`

B. `x=25`

C. `int x=25`

D. `x:=25`

 Save Answer



← Back

Q2 Float Data Type

2 marks

What is the output of `type(3.0).__name__`?

A. int

B. float

C. complex

D. str

 Save Answer



← Back

Q3 Complex Number Assignment

2 marks

Which assignment creates a complex value?

A. `a=3+4j`

B. `a='3+4j'`

C. `a=(3,4)`

D. `a=complex`

 Save Answer



← Back

Q5 Input Function Return Type

2 marks

Which datatype is returned by input() before conversion?

A. int

B. float

C. str

D. bool

 Save Answer



← Back

Q6 Immutable Object

2 marks

Which object is immutable?

A. list

B. dict

C. tuple

D. set

 Save Answer



Q7 String to Integer Conversion

2 marks

Which converts '45' to an integer?

A. `int('45')`

B. `str('45')`

C. `float(45)`

D. `eval45`

 Save Answer



← Back

Q8 Boolean Variable Assignment

2 marks

Which assignment stores a Boolean?

A. flag='True'

B. flag=1

C. flag=True

D. flag='False'

 Save Answer



← Back

Q9 Boolean Data Type

2 marks

What is type(True)?

A. int

B. bool

C. str

D. float

 Save Answer



← Back

Q10 Hexadecimal Literal

2 marks

Which literal represents hexadecimal 26?

A. 0b11010

B. 0o32

C. 0x1A

D. 26H

 Save Answer



← Back

Q11 Binary to Decimal Conversion

2 marks

Decimal value of 0b1011 is

A. 9

B. 10

C. 11

D. 13

 Save Answer



← Back

Q12 Hexadecimal to Decimal Conversion

2 marks

Decimal value of 0x1F is

A. 31

B. 30

C. 32

D. 29

 Save Answer



← Back

Q13 Octal to Decimal Conversion

2 marks

Decimal value of 0o17 is

A. 15

B. 16

C. 17

D. 18

 Save Answer



← Back

Q14 Decimal to Binary Conversion

2 marks

Binary representation of decimal 10 is

A. 0b1001

B. 0b1010

C. 0b1110

D. 0b1100

 Save Answer



← Back

Q15 Complex Number Addition

2 marks

Result of $(2+3j)+(1-2j)$ is

A. $3+j$

B. $3+1j$

C. $1+5j$

D. $2+1j$

 Save Answer



← Back

Q16 Complex Number Identification

2 marks

Which is a complex number?

A. 3.5

B. 5

C. $2+7j$

D. '2+7j'

 Save Answer



← Back

Q17 Absolute Value Function

2 marks

Result of abs(-12) is

A. -12

B. 12

D. 1

 Save Answer



← Back

Q18 Division Result Type

2 marks

Type of 5/2 is

A. int

B. float

C. complex

D. bool

 Save Answer



← Back

Q19 Round Function

2 marks

Value of round(7.6) is

A. 7

B. 8

C. 7.5

D. Error

 Save Answer



← Back

Q20 Integer Conversion

2 marks

Value of `int(8.9)` is

A. 9

B. 8

C. 8.9

D. Error

 Save Answer



← Back

Q21 Floor Division with Negative Dividend

2 marks

Result of $-10//4$ is

A. -2

B. -3

C. 2

D. 3

 Save Answer



← Back

Q22 Floor Division with Negative Divisor

2 marks

Result of $10// -4$ is

A. -2

B. -3

C. 2

D. 3

 Save Answer



← Back

Q23 Modulo with Negative Dividend

2 marks

Result of $-10\%3$ is

A. -1

B. 1

C. 2

D. -2

 Save Answer



← Back

Q24 Modulo with Negative Divisor

2 marks

Result of $10\% - 3$ is

A. 1

B. -1

C. 2

D. -2

 Save Answer



← Back

Q25 Exponentiation with Negative Base

2 marks

Result of $(-2)^{**3}$ is

A. 8

B. -8

C. -6

D. 6

 Save Answer



← Back

Q26 Operator Precedence

2 marks

Result of $-2^{**}2$ is

A. 4

B. -4

C. -2

D. 2

 Save Answer



← Back

Q27 Exponentiation Result

2 marks

Result of $(-2)**2$ is

A. -4

B. 2

C. 4

D. 8

 Save Answer



← Back

Q28 True Division

2 marks

Result of $7/2$ is

A. 3

B. 3.0

C. 3.5

D. 4

 Save Answer



← Back

Q29 Operator Precedence

2 marks

Highest precedence among +, *, ** is

A. +

B. *

C. **

D. Equal

 Save Answer



← Back

Q30 Modulo Operation

2 marks

Result of $17\%5$ is

A. 2

B. 3

C. 4

D. 1

 Save Answer



← Back

Q31 Logical AND Operator

2 marks

5>3 and 2<1 evaluates to

A. True

B. False

C. None

D. Error

 Save Answer



← Back

Q32 Identity Operator

2 marks

Which operator checks object identity?

A. 0x

B. is

C. in

D. =

 Save Answer