COMPUTATIONAL GENOMICS / TECHNICAL RESEARCH REPORT

Genomic Investigation of MRSA Virulence and Antimicrobial Resistance

Comparative analysis framework for public *Staphylococcus aureus* assemblies

Tharuniya Anpalakan

Project Author

Study status (revalidated 6 November 2026). Four versioned public assemblies have been acquired and assessed with an exploratory FASTA quality-control screen. Resistance and virulence gene calling, molecular typing, and genotype–phenotype comparisons were not performed because the required software, databases, or metadata were unavailable. This document reports the completed assembly analysis without treating planned analyses as findings.

Abstract

Background. Genome sequencing can identify known resistance and virulence determinants in *Staphylococcus aureus*, provided that assemblies, reference databases, and isolate metadata are adequately documented. **Objective.** This study establishes a reproducible workflow for a small collection of public reference assemblies and evaluates the assembly data available before molecular screening. **Methods.** Four distinct, versioned NCBI RefSeq assemblies were selected as a technical demonstration cohort. Downloaded FASTA files were checked for valid sequence identifiers, non-empty records, and valid characters; assembly length, GC content, N50, sequence-record count, and ambiguous-base fraction were calculated. A predefined exploratory screen was applied. **Results.** All four files passed this screen. Assembly lengths ranged from 2,820,462 to 2,917,469 bp; sequence-record counts ranged from one to four. Four automated tests and six notebooks executed successfully. **Interpretation.** These are assembly-level observations, not evidence of completeness, antimicrobial susceptibility, toxin expression, or clinical phenotype. AMRFinderPlus, MLST, and SCCmec analyses were not performed in the present environment. The cohort cannot support population prevalence or acquisition-setting comparisons.

1 Introduction

Staphylococcus aureus is a Gram-positive coccus that colonizes the skin and mucosa of humans and can cause skin and soft-tissue infection, bacteremia, pneumonia, endocarditis, and other invasive disease. Colonization, host susceptibility, local tissue conditions, and strain-specific traits all contribute to the transition from carriage to infection [1, 2]. Methicillin-resistant *S. aureus* (MRSA) is of particular interest because resistance to many beta-lactams can constrain treatment and because successful lineages circulate through both community and healthcare settings. Epidemiological labels describe where and how infection was acquired; they cannot be assigned solely from a genomic signature.

This project asks: *What antimicrobial resistance and virulence determinants can be identified in a reproducibly selected collection of publicly available S. aureus genomes, and how do their distributions vary across the available isolate metadata?* The question is conditional. A sequence-based screen can address gene content only when the caller, reference database, and target coverage are validated. A comparison between isolate groups requires authentic group metadata, and a genotype–phenotype comparison additionally requires susceptibility measurements. The current report therefore presents the completed acquisition and assembly QC, then specifies how the later stages should be carried out.

2 Research objectives

1. Preserve exact assembly accession, BioSample, source-URL, and checksum provenance.
2. Validate FASTA structure and summarize elementary assembly properties using explicit decision rules.
3. Screen for AMR and virulence-associated determinants only if a versioned caller and database can be executed and reviewed.

4. Assign MLST or SCCmec types only when dedicated typing evidence is complete and adequately supported.
5. Compare the four assemblies descriptively without interpreting this selected cohort as a population sample or inferring phenotype from genotype alone.

3 Biological and genomic context

3.1 Methicillin resistance and SCCmec

The most familiar MRSA mechanism is acquisition of *mecA*, which encodes the alternative penicillin-binding protein PBP2a. Because PBP2a has reduced affinity for many beta-lactam agents, cross-linking of peptidoglycan can continue when native penicillin-binding proteins are inhibited. PBP2a is not simply a point-mutated version of a native protein: it is the product of an acquired determinant. A divergent homologue, now termed *mecC*, has also been documented in human and animal-associated *S. aureus* [2, 3].

Methicillin-resistance determinants are commonly embedded within staphylococcal cassette chromosome *mec* (SCCmec), a mobile element with a *mec* gene complex, cassette chromosome recombinase genes, and variable joining regions. SCCmec type is defined by the element's architecture. A *mecA* hit alone cannot identify SCCmec IV or any other type [4]. The historical association of some community-associated MRSA lineages with compact SCCmec IV elements is useful background, but it is neither an exclusive molecular rule nor a substitute for patient-level epidemiological information [5, 6].

3.2 Tetracycline resistance

Tetracyclines act at the bacterial 30S ribosomal subunit and interfere with aminoacyl-tRNA entry during translation. Acquired resistance can arise through efflux, including Tet(K) and Tet(L), or ribosomal protection, including Tet(M) and Tet(O) [7]. The presence of a named determinant does not, by itself, specify a susceptibility category for every tetracycline-class compound; expression, allele, drug, breakpoint, and laboratory method matter. Conversely, a negative sequence call cannot establish susceptibility, especially when assembly quality, gene divergence, or database coverage is uncertain. In this study no isolate-level tetracycline phenotype has been verified.

3.3 Virulence markers and PVL

Panton–Valentine leukocidin (PVL) is a bicomponent toxin encoded by *lukS-PV* and *lukF-PV*. The protein subunits act together to form pores in susceptible host leukocytes; receptor studies implicate human C5a receptors in cell targeting [8, 9]. Hemolysins and adhesion-associated factors, including alpha-toxin and clumping or fibronectin-binding proteins, contribute to the wider virulence repertoire. Genomic detection is evidence of a sequence related to a known determinant, not direct evidence of transcription, protein abundance, toxin activity, host response, or disease severity.

PVL has been associated with particular skin and soft-tissue presentations, but its clinical contribution differs by syndrome and study design. A systematic review found that broad claims about PVL as an independent determinant of severe disease are not warranted [10]. PVL status is therefore treated as a potential genomic observation to be assessed after validated screening, not as a defining feature of community acquisition.

4 Study design and data provenance

4.1 Selection rule

Four distinct, named, complete NCBI RefSeq assemblies were fixed before QC: N315, MW2, MRSA252, and USA300_FPR3757. Their original sequencing publications provide historical context [11, 5, 12, 6]. This is a compact, deliberately selected reference-genome demonstration rather than a random or representative sample of circulating MRSA. One versioned assembly per named strain was retained. No selection was made on the basis of the QC outcomes reported below.

Table 1: Public genome cohort. BioSample accessions and genome URLs are recorded in the repository manifest. Empty clinical fields were not imputed.

Strain	RefSeq assembly	BioSample	Verified contextual note
N315	GCF_000009645.1	SAMD00061099	Published reference strain
MW2	GCF_000011265.1	SAMD00061104	Published reference strain
MRSA252	GCF_000011505.1	SAMEA1705935	Hospital-associated description in BioProject
USA300_FPR3757	GCF_000013465.1	SAMN02604150	USA300 lineage; acquisition setting of specimen unverified

The manifest preserves source URLs, accession versions, assembly level, source database, metadata availability, and a selection rationale. Collection year, country, isolation source, sequencing platform, and antimicrobial susceptibility data remain blank where no reliable isolate-level record was established. The genomes were obtained on 17 October 2026. The compressed files are excluded from Git; observed SHA-256 digests are retained in the tracked metadata directory so a future download can be checked. Versioned accession identifiers and source links permit independent retrieval [13].

4.2 Scope of comparison

The four assemblies represent different named reference strains and historical studies. That diversity is useful for exercising a pipeline, but the cohort is too small and purposively assembled for population prevalence estimates or significance testing. The available metadata do not establish a consistent community-versus-healthcare classification for each sequenced specimen. The report consequently makes no such group comparison and no inference about transmission.

5 Computational methods

5.1 Acquisition and FASTA validation

The acquisition module downloads each exact HTTPS 'genomic.fna.gz' file listed in the manifest. It rejects malformed or duplicate RefSeq accessions, checks the expected NCBI URL prefix, verifies gzip integrity and FASTA start, and compares downloaded compressed-file hashes with the tracked SHA-256 values. The QC module streams each sequence record and rejects empty files, duplicate or missing record identifiers, and characters outside A, C, G, T, and N. These checks detect obvious file and formatting problems; they cannot establish read-level quality or assembly completeness.

5.2 Assembly metrics and decision rule

Total length is the sum of all FASTA-record lengths, including plasmid records where present. GC fraction is the number of G and C bases divided by that total; ambiguous fraction is the number of N bases divided by the same total. N50 is the length of the sequence record at which the cumulative length of records sorted from longest to shortest reaches at least half of the total assembly length. The quality screen flags an assembly for review if its total length lies outside 2.5–3.2 Mbp, if it has more than 20 records, or if more than 1% of bases are N. These pragmatic bounds were chosen for this complete-reference demonstration and are not a universal acceptance criterion for *S. aureus*. No independent completeness or contamination estimator was run.

5.3 Determinant screening and typing: specified, not executed

The repository includes an AMRFinderPlus wrapper for consistent assembly-level screening. Its specified invocation uses nucleotide assemblies, the *S. aureus* organism option, and `-plus` for curated non-AMR categories. The wrapper preserves the native tab-separated output, extracts positive AMR calls, and records software/database versions and reported identity and coverage values. A separate parser extracts explicitly labelled virulence calls. These methods follow the curated-reference approach described for AMRFinder and AMRFinderPlus [14, 15, 16]. The executable and database were not available in the present Windows environment; an execution attempt stopped explicitly before analysis, so no gene calls were made.

Before interpreting a non-call for *mecA*, *mecC*, a tet-family member, or either PVL component, the reference database must be audited for that target and the native output checked for partial or ambiguous alignments. A positive hit should be considered with its identity, reference coverage, method and genomic context. Sequence typing requires a validated multilocus-sequence-typing implementation and allele database [17]; SCCmec assignment requires cassette-level evidence [4]. Neither typing analysis was performed here.

5.4 Descriptive analysis

The completed statistics are the number of selected assemblies, the number passing exploratory QC, and observed metric ranges. A Wilson score interval is implemented in the code for future fully assessed binary outcomes [18], but no marker interval is reported because no complete,

validated marker screen exists. Gene-call status remains 'not assessed' rather than being silently recoded as absence. No hypothesis test was attempted on this four-genome convenience cohort.

6 Results: completed assembly quality control

All four versioned assemblies were retrieved and parsed. Each passed the predefined size, sequence-record-count, and ambiguous-base screen (Table 2). N315 and USA300_FPR3757 contain more than one FASTA record; the total lengths therefore include all supplied sequence records, not solely the main chromosome. The complete numerical output is available as a machine-readable CSV in the repository.

Table 2: Observed FASTA assembly metrics. GC and N percentages use total assembly length as denominator. 'Pass' applies only to the predefined exploratory screen.

Strain	Length (bp)	GC (%)	Records	N50 (bp)	N (%)	Screen
N315	2,839,469	32.808	2	2,814,816	0.000	Pass
MW2	2,820,462	32.830	1	2,820,462	0.000	Pass
MRSA252	2,902,619	32.807	1	2,902,619	0.000	Pass
USA300_FPR3757	2,917,469	32.691	4	2,872,769	0.000	Pass




Figure 1: Assembly length and FASTA-record count for the four downloaded RefSeq assemblies. The figure is generated directly from `results/genome_qc.csv`; it does not display resistance or virulence findings.

The observed length range is 2,820,462–2,917,469 bp, and the sequence-record count ranges from one to four. All downloaded sequences contain zero N bases under the implemented counting rule. All four compressed-file SHA-256 values matched the tracked provenance table on revalidation. The automated suite passed four of four tests, and all six notebooks executed successfully in memory. Repeated figure builds were byte-identical and the displayed figures were visually inspected. These observations neither validate species identity nor rule out contamination, structural errors, or missing genes. There are no AMR, virulence, MLST, SCCmec, or susceptibility results to report.

7 Discussion

The acquisition and QC stages establish a traceable starting point for comparative genomics. Exact accession versions, BioSample identifiers, URLs, checksums, and the screening rule make the completed stage inspectable. At the same time, this cohort is a poor basis for biological frequency estimates: it is small, historically selected, and drawn from distinct studies and lineages. The all-pass QC result indicates only that the downloaded assemblies meet the stated elementary thresholds, not that all biologically relevant loci are complete and accurately assembled.

Future AMRFinderPlus output could support statements such as 'a reference sequence assigned to a named determinant was detected under a documented database version and method'. It would not alone prove a resistant phenotype or a virulence effect. In particular, PVL gene detection would not establish active toxin production, and *mecA* detection would not establish an SCCmec type. Genotype–phenotype evaluation requires authenticated antimicrobial susceptibility records, drug-specific interpretive criteria, and attention to discordant calls. The present manifest contains no such phenotype data.

The historic source coursework described a patient-specific MRSA scenario but supplied no isolate accession, AST measurements, or sequence data for that patient. Those statements were therefore treated as background claims requiring review rather than transferred into this study's results. Original source figures were retained privately because reuse rights were not verified. No bacterial culture, sequencing, clinical testing, or wet-laboratory work is attributed to the computational project author.

8 Limitations and next analysis steps

- The four reference assemblies are a convenience cohort; inference to contemporary MRSA populations is inappropriate.
- The QC stage has not used raw reads or a validated completeness/contamination estimator.
- AMRFinderPlus and its database were unavailable and have not run on these assemblies. The code is ready for execution in a compatible environment, followed by native-output review and database coverage checks.
- MLST and SCCmec software/databases were unavailable. Typing needs dedicated tools and manual evaluation of confidence, especially if cassette structure is fragmented.
- No consistent collection, acquisition-setting, or authentic susceptibility metadata were verified; those comparisons must wait for suitable records.
- A phylogeny was omitted because homologous alignment, recombination assessment, and an appropriate sampling design have not been completed.

9 Conclusion

Four public, versioned *S. aureus* assemblies were acquired and passed a documented exploratory FASTA QC screen. The principal research questions about resistance and virulence determinants

remain open until validated screening is executed and its reference coverage reviewed. The resulting repository is a reproducible technical foundation for analysis, not a completed molecular survey or clinical diagnostic report.

Reproducibility statement

The repository contains the exact manifest, tracked download checksums, source code, tests, notebook views, figure-generation code, and a dated execution log. On a fresh clone, install the documented Python environment and run the acquisition, QC, and visualization modules from the repository root. This PDF was compiled from the LaTeX source in the `reports` directory with BibTeX entries in `references`. Run `python reports/build_latex.py` to rebuild it. Revalidation used Python 3.14.2, pandas 3.0.6, Matplotlib 3.11.2, and pytest 9.1.1; AMRFinderPlus, MLST, and SCCmec typing software were not installed or run.

References

[1] Steven Y. C. Tong, Joshua S. Davis, Emily Eichenberger, Thomas L. Holland, and Vance G. Fowler. Staphylococcus aureus infections: epidemiology, pathophysiology, clinical manifestations, and management. *Clinical Microbiology Reviews*, 28(3):603–661, 2015. doi:10.1128/CMR.00134-14.

[2] Timothy J. Foster. The Staphylococcus aureus superbug. *Journal of Clinical Investigation*, 114:1693–1696, 2004. doi:10.1172/JCI23825.

[3] Laura García-Álvarez et al. Meticillin-resistant Staphylococcus aureus with a novel mecA homologue in human and bovine populations in the UK and denmark: a descriptive study. *The Lancet Infectious Diseases*, 11:595–603, 2011. doi:10.1016/S1473-3099(11)70126-8.

[4] International Working Group on the Classification of Staphylococcal Cassette Chromosome Elements. Classification of staphylococcal cassette chromosome mec (SCCmec): guidelines for reporting novel SCCmec elements. *Antimicrobial Agents and Chemotherapy*, 53:4961–4967, 2009. doi:10.1128/AAC.00579-09.

[5] Tadashi Baba et al. Genome and virulence determinants of high virulence community-acquired MRSA. *The Lancet*, 359:1819–1827, 2002. doi:10.1016/S0140-6736(02)08713-5.

[6] Binh An Diep et al. Complete genome sequence of USA300, an epidemic clone of community-acquired meticillin-resistant Staphylococcus aureus. *The Lancet*, 367:731–739, 2006. doi:10.1016/S0140-6736(06)68231-7.

[7] Chuka U. Chukwudi. rRNA binding sites and the molecular mechanism of action of the tetracyclines. *Antimicrobial Agents and Chemotherapy*, 60:4433–4441, 2016. doi:10.1128/AAC.00594-16.

[8] Mohammad J. Aman and Rajan P. Adhikari. Staphylococcal bicomponent pore-forming toxins: Targets for prophylaxis and immunotherapy. *Toxins*, 6:950–972, 2014. doi:10.3390/toxins6030950.

[9] Andras N. Spaan et al. The staphylococcal toxin Panton-Valentine leukocidin targets human C5a receptors. *Cell Host and Microbe*, 13:584–594, 2013. doi:10.1016/j.chom.2013.04.006.

[10] Laura J. Shallcross, Ellen Fragaszy, Anne M. Johnson, and Andrew C. Hayward. The role of the Panton-Valentine leucocidin toxin in staphylococcal disease: a systematic review and meta-analysis. *The Lancet Infectious Diseases*, 13:43–54, 2013. doi:10.1016/S1473-3099(12)70238-4.

[11] Makoto Kuroda et al. Whole genome sequencing of meticillin-resistant Staphylococcus aureus. *The Lancet*, 357:1225–1240, 2001. doi:10.1016/S0140-6736(00)04403-2.

[12] Matthew T. G. Holden et al. Complete genomes of two clinical Staphylococcus aureus strains: evidence for the rapid evolution of virulence and drug resistance. *Proceedings of the National Academy of Sciences USA*, 101:9786–9791, 2004. doi:10.1073/pnas.0402521101.

[13] National Center for Biotechnology Information. NCBI datasets genome download documentation. `https://www.ncbi.nlm.nih.gov/datasets/docs/v2/reference-docs/command-line/datasets/download/genome/`. Accessed 6 November 2026.

[14] Michael Feldgarden et al. Validating the AMRFinder tool and resistance gene database by using antimicrobial resistance genotype-phenotype correlations in a collection of isolates. *Antimicrobial Agents and Chemotherapy*, 63:e00483–19, 2019. doi:10.1128/AAC.00483-19.

[15] Michael Feldgarden et al. AMRFinderPlus and the reference gene catalog facilitate examination of the genomic links among antimicrobial resistance, stress response, and virulence. *Scientific Reports*, 11:12728, 2021. doi:10.1038/s41598-021-91456-0.

[16] National Center for Biotechnology Information. AMRFinderPlus. `https://www.ncbi.nlm.nih.gov/pathogens/antimicrobial-resistance/AMRFinder/`. Accessed 6 November 2026.

[17] Mark C. Enright, Nicholas P. J. Day, Clare E. Davies, Sharon J. Peacock, and Brian G. Spratt. Multilocus sequence typing for characterization of methicillin-resistant and methicillin-susceptible clones of Staphylococcus aureus. *Journal of Clinical Microbiology*, 38:1008–1015, 2000. doi:10.1128/JCM.38.3.1008-1015.2000.

[18] Edwin B. Wilson. Probable inference, the law of succession, and statistical inference. *Journal of the American Statistical Association*, 22:209–212, 1927. doi:10.1080/01621459.1927.10502953.