Graviton Pressure Theory

The Unified Framework

Individual Submission

This document is part of a multi-part scientific framework

Part 23 of 30

The Mathematical Foundation

The Equations of the Medium, and What They Reproduce

This submission is part of the broader Graviton Pressure Theory (GPT) project, a redefinition of gravitational interaction rooted in causal field dynamics and physical force transmission. Each document is designed to stand on its own; its full context emerges as part of the larger framework. For the complete series, see https://peopleoftheworld.net/graviton-pressure-theory/.

Author: Shareef Ali Rashada

shareef@peopleoftheworld.net

Second edition, November 2026 (first edition 12 June 2025)

Contents

About This Paper **2**

23.1 The Quantities **2**

23.2 The Inflow and Its Acceleration **2**

23.3 Pressure **3**

23.4 Continuity **4**

23.5 Combining Flows **4**

23.6 Clocks **4**

23.7 Rulers, and the Whole Geometry **5**

23.8 Light **6**

23.9 Motion **7**

23.10 Rotation **7**

23.11 Waves in the Medium **8**

23.12 The Medium's Scale **8**

23.13 The Relaxation Boundary **9**

23.14 Summary **9**

Changes in the Second Edition **10**

About This Paper

This paper sets out the equations of Graviton Pressure Theory: the flow of the medium into a mass, its pressure, the motion of bodies and of light within it, the slowing of clocks, the effect of rotation, the waves the medium carries, and its scale. Each equation is checked against what is measured, and every other part of the framework cites this one set.

The central result is simple to state. A flowing medium in flat space, with light moving at c through it and with rods and clocks behaving as moving rods and clocks do, reproduces the predictions of General Relativity for a single mass exactly. The numbers that General Relativity calculates from the geometry of spacetime, GPT obtains from the motion of a medium. What General Relativity treats as geometry, GPT treats as the kinematics of rods and clocks moving through a medium. This form of the equations is known in the physics literature as the river model,[1] and the wider study of flows that mimic gravity as analogue gravity;[2] GPT reads it not as an analogy but as the mechanism.

The paper states, too, what the equations do not yet contain: the derivations still owed are listed with each topic and gathered at the end. Claims are marked by their status, as the key shows.

How to read the marks. Claims about Graviton Pressure Theory are marked by how far they are established.

Established	No mark. Accepted, measured physics.
Argued	Solid blue underline. A conclusion this framework argues in detail, which readers can check.
Hypothesis	Dashed orange underline. A proposal of GPT, not yet tested.
Open ○	Dotted grey underline and ○. A question not yet answered; work still owed.

A whole passage with one status carries a bar of the same style in the left margin. The marks are drawn in distinct line styles, so they read in black and white.

23.1 The Quantities

Table 23.1 lists the quantities used throughout. All are in SI units; the medium's pressure is in pascals.

23.2 The Inflow and Its Acceleration

HYPOTHESIS

The medium flows into a mass M along radial lines, with a speed calibrated once to Newton:

$$\mathbf{v}(r) = -\sqrt{\frac{2GM}{r}}\,\hat{\mathbf{r}}. \tag{1}$$

At the Earth's surface this is 11.2 kilometres per second, the escape speed; at the Sun's surface, 618 kilometres per second.

[1] Hamilton and Lisle, 2008.
[2] Barceló et al., 2011.

Symbol	Meaning	Value or unit
$\mathbf{v}$	velocity of the medium's inflow	m/s
ρ_g	density of the medium's active (pressure-bearing) part	about 10^{-26} kg/m^3
n	count density of gravitons (active and neutral)	m^{-3}
P	active pressure of the medium	Pa
P_0	active pressure in free space, $\frac{1}{2}\rho_g c^2$	about 4.5×10^{-10} Pa
c	speed of waves in the medium (light, gravitational waves)	2.998×10^8 m/s
Φ	Newtonian potential, $-GM/r$ for one mass	m^2/s^2
$\mathbf{u}$	velocity of a body, in the frame of distant matter	m/s
$\mathbf{w}$	velocity of a body through the medium, $\mathbf{u} - \mathbf{v}$	m/s
a_0	acceleration at the relaxation boundary	about 1.2×10^{-10} m/s^2

Table 23.1: Quantities used in this paper.

The medium accelerates as it flows in, and its acceleration is exactly Newton's gravity:

$$\mathbf{a}_{\text{medium}} = (\mathbf{v} \cdot \nabla)\mathbf{v} = v_r \frac{dv_r}{dr}\,\hat{\mathbf{r}} = -\frac{GM}{r^2}\,\hat{\mathbf{r}}. \tag{2}$$

Every body in the flow receives momentum from it in proportion to its own mass, at every particle at once (Part 7, Part 11), and so shares this acceleration:

$$\mathbf{F} = m\,\mathbf{a}_{\text{medium}} = -\frac{GMm}{r^2}\,\hat{\mathbf{r}}. \tag{3}$$

Because the delivery is proportional to mass, every body falls alike, whatever it is made of; the MICROSCOPE mission found this true to about one part in 10^{15}.[3]

The first edition wrote the force as $\mathbf{F} = -m\nabla P_g$. That expression has the wrong units (kilograms times pascals per metre is not a newton), and it cannot be repaired by a constant, because the pressure gradient acts on the medium, not on the body. The medium's own equation of steady flow (Euler's equation) gives the correct link: the medium's acceleration is set by its pressure gradient and its own density, and bodies share that acceleration.

$$(\mathbf{v} \cdot \nabla)\mathbf{v} = -\frac{\nabla P}{\rho_g}. \tag{4}$$

23.3 Pressure

Integrating Equation (4) along the inflow gives Bernoulli's relation:

$$P(r) = P_0 - \tfrac{1}{2}\rho_g v^2 = P_0 - \frac{\rho_g GM}{r} = P_0 + \rho_g\,\Phi. \tag{5}$$

The active pressure is lower near a mass, by an amount set by the Newtonian potential Φ. The free-space value is $P_0 = \frac{1}{2}\rho_g c^2$, the value at which the pressure law and the clock law of Section 23.6 agree. With it,

$$\frac{P}{P_0} = 1 - \frac{v^2}{c^2} = 1 - \frac{2GM}{c^2 r}. \tag{6}$$

[3] Touboul et al., 2022.

23.4 Continuity

A steady inflow neither piles up nor runs short only if as much medium crosses every sphere around the mass as any other. With the inflow speed of Equation (1), that requires the count of gravitons to rise toward the mass as $r^{-3/2}$:

$$4\pi r^2\, n(r)\, |v(r)| = Q = \text{constant} \quad \Longrightarrow \quad n(r) \propto r^{-3/2}. \tag{7}$$

This is the premise the framework already holds: the count of gravitons rises toward a mass while the active pressure falls (Part 14). The two are consistent because most of the medium near a mass has passed through matter and been ejected neutral: it is counted, but it does not press.

The relation between the intake Q and the mass M is still owed. ◦ If Q is proportional to M, the normalisation of n must scale as $\sqrt{M}$.

23.5 Combining Flows

HYPOTHESIS

Flows from several masses do not add as velocities; their pressures combine. If the pressure deficits add,

$$P = P_0 + \rho_g \sum_i \Phi_i, \qquad \Phi_i = -\frac{GM_i}{|\mathbf{r} - \mathbf{r}_i|}, \tag{8}$$

then the acceleration set up by the combined pressure is

$$\mathbf{a} = -\frac{\nabla P}{\rho_g} = -\nabla \sum_i \Phi_i, \tag{9}$$

which is Newton's law for any number of bodies. In a static distribution of matter of density ρ_m, the pressure obeys

$$\nabla^2 P = 4\pi G\, \rho_g\, \rho_m, \tag{10}$$

which is Poisson's equation of Newtonian gravity, written for the medium's pressure.

Combining by pressure recovers Newtonian dynamics for many bodies exactly: the orbits of the planets, the Lagrange points, the tides (Part 13). Adding the inflow velocities instead would not: at the Earth, the cross term between the Sun's inflow and the Earth's would be of order 40 to 70 metres per second squared, which is not observed. The direction of the velocity field around many bodies, and the full non-linear rule for strong fields, are still owed. ◦

23.6 Clocks

HYPOTHESIS

A clock is a structure of medium waves, and it ticks more slowly the faster the medium streams through it. A clock moving with velocity $\mathbf{u}$ through a medium flowing at $\mathbf{v}$ moves through the medium at $\mathbf{w} = \mathbf{u} - \mathbf{v}$, and its rate is

$$\frac{d\tau}{dt} = \sqrt{1 - \frac{|\mathbf{u} - \mathbf{v}|^2}{c^2}}. \tag{11}$$

For a clock held still ($\mathbf{u} = 0$), the medium streams past at the inflow speed, and Equations (6) and (11) give

$$\frac{d\tau}{dt} = \sqrt{1 - \frac{2GM}{c^2 r}} = \sqrt{\frac{P}{P_0}}. \tag{12}$$

Gravitational and motional time dilation are one effect: both are the speed of the medium through the clock. The rate depends on where the clock is and how it moves, not on what it is made of. Every kind of clock compared so far, from caesium and hydrogen masers to optical clocks of strontium, ytterbium, aluminium and mercury ions, slows by the same factor.

Checks.

- **Height.** Raising a clock by h near the Earth speeds it by gh/c^2: 2.5×10^{-15} over the 22.5 metres of the Pound–Rebka tower,[4] and 3.6×10^{-17} for the 33-centimetre rise measured with optical clocks.[5]
- **GPS.** A GPS satellite's clock runs faster by 45.7 microseconds a day, because the medium streams more slowly past it at altitude, and slower by 7.2 microseconds a day because of its orbital speed: a net 38.5 microseconds a day, which the system corrects for.
- **The Earth's surface.** With the medium streaming past at 11.2 kilometres per second, a clock on the ground runs slow by $1 - \sqrt{1 - v^2/c^2} = 6.96 \times 10^{-10}$, exactly GM/c^2R.
- **Fast muons.** Muons circulating at $\gamma \approx 29.3$ in a storage ring, under accelerations of about $10^{18}\ g$, lived γ times longer and no more: the rate depends on speed through the medium, not on acceleration.[6]

23.7 Rulers, and the Whole Geometry

HYPOTHESIS

A rod moving through the medium is shortened along its motion by the same factor that slows a clock, as moving rods are; this is the kinematics that Lorentz and FitzGerald proposed and that Bell showed follows from the dynamics of the rod's own structure.[a][b] A rod held still near a mass has the medium streaming through it along the radius, and so is shortened along the flow, and only along it.

[a] Bell, 1987.
[b] Brown, 2005.

Clocks slowed by Equation (11) and rods shortened along the flow, in a flat space through which the medium flows at Equation (1), are described together by one expression:

$$c^2 d\tau^2 = c^2 dt^2 - |\, d\mathbf{x} - \mathbf{v}\, dt \,|^2 . \tag{13}$$

This is the Schwarzschild solution of General Relativity, in the form Painlevé and Gullstrand found in 1921 and 1922.[7][8] Every prediction General Relativity makes for a single non-rotating mass therefore follows from GPT unchanged. General Relativity reads Equation (13) as the

[4] Pound and Rebka, 1960.
[5] Chou et al., 2010.
[6] Bailey et al., 1977.
[7] Painlevé, 1921.
[8] Gullstrand, 1922.

geometry of spacetime; GPT reads it as what rods and clocks do in a flowing medium. The numbers are the same; the cause is named.

The first edition's form, in which rulers shrank equally in every direction, gave the wrong advance of Mercury's perihelion (57″ per century instead of 43″). Rulers that shrink along the flow only give the right one.

23.8 Light

HYPOTHESIS

Light is a transverse wave of the medium and moves at c relative to the medium. A wave with wave vector $\mathbf{k}$ in a medium moving at $\mathbf{v}$ has frequency

$$\omega = c\,|\mathbf{k}| + \mathbf{k}\cdot\mathbf{v}. \tag{14}$$

Measured locally, by rods and clocks that are themselves in the medium, its speed is always c. Tracing rays with Equation (14) past a mass gives:

- **Bending**: $\Delta\theta = 4GM/c^2 b$ for a ray passing at distance b, which is 1.75″ at the Sun's edge. The traced rays give $\Delta\theta\, b\, c^2/GM = 4.016$, 4.002 and 4.0005 for $b = 10^3$, 10^4 and 10^5 times GM/c^2, approaching 4 as the path lengthens.
- **Delay**: a signal passing the Sun is delayed by $(2GM/c^3)\ln(4r_1r_2/b^2)$ each way, as Shapiro predicted and radar and spacecraft measured; the Cassini spacecraft confirmed it to about one part in 10^5.[9][10]
- **Redshift**: a signal climbing from a lower to a higher clock is received by a faster clock, so it appears redder by $\Delta\Phi/c^2$, as in Section 23.6.

In the language of refraction, the medium near a mass acts on light with an index $n \approx 1 + 2GM/c^2 r$. From Equation (6), that is

$$n \approx 2 - \frac{P}{P_0}\,: \tag{15}$$

the lower the active pressure, the higher the index, and the slower light travels as measured from far away. The first edition had the sign reversed, with light slowing where the pressure was higher.

No colour in the bending. Because light moves at c through the medium whatever its frequency, the bending is the same for radio and for visible light. It is: radio and optical measurements agree to a few parts in a thousand, and radio interferometry fixes the bending to about one part in 10^4. The first edition's suggestion of colour-dependent lensing is withdrawn.

[9] Shapiro, 1964.
[10] Bertotti et al., 2003.

23.9 Motion

HYPOTHESIS

A body moving through the medium at **w** takes in the medium faster at its leading face than at its trailing face, by the factor

$$k = \sqrt{\frac{1 + w/c}{1 - w/c}}. \tag{16}$$

The imbalance is its momentum, the path it carves (Part 11).

The arithmetic is exact. Half the sum of the front and back rates, $\frac{1}{2}(k + 1/k)$, is $\gamma = 1/\sqrt{1 - w^2/c^2}$; half the difference, $\frac{1}{2}(k - 1/k)$, is $\gamma w/c$. So the body's energy and momentum are

$$E = \gamma mc^2, \qquad \mathbf{p} = \gamma m\mathbf{w}. \tag{17}$$

Their shared rate, the geometric mean $\sqrt{k \cdot 1/k} = 1$, is the rest rate that runs the body's internal cycles, which is why a moving clock slows by $1/\gamma$; the excess $\gamma - 1$ is its kinetic energy. A force along the motion changes the speed at a cost γ^3 times that at rest, and a force across it at a cost γ. Approaching c, the front rate grows without bound, and so does the energy needed.

The path of a body moving freely in the medium follows from Equation (13): it is the path General Relativity calls a geodesic. For an orbit, it gives Kepler's ellipses with a slow advance of the perihelion,

$$\Delta\varpi = \frac{6\pi GM}{c^2 a(1 - e^2)} \text{ per orbit}, \tag{18}$$

which is 43.0″ per century for Mercury, as observed after the pull of the other planets (532″ per century) is removed.[11]

Matter and light differ in one respect. Matter is a standing pattern of the medium, with both a leading and a trailing exchange, and so a rest rate: a mass. Light is a travelling wave of the medium, with only a front, so its rest rate is zero: it has energy and momentum, $E = pc$, but no mass.

23.10 Rotation

HYPOTHESIS

A rotating mass gives the medium flowing into it a swirl, as a spinning drain gives swirl to the water entering it. Far from a body with angular momentum J, the swirl turns at

$$\omega(r) \approx \frac{2GJ}{c^2 r^3}, \tag{19}$$

and a gyroscope carried in the medium is turned with it.

This is the frame-dragging of General Relativity: the river model of a rotating black hole contains exactly this swirl.[12] Gravity Probe B measured the dragging of its gyroscopes around the Earth at 37.2 ± 7.2 milliarcseconds a year, against 39.2 predicted, and the precession of their spin axes by the inflow itself at 6602 ± 18 milliarcseconds a year, against 6606.[13] The orbits of the LAGEOS satellites are dragged as predicted, to about ten per cent.[14]

[11] Will, 2014.
[12] Hamilton and Lisle, 2008.
[13] Everitt et al., 2011.
[14] Ciufolini and Pavlis, 2004.

23.11 Waves in the Medium

HYPOTHESIS

The medium carries stress as well as pressure, and so it carries transverse waves. Light is one; gravitational waves, the medium's response to masses accelerating, are another. Both move at c.

What is measured fixes the form of these waves:

- **Speed.** Gravitational waves and light from the neutron-star merger of August 2017 arrived within two seconds of each other after travelling 130 million years: their speeds agree to about one part in 10^{15}.[15] In GPT this is expected, since both are waves of the same medium.
- **Polarisation.** With three detectors, the pattern of stretching fitted the two transverse, tensor polarisations better than scalar or vector patterns.[16] The waves are not pressure waves of the medium; a pressure wave would be a single, breathing polarisation.
- **Energy carried.** The orbit of the binary pulsar PSR B1913+16 shrinks exactly as the energy carried away by gravitational waves requires: the measured rate is 0.9983 ± 0.0016 times the predicted one.[17]

How one medium carries two kinds of transverse wave, vector waves for light and tensor waves for gravity, at the same speed, is still owed. ◦ The elastic models of the luminiferous medium in the nineteenth century are the precedent to study. The first edition's scalar wave equation for the pressure, with a speed that might exceed c, is withdrawn: it gives the wrong polarisation and the wrong speed.

23.12 The Medium's Scale

Only the inflow speed enters the motion of planets, clocks and light; the medium's density and free-space pressure do not. They cannot be measured from orbits or from the bending of light. They are fixed instead by what the medium may deliver as heat.

If matter absorbed the medium flowing into it and kept its energy as heat, the intake $4\pi R^2 \rho_g v_{\text{esc}}$, times c^2, would heat every planet and star. At $P_0 \approx 4.5 \times 10^{-10}$ pascals, $\rho_g \approx 1.0 \times 10^{-26}$ kilograms per cubic metre:

- the Earth would absorb 5.1×10^9 watts, one ten-thousandth of its measured heat flow of 47 terawatts;
- the Sun would absorb 3.4×10^{15} watts, a hundred-billionth of its output.

Both pass. The first edition's higher values, 10^{10} to 10^{12} pascals in Part 19 and 10^{-6} to 10^{-4} pascals in Part 25, are retired as the medium's free-space pressure: the first would heat the Earth beyond its measured heat flow by fifteen orders of magnitude and more. The 10^{10} to 10^{11} pascals of Part 19 survives as a different quantity, the energy density of the bonds that hold matter together (iron's cohesive energy is about 5.8×10^{10} joules per cubic metre).

The scale has a coincidence worth noting: $\frac{1}{2}\rho_c c^2$, for the critical density of the universe, is 3.8×10^{-10} pascals, and the pressure attributed to dark energy has a magnitude of 5.3×10^{-10}

[15] Abbott et al., 2017b.
[16] Abbott et al., 2017a.
[17] Weisberg and Huang, 2016.

pascals. In General Relativity the dark-energy pressure is negative, and the medium's is positive; any link must come through a different route, the expansion of a self-repelling medium that is not confined, and must be derived (Part 14).

The medium's equation of state, and the range and law of its self-repulsion, are still owed. ◦ One constraint is already clear: if the pressure were simply proportional to the density, compression waves would travel at $c/\sqrt{2}$, so light and gravitational waves cannot be compression waves. That agrees with Section 23.11. The first edition's inverse-square repulsion between gravitons is withdrawn: a medium whose parts repel at long range without limit has no equilibrium.

23.13 The Relaxation Boundary

Galaxies rotate as though held by more matter than they contain, and the departure from Newton follows a single pattern. Across 153 galaxies of every size, the acceleration observed at each radius is fixed by the acceleration expected from the visible matter alone, with a scatter of about 0.13 dex, by the relation[18]

$$g_{\mathrm{obs}} = \frac{g_{\mathrm{bar}}}{1 - e^{-\sqrt{g_{\mathrm{bar}}/a_0}}}, \qquad a_0 \approx 1.2 \times 10^{-10}\ \mathrm{m/s^2}. \tag{20}$$

Where g_{bar} is large, $g_{\mathrm{obs}} = g_{\mathrm{bar}}$: Newton holds. Where it is small, $g_{\mathrm{obs}} \to \sqrt{g_{\mathrm{bar}}\, a_0}$, and a star orbiting at a large radius moves at a speed that no longer falls with distance, $v^4 = GMa_0$ (Figure 23.1).

HYPOTHESIS

GPT places a boundary around every mass where its inflow's acceleration falls to a_0:

$$r_{a_0} = \sqrt{\frac{GM}{a_0}}. \tag{21}$$

Beyond it, the medium ejected neutral by the mass is recharged, and gravity changes its form. For the Sun, $r_{a_0} \approx 7{,}000$ astronomical units, far beyond the planets, so the solar system's tracking is untouched. For the Milky Way, with 6×10^{10} to 10^{11} solar masses of stars and gas, it is 8 to 11 kiloparsecs, where the rotation curve flattens.

The law of the boundary, from which Equation (20) must follow, is still owed. ◦ It is GPT's central distinctive prediction (Part 14). Tests at the scale of the Sun's boundary already exist: wide binary stars separated by thousands of astronomical units, whose orbits are currently disputed in both directions.[19][20]

23.14 Summary

Table 23.2 gathers the results. Every measurement that General Relativity passes for a single mass, GPT passes with the same numbers, because it contains the same equations; what it adds is the cause, a medium flowing through flat space, and a single prediction of its own at the scale of galaxies.

[18] McGaugh et al., 2016.
[19] Chae, 2023.
[20] Banik et al., 2024.




Figure 23.1: What the relaxation boundary must reproduce: the acceleration observed in galaxies against the acceleration expected from their visible matter, in the fitted form of the measured relation (Equation 20). Above a_0, Newton holds; below it, the observed acceleration approaches $\sqrt{g_{\rm bar}a_0}$, and rotation curves flatten. GPT places the change at the boundary where each mass's inflow falls to a_0.
HYPOTHESIS

The derivations still owed are: the relation between intake and mass (Section 23.4); the full rule for combining flows in strong fields (Section 23.5); the two transverse waves of one medium (Section 23.11); the equation of state (Section 23.12); and the law of the relaxation boundary (Section 23.13).

The path ahead is not curved. It is pressurised. And it is measurable.

Changes in the Second Edition

This list records what changed from the first edition (12 June 2025) and why. Changes to the argument alter what the paper claims or how it supports a claim; clarifications improve wording, sources or presentation and leave the argument as it was.

Changes to the argument

- The equations are replaced by the agreed account: the inflow calibrated to Newton, its acceleration, Bernoulli's pressure, continuity, combining by pressure, one clock law, rulers shortened along the flow, light at c through the medium, motion as the carved path, rotation as swirl, transverse waves, the scale and the relaxation boundary. Together they reproduce the Schwarzschild solution exactly, in Painlevé–Gullstrand form.
- $\mathbf{F} = -m\nabla P_g$ and $a = -\nabla P_g$ are withdrawn: their units do not balance, and the pressure gradient acts on the medium, which bodies share by delivery in proportion to mass.
- The pressure "field tension" $G_{ij} = \partial_i P^j - \partial_j P^i$, which treated a scalar pressure as a vector,

Effect	GPT	Measured
Falling	medium's acceleration GM/r^2, shared in proportion to mass	same for all bodies to 10^{-15}
Many bodies	pressures combine; $\mathbf{a} = -\nabla \sum \Phi_i$	planetary motion, tides
Clocks	$\sqrt{1 - \lvert\mathbf{u} - \mathbf{v}\rvert^2/c^2}$	Pound–Rebka; 33 cm; GPS; muons
Light	c through the medium; $4GM/c^2b$	$1.75''$; Shapiro delay to 10^{-5}
Orbits	Equation (13); $6\pi GM/c^2a(1-e^2)$	Mercury $43''$ per century
Rotation	swirl $2GJ/c^2r^3$	Gravity Probe B; LAGEOS
Waves	transverse, at c	GW170817; tensor; binary pulsar
Scale	$P_0 \approx 4.5 \times 10^{-10}$ Pa	within the heat budget
Galaxies	boundary at $\sqrt{GM/a_0}$	radial acceleration relation

Table 23.2: The equations of the medium and the measurements they meet.

is withdrawn; the medium's stress and the swirl of its flow take its place.

- The lensing formula $\Delta\theta = \int (1/E)(\nabla P_g \cdot \hat{n})\, ds$, whose units do not give an angle, is replaced by ray tracing with Equation (14).
- Three different clock laws ($\sqrt{1 - P_g/P_0}$, $\exp(-P_g/P_0)$, and $\Delta\lambda/\lambda_0 = \sqrt{P_g/P_0}$ for redshift), each with time slowing where the pressure was higher, are replaced by one law, with the active pressure lower near a mass.
- Redshift as energy lost by light to the pressure of the medium is replaced by the slower clocks below.
- The scalar wave equation for the pressure, with a speed that might exceed c, and the predicted dispersion, echoes and non-tensor waves, are withdrawn: the measured speed, polarisation and absence of dispersion rule them out.
- The frame-dragging force $\mathbf{F} = m(\mathbf{v}_g \times \boldsymbol{\omega})$ is replaced by the swirl of the inflow, which gives the measured rate.
- The proposed tests of particle lifetimes underground and at altitude are moved to Part 29: the predicted differences are about one part in 10^{12} per ten kilometres of height, the same as in General Relativity. The proposals for detectors, biological sensor arrays, chronobiology in space and accelerator tests are likewise moved to Part 29, where they are set beside the framework's distinctive predictions.

Clarifications

- The subtitle changes from *Empirical Foundation of Graviton Pressure Theory* to *The Equations of the Medium, and What They Reproduce*.
- A table of quantities, a summary table, a figure of the radial acceleration relation, status marks and this list are added; sources are given for every measurement.

References

B. P. Abbott et al. Gw170814: A three-detector observation of gravitational waves from a binary black hole coalescence. *Physical Review Letters*, 119:141101, 2017a.

B. P. Abbott et al. Gravitational waves and gamma-rays from a binary neutron star merger: Gw170817 and grb 170817a. *The Astrophysical Journal Letters*, 848:L13, 2017b.

J. Bailey et al. Measurements of relativistic time dilatation for positive and negative muons in a circular orbit. *Nature*, 268:301–305, 1977.

Indranil Banik et al. Strong constraints on the gravitational law from gaia dr3 wide binaries. *Monthly Notices of the Royal Astronomical Society*, 527:4573–4615, 2024.

Carlos Barceló, Stefano Liberati, and Matt Visser. Analogue gravity. *Living Reviews in Relativity*, 14:3, 2011.

John S. Bell. How to teach special relativity. In *Speakable and Unspeakable in Quantum Mechanics*. Cambridge University Press, 1987. First published 1976 in Progress in Scientific Culture.

B. Bertotti, L. Iess, and P. Tortora. A test of general relativity using radio links with the cassini spacecraft. *Nature*, 425:374–376, 2003.

Harvey R. Brown. *Physical Relativity: Space-time Structure from a Dynamical Perspective*. Oxford University Press, 2005.

Kyu-Hyun Chae. Breakdown of the newton–einstein standard gravity at low acceleration in internal dynamics of wide binary stars. *The Astrophysical Journal*, 952:128, 2023.

C. W. Chou, D. B. Hume, T. Rosenband, and D. J. Wineland. Optical clocks and relativity. *Science*, 329:1630–1633, 2010.

I. Ciufolini and E. C. Pavlis. A confirmation of the general relativistic prediction of the lense–thirring effect. *Nature*, 431:958–960, 2004.

C. W. F. Everitt et al. Gravity probe b: Final results of a space experiment to test general relativity. *Physical Review Letters*, 106:221101, 2011.

Allvar Gullstrand. Allgemeine lösung des statischen einkörperproblems in der einsteinschen gravitationstheorie. *Arkiv för Matematik, Astronomi och Fysik*, 16(8):1–15, 1922.

Andrew J. S. Hamilton and Jason P. Lisle. The river model of black holes. *American Journal of Physics*, 76:519–532, 2008.

Stacy S. McGaugh, Federico Lelli, and James M. Schombert. Radial acceleration relation in rotationally supported galaxies. *Physical Review Letters*, 117:201101, 2016.

Paul Painlevé. La mécanique classique et la théorie de la relativité. *Comptes Rendus de l'Académie des Sciences*, 173:677–680, 1921.

R. V. Pound and G. A. Rebka. Apparent weight of photons. *Physical Review Letters*, 4:337–341, 1960.

Irwin I. Shapiro. Fourth test of general relativity. *Physical Review Letters*, 13:789–791, 1964.

Pierre Touboul et al. Microscope mission: Final results of the test of the equivalence principle. *Physical Review Letters*, 129:121102, 2022.

J. M. Weisberg and Y. Huang. Relativistic measurements from timing the binary pulsar psr b1913+16. *The Astrophysical Journal*, 829:55, 2016.

Clifford M. Will. The confrontation between general relativity and experiment. *Living Reviews in Relativity*, 17:4, 2014.