Graviton Pressure Theory

The Unified Framework

Individual Submission

This document is part of a multi-part scientific framework

Part 17 of 30

The Definition of Mass

Energy Slowed Down: Matter as Sustained Circulation in the Medium

This submission is part of the broader Graviton Pressure Theory (GPT) project, a redefinition of gravitational interaction rooted in causal field dynamics and physical force transmission. Each document is designed to stand on its own; its full context emerges as part of the larger framework. For the complete series, see https://peopleoftheworld.net/graviton-pressure-theory/.

Author: Shareef Ali Rashada

shareef@peopleoftheworld.net

Second edition, November 2026 (first edition 2025)

Contents

About This Paper 2

Introduction 2

17.1 What Mass Measures 3

17.2 Matter Is Energy Slowed Down 3

17.3 The Numbers of a Particle 5

17.4 The Turnover in Three Dimensions 6

17.5 Inertia, Falling and Weight 9

17.6 What Changes a Body's Mass 9

17.7 Mass in Atoms, Nuclei and Stars 10

17.8 Why Gravity Is Weak 12

17.9 The Numbers to Be Derived 12

17.10 Light, Neutrinos and the Massless 13

17.11 What Remains to Be Derived 14

Conclusion 14

Changes in the Second Edition 14

About This Paper

This paper says what mass is. Physics measures mass with great precision and names its properties, but does not say what it is. GPT proposes that matter is energy slowed down: motion of the medium at the speed of light, wound into a knot so that it goes nowhere, and kept in being by a continual turnover of the medium through it. The paper sets out the evidence that matter is confined motion, the scale at which that motion runs, the turnover that sustains them, why inertia and weight are one exchange, what changes a body's mass and by how much, the numbers of particles, atoms, nuclei and stars, why gravity is weak, and the numbers still to be derived. It marks what is measured, what GPT proposes, and what is still to be derived.

The graviton is in Part 15, the field in Part 16, the equations of the medium in Part 23, and the model as a whole in Part 14.[1][2][3][4] Claims are marked by their status, as the key shows.

How to read the marks. Claims about Graviton Pressure Theory are marked by how far they are established.

Established	No mark. Accepted, measured physics.
Argued	Solid blue underline. A conclusion this framework argues in detail, which readers can check.
Hypothesis	Dashed orange underline. A proposal of GPT, not yet tested.
Open ◦	Dotted grey underline and ◦. A question not yet answered; work still owed.

A whole passage with one status carries a bar of the same style in the left margin. The marks are drawn in distinct line styles, so they read in black and white.

Introduction

Mass is measured more precisely than almost anything in physics. A body's resistance to being pushed and its weight agree, for every material tested, to one part in 10^{15}.[5] A body's mass rises with every joule of energy added to it and falls with every joule taken away, in the ratio $E = mc^2$. These are among the best-established facts there are.

What mass *is* has no answer in the standard account. Newton defined it by the force needed to accelerate it; General Relativity makes it a source of spacetime curvature; the Standard Model gives each elementary particle a number for its coupling to a field that fills all space, measured and not derived. Each describes how mass behaves. None says what is there.

GPT's first edition defined mass as a structure's resistance to the flow of the medium. That instinct is kept: mass is a relation between a pattern and the medium it is made of. This edition says what the pattern is, what the relation is, and why the measured facts follow.

One principle guides it. The universe began hot and energetic, expanded and cooled, and everything in it came from that one beginning. So nothing can be made of anything else: each thing is a later phase or process of what came before, cooled energy settled into some form. The medium is the first such form; light is its travelling wave; matter is its knotted circulation; nuclei, atoms and everything after are made of the knots. "Fundamental" means earliest, not uncaused. How knots form, why they need three dimensions, and how they build nuclei and

[1] Rashada, 2026b.
[2] Rashada, 2026c.
[3] Rashada, 2026d.
[4] Rashada, 2026a.
[5] Touboul et al., 2022.

atoms is the subject of Part 19, *The Pathway to Matter*. This paper is about what confined energy does, and about its numbers.

17.1 What Mass Measures

Mass is measured in three ways, and they always agree. *Inertial* mass is how hard a body is to accelerate. *Passive gravitational* mass is how strongly it is carried toward another mass; torsion balances and the MICROSCOPE satellite find it in the same ratio to inertial mass for every material, to one part in 10^{15}. *Active gravitational* mass is how strongly a body draws other bodies toward it; ranging to the Moon, whose iron core and rocky mantle would pull unequally if active and passive mass differed, finds them equal to about one part in 10^{12}.[6]

Mass is energy. Every change in a body's energy changes its mass by $\Delta m = \Delta E/c^2$:

- **Heat.** Warming a kilogram of water by 100 degrees adds 419 kilojoules and 4.7×10^{-12} kilograms.
- **Binding.** A helium nucleus weighs 0.76 per cent less than the two protons and two neutrons it is made of; the difference is the energy released in binding them.
- **Motion.** A body's energy, and its resistance to further acceleration, rise as $\gamma = 1/\sqrt{1 - v^2/c^2}$. Electrons at the LEP collider reached $\gamma \approx 2 \times 10^5$.
- **Stored energy.** A phone battery charged with 15 watt-hours gains 6×10^{-13} kilograms.
- **Phase changes.** Freezing a kilogram of water releases 334 kilojoules; the ice has less mass by 3.7×10^{-12} kilograms.

The conservation of mass in chemistry is the conservation of energy, at energies too small for $\Delta E/c^2$ to be weighed. Any account of mass must give all of this exactly.

17.2 Matter Is Energy Slowed Down

HYPOTHESIS

Matter is motion of the medium at the speed of light, confined to a small region so that it goes nowhere. Its energy is the energy of that motion; its mass is that energy divided by c^2; and its rest is an average over motion at c.

The mainstream already contains most of the evidence for this, without drawing the conclusion.

A box of light has mass. Light trapped between two mirrors carries energy E but, bouncing back and forth, goes nowhere. The box with the light in it weighs more, by E/c^2, and it is harder to accelerate by the same amount. The reason is mechanical (Figure 17.1). Push the box, and the rear mirror moves into the light arriving at it, which strikes it shifted up in frequency and pushes harder; the front mirror moves away from the light arriving at it, which strikes shifted down and pushes less. The light resists the acceleration, and the resistance comes out exactly $m = E/c^2$. Trapped motion at c behaves as mass, with inertia, from the imbalance of what it delivers to front and back.

[6] Bartlett and Buren, 1986.




Figure 17.1: A mirrored box of light. At rest, the light pushes both mirrors equally. Accelerated, the rear mirror meets the light and is pushed harder than the front, which runs from it: the trapped light resists the acceleration as a mass E/c^2.

The arithmetic of the box. Let the trapped light be two pulses, each of energy $E/2$, running opposite ways. Move the box at speed v, with $\beta = v/c$. The pulse running forward is shifted up in frequency by the factor $k = \sqrt{(1+\beta)/(1-\beta)}$, and the pulse running back is shifted down by $1/k$. Their total energy and momentum are then

$$E_{\text{box}} = \frac{E}{2}\left(k + \frac{1}{k}\right) = \gamma E, \qquad p_{\text{box}} = \frac{E}{2c}\left(k - \frac{1}{k}\right) = \gamma \frac{E}{c^2} v, \tag{1}$$

with $\gamma = 1/\sqrt{1-\beta^2}$. That is exactly the energy and momentum of a body of mass $m = E/c^2$ moving at v. Nothing was assumed about mass: the light had none. The factor k is the same one that sets the leading and trailing exchanges of a moving body in GPT (Part 23): a moving body's momentum is the imbalance between its front and back. Pushing on Equation (1) gives inertia its measured shape at high speed: a body resists a push along its motion as $\gamma^3 m$ and across it as γm.

The proton is almost all confined motion. The three quarks in a proton account, by their own masses, for about one per cent of its mass. The rest is the energy of the motion and binding confined inside it.[7] A calculation from the theory of the strong force divides the proton's mass roughly into a tenth from the quarks' own masses and their interaction with the vacuum, about a third from the quarks' motion, about a third from the energy of the field that binds them, and about a quarter from a quantum effect of that field.[8] Most of the mass of everything that can be touched is confined energy.

The electron moves at c. Dirac's equation for the electron, one of the best-tested equations in physics, says that at every instant the electron's velocity is $\pm c$. What is seen as its motion is an average over a rapid trembling, which Schrödinger named *Zitterbewegung*: about 10^{21} times a second over a distance near the electron's Compton wavelength.[9] It is too fast to observe in a free electron, but the same equation has been reproduced in trapped ions, and the trembling seen.[10]

[7] Yang et al., 2018.
[8] Yang et al., 2018.
[9] Schrödinger, 1930.
[10] Gerritsma et al., 2010.

An electron cannot be made smaller. Trying to confine an electron within less than its Compton wavelength, 2.4×10^{-12} metres, does not produce a smaller electron. It produces new pairs of electrons and positrons.

Matter and light turn into each other. An electron and a positron that meet become two flashes of light. Light with enough energy, passing a nucleus, becomes an electron and a positron;[11] and in collisions of light with light alone, pairs of matter have been made from nothing but light.[12] Matter is made of the same thing light is made of, in another form. In GPT that thing is the medium: light is the medium's travelling wave, and matter is its confined circulation.

17.3 The Numbers of a Particle

If an electron is circulation at c carrying its spin, $\hbar/2$, its momentum $m_e c$ goes round a loop of radius

$$r = \frac{\hbar}{2mc}, \tag{2}$$

and every other number follows from its mass. Table 17.1 gives them for the elementary particles, the electron, muon and tau, from Equation (2), and for two composite particles, the pion and the proton, from their measured sizes.

	Mass (MeV)	**Loop radius** (m)	**Turns** (per second)	**Acceleration** (m/s^2)	**Force to hold it** (N)
Electron	0.511	1.9×10^{-13}	2.5×10^{20}	4.7×10^{29}	0.42
Muon	105.7	9.3×10^{-16}	5.1×10^{22}	9.6×10^{31}	1.8×10^{4}
Tau	1777	5.6×10^{-17}	8.6×10^{23}	1.6×10^{33}	5.1×10^{6}
Pion (measured size)	139.6	6.6×10^{-16}	7.2×10^{22}	1.4×10^{32}	3.4×10^{4}
Proton (measured size)	938.3	8.4×10^{-16}	5.7×10^{22}	1.1×10^{32}	1.8×10^{5}

Table 17.1: What a particle is doing while it appears to rest, if its energy circulates at c: loop radius, turns per second $c/2\pi r$, acceleration c^2/r, and the force mc^2/r needed to keep its momentum on the loop. HYPOTHESIS

Dirac's equation already contains the loop. The trembling of Dirac's electron runs at $2m_e c^2/\hbar \approx 1.55 \times 10^{21}$ radians a second, which is 2.5×10^{20} cycles a second, over an amplitude of $\hbar/2m_e c$. Those are exactly the turns per second and the loop radius of Equation (2). A loop at c carrying the electron's spin and the trembling Dirac found in his equation are the same motion, described twice.

The magnetic moment. The electron's magnetic moment, measured to thirteen figures, is $g = 2.00231930436$ times what its spin would give if its charge circulated like its mass. A simple charged loop at c carrying spin $\hbar/2$ gives $g = 1$; models in which the energy is shared between the circulation and the field around it give $g = 2$.[13] The remaining 0.00116 is, to within 0.15 per cent, $\alpha/2\pi$, where $\alpha \approx 1/137$ is the strength of the electron's coupling to light:

[11] Anderson, 1933.
[12] STAR Collaboration, 2021.
[13] Hestenes, 1990.




Figure 17.2: Accelerations, from the Earth's surface to the circulation inside matter, on a logarithmic scale. The circulation figures assume matter's energy runs at c around a loop of the particle's size. HYPOTHESIS

the first correction for the electron's interaction with its own field. In GPT, that is the knot's interaction with its own counterflow. Deriving the 2, and then the correction, is the test of the full pattern. ◦

"Rest" is the most extreme motion there is, wrapped so that it looks still (Figure 17.2). That is the deepest form of GPT's premise that nothing is at rest. It has three consequences already met in this framework.

- **Clocks and rods.** A moving body's circulation must still run at c through the medium, so each of its cycles takes longer and the pattern settles squashed along its motion: the slowing of clocks and the squashing of rods, by the same factor (Part 23).
- **Phase.** The phase of matter is the phase of a rhythm of 10^{20} or more a second. That is why the medium's interactions are so fine-grained, and why the pattern a body writes into its counterflow is regular enough to build shells (Part 16).
- **The electron still looks like a point.** Collisions find no size in the electron down to about 10^{-18} metres. So the loop is the *path* of something much smaller moving at c, not a ring spread out over 10^{-13} metres.

17.4 The Turnover in Three Dimensions

HYPOTHESIS

Matter is kept in being by a cycle that runs through it and out into space (Figure 17.3).

1. **Arrival.** In-phase gravitons arrive at a knot from every direction, pressing. Around a lone knot the arrivals balance, and there is no net push. But the knot continually turns pressing medium into neutral medium, so the region around it holds slightly fewer pressing gravitons than the region far away. That deficit is a pressure gradient, and the medium drifts gently inward down it: the drift is the field, $v^2 = 2GM/r$.
2. **Transfer.** A graviton reaching the knot meets its phase boundary and is drawn into step with the circulation, as one oscillator locks onto a stronger one nearby. The self-repulsive energy it had from being in step with its surroundings goes into the circulation.




Figure 17.3: The turnover. In-phase medium is pressed in; its pressing energy goes into the circulation of the body's knots; it leaves neutral, carrying the knots' phase and the energy as motion; far away it is re-phased and available to flow in again. HYPOTHESIS

3. **Departure.** It leaves carrying the knot's phase: neutral to the surrounding pressure, but moving and carrying momentum. It travels out through the rest of the body, past its edges and into space, as the counterflow, carrying the pattern of the body's spin and motion (Part 16).

4. **Return.** Far out, the surrounding medium brings the neutral gravitons back into step. That re-phasing is the relaxation boundary (Part 15). In phase again, they press again, and are available to flow in again.

Matter is a turnover, not a lump, and matter and the medium are partners in one circulation. The pattern lasts while the medium passing through it changes, as a whirlpool or a flame keeps its shape while water or fuel passes through.

The first edition said that mass is "not a frozen property": that "each moment of graviton interaction reaffirms" it, and that mass is "the stabilized echo of resistance in a perpetually refreshed field". It also said that gravitons are absorbed, depositing energy. Both are kept, and joined: the deposit is what keeps the pattern in being.

Where the energy goes. In a steady state a knot does not grow, so the energy it takes in must leave. It leaves as the motion of the neutral gravitons: pressing energy in, energy of motion out, with the knot as the converter. Far away, re-phasing turns that motion back into pressing energy, closing the loop of energy as well as of flow. Whatever escapes the loop as heat is bounded by the Earth's heat budget, which is how the medium's density was fixed (Part 23); whatever stays is accretion, at most about two kilograms a year for the whole Earth (Part 16).

Why every electron is the same. Every electron has the same mass, and so does every proton. On a picture in which each particle carries a stored amount from its birth, that is a coincidence of origin. On the turnover it follows: a knot's energy is the balance between what it takes in and what it sheds for its winding, as every whirlpool in a given current settles to the same size. Mass is an equilibrium set by the medium, not an amount each particle carries. That gives a test. Where the medium's state differed, masses would differ. The ratio of the proton's mass to the electron's was the same seven billion years ago, in light absorbed by molecules in a distant galaxy, to about one part in 10^7,[14] so the medium's state, as knots feel it, has been steady at least that long.

Why a body's draw follows its energy. A knot takes in a fixed amount of the medium with each turn of its circulation. It cannot be area that sets the intake: a heavier particle is a smaller knot, its size falling as one over its mass. But its turns per second are its energy divided by h, so a fixed take per turn makes its intake proportional to its energy. In a body of many knots every circulation contributes; binding lowers the circulating energy, and the intake falls with it. So a body's draw on the medium, its active gravitational mass, follows its total energy, binding included, which is what the Moon's ranging shows, and it puts the relaxation boundary at one acceleration, a_0, around every mass (Part 15). The size of the take per turn is owed. ◦

Why matter is nearly transparent. If each knot took most of what reached it, the outer knots of a body would starve the inner ones, and mass would depend on shape, as in Le Sage's shadowing theory. Mass adds exactly, and gravitational shielding is bounded far below what such shadowing needs. So each knot takes only a tiny share of what passes, and the medium flows through matter almost freely, as neutrinos pass through the Earth. Every knot in a body receives its share, and the body's total is their sum.

What makes a particle an entity. A knot's material is replaced continually, so its identity cannot lie in its material. It lies in what the turnover cannot change: its winding (its charge and its baryon or lepton number), its handedness, its spin and its balance energy. These are exactly the quantities conserved in every interaction. A particle persists as a whirlpool does: as the invariants of a pattern the flow passes through.

Holding and supplying are different jobs. The medium's free-space pressure, $P_0 \approx 4.5 \times 10^{-10}$ pascals, acting over an electron-sized area, gives a force of about 10^{-35} newtons, some 10^{34} times too small to hold an electron's circulation on its loop. So the surrounding pressure does not hold a particle together. The knot holds itself, by the phase boundary it makes and the stiffness of the in-phase medium against it, and its strands are kept apart by their common phase (Part 19); the inflow supplies the turnover. Kelvin proposed in 1867 that atoms are knotted vortices of a medium;[15] his version failed on the spectra of atoms, but stability from knotting survived, in Skyrme's particles and their successors.[16]

A second role for the counterflow. The departing gravitons are neutral to the surrounding pressure, but they carry the knot's phase. Passing through other knots, they meet a phase that is in step or out of step with it. If charge is the phase a knot emits (Part 19), the counterflow is

[14] Bagdonaite et al., 2013.
[15] Thomson, 1867.
[16] Skyrme, 1962.

gravitationally neutral and electrically active at once, and in a neutral atom equal and opposite phases cancel at a distance.

17.5 Inertia, Falling and Weight

HYPOTHESIS

Each of the three masses has its interface.

- **Inertial mass** is the confined circulation. Pushed, the circulation's turning is unbalanced front to back, as in the box of light, where what turns it is the knot's own boundary in place of mirrors. A moving body's imbalance is its momentum, and it costs nothing to keep (Part 23); the resistance to changing it is inertia, and its size is the circulating energy.
- **Passive gravitational mass**: **a pattern of the medium is carried by the medium.** Matter is circulation of the medium, so when the medium accelerates toward a mass, the pattern goes with it. Every body falls alike because every body is made of the medium it falls in.
- **Active gravitational mass** is the body's draw on the medium: a fixed take per turn, so proportional to its energy (Section 17.4).

Weight is the support's relay of the push needed to keep the pattern from being carried (Part 11).

Inertial and passive gravitational mass agree for every material because both follow from what matter is: circulation of the medium. Active mass agrees with them because the take per turn makes a body's draw follow the same energy. The first edition held that inertia and gravity are "resistance within the graviton field". What changes is what the resistance is: not a static impedance, but the circulation of the pattern and its turnover, whose size is the pattern's energy.

The exchange also gives inertia its measured shape at high speed. Along the motion a body resists further acceleration as $\gamma^3 m$, across it as γm: the longitudinal and transverse masses of relativistic mechanics, confirmed in every particle accelerator.

Motion is a relation with the medium. The first edition said there is no universal tendency to continue motion. That is withdrawn: spacecraft coast for decades. But the instinct behind it is kept. A body's motion is not a property it has in isolation; it is its relation with the medium, and it continues because of that relation, not in spite of it.

17.6 What Changes a Body's Mass

The first edition listed what changes mass, and the direction of every change it gave is right. What it lacked was the size. In this edition each change has the size measurement requires: $\Delta E/c^2$, no more (Table 17.2).

Change	First edition	Measured
Heating	Raises mass	Raises it by $\Delta E/c^2$: $+4.7 \times 10^{-12}$ kg for a kilogram of water warmed by 100 degrees
Binding	Reduces mass	Reduces it by the binding energy$/c^2$: 0.76% for helium
Stress	Raises mass	Raises it by the stored elastic energy$/c^2$
Speed	Raises resistance	Raises it as γ (across) and γ^3 (along)
Melting	Raises mass	Raises it by the latent heat$/c^2$: $+3.7 \times 10^{-12}$ kg per kilogram of ice
Depth in a field	Raises resistance	Unchanged locally; seen from far away, a body lowered toward a mass has less energy, by GMm/r: 7×10^{-10} of its mass at the Earth's surface

Table 17.2: What changes a body's mass. The first edition's directions are right except for depth in a field; the sizes are $\Delta E/c^2$.

Coherence and structure. For materials as they are, mass tracks energy, binding included, to about one part in 10^{12}, because materials whose nuclei hold different fractions of binding energy fall alike to one part in 10^{15}. A material's order changes its mass by the energy the order holds, and no more. The first edition's "equal matter, different mass" is restated: equal *energy*, equal mass; and the structure of a body decides where in it the medium's delivery lands, not its total (Part 15).

Mass from surroundings: a real effect at another level. The first edition's environmental mass has a true counterpart. An electron moving through a crystal responds to a push as if it had a different mass, because it moves in constant exchange with the lattice around it: in gallium arsenide about 0.067 of its free mass, and in some metals hundreds of times its free mass.[17] The electron's own mass does not change; its *effective* mass is set by its interaction with its surroundings. That is inertia arising from exchange with a surrounding medium, measured in the laboratory every day. GPT proposes that a body's own mass arises the same way, from its exchange with the medium of space.

Changing mass beyond $\Delta E/c^2$. The first edition proposed mass-tuning devices. Changing mass by more than $\Delta E/c^2$ is not excluded by measurement in every form, but it is bounded: spinning rotors weigh normally.[18] Whether a body changing its order weighs differently while it changes is open, and an experiment is under way to weigh the energy released when a material becomes superconducting. ◦[19] It is a test for Part 29.

17.7 Mass in Atoms, Nuclei and Stars

The atom. A hydrogen atom weighs less than a free proton and a free electron by its binding energy, 13.6 electronvolts: 1.45×10^{-8} of its mass. Its numbers fit the circulation picture closely. The electron in hydrogen's lowest state moves at αc, about 2,190 kilometres a second, so the

[17] Stewart, 1984.
[18] Faller et al., 1990.
[19] Calloni et al., 2014.




Figure 17.4: Binding energy per nucleon. Measured values and the smooth part of the liquid-drop formula (volume, surface, charge and balance terms, with standard coefficients).

electron moves round the atom at 1/137 of the speed at which its own energy circulates. And the atom's radius, 5.29×10^{-11} metres, is the electron's circulation scale, $\hbar/m_e c$, divided by α: the atom is 137 times larger than the electron's own loop. The number $\alpha \approx 1/137.036$ is the strength with which one charged pattern couples to another; why it has that value is unexplained in every theory, and is owed. ◦

Nuclei. Nuclei weigh less than their parts by much more: about 8 MeV for each nucleon, nearly one per cent of its mass. The binding per nucleon rises from 1.1 MeV in the deuteron to a peak of 8.8 MeV at iron and nickel, and falls slowly to 7.6 MeV in uranium (Figure 17.4). Joining light nuclei, up to iron, releases energy; splitting heavy ones does too. A formula with five terms, the liquid-drop formula, fits the binding of the heavier nuclei to about one per cent, and in GPT each of its terms has a mechanism (Table 17.3; Part 19).

Term	Form	Size (MeV)	In GPT
Volume	$+a_V A$	15.8	Each nucleon's outer pattern overlaps its neighbours' and reinforces between them, binding them; each has only so many neighbours, so binding grows with the number of nucleons
Surface	$-a_S A^{2/3}$	17.8	Nucleons at the surface have fewer neighbours to interlock with
Charge	$-a_C Z^2/A^{1/3}$	0.71	Protons emit in phase with one another and push apart
Balance	$-a_A(N-Z)^2/A$	23.7	A proton and a neutron, different knots, can bind with parallel spins; two of a kind cannot, since identical knots of the same circulation push each other out, so equal numbers bind best
Pairing	$\pm a_P/A^{1/2}$	11.2	Two identical nucleons of opposite circulation do not push each other out and can share a pattern

Table 17.3: The liquid-drop formula for the binding energy of a nucleus with A nucleons, Z protons and N neutrons. Its coefficients are fitted to measurement; in GPT each term has a mechanism. HYPOTHESIS

Stars. Fusing four hydrogen nuclei into helium releases 0.7 per cent of their mass; splitting a uranium nucleus releases about 0.09 per cent. The Sun shines by the first: it turns about 4.3 million tonnes of mass into light every second, and in its 4.6 billion years has lost about three ten-thousandths of its mass that way.

17.8 Why Gravity Is Weak

Between two protons, the electric push is 1.2×10^{36} times the gravitational pull; between two electrons, 4.2×10^{42} times. Put another way, gravity's strength between two protons, $Gm_p^2/\hbar c$, is 5.9×10^{-39}, against $\alpha \approx 7.3 \times 10^{-3}$ for the electric coupling. No theory explains the gap; it is known as the hierarchy problem.

GPT gives the direction of the answer. A body's gravity is its draw on the medium, and the draw is a tiny take per turn (Section 17.4): matter is nearly transparent to the medium, and must be, or bodies would shade one another. A charged pattern's effect on another is its emitted phase acting directly on the other's circulation. Gravity is weak because each knot takes so little of what passes through it; charge is strong because phase acts on phase directly. The size of the gap, 10^{36}, should follow from the size of the take per turn and the strength of phase coupling; both are owed. ◦

17.9 The Numbers to Be Derived

These are the numbers a theory of mass is asked for (Table 17.4). No theory derives any of them. In the Standard Model the masses of the electron, muon and tau and the strength of charge are measured inputs: the electron's mass enters through a coupling to the Higgs field of 2.9×10^{-6}, itself measured. The theory of the strong force computes the proton's mass, to about

one per cent, from the quarks' masses and the strong coupling, which are again measured. GPT does not yet derive them either; Table 17.4 is its target.

Quantity	Measured	In GPT, it should follow from
Electron's mass	0.51099895 MeV	The threshold of the simplest knot
Proton-to-electron mass ratio	1836.15267	The three-lobed knot against the single one
Neutron minus proton	1.29333 MeV	The difference of one lobe's twist
Muon and tau to electron	206.768 and 3477.2	Excited windings of the electron's knot
Strength of charge, $1/\alpha$	137.035999	The coupling of phase to phase
Electric to gravitational, two protons	1.24×10^{36}	Phase coupling against the take per turn
Electron's magnetic moment, g	2.00231930436	The knot with its field
Neutrino masses	below 0.45 eV; differences squared 7.4×10^{-5} and 2.5×10^{-3} eV^2	Patterns holding very little energy

Table 17.4: The numbers of mass. HYPOTHESIS

Regularities that any account must explain. The three charged leptons satisfy Koide's relation, $(m_e + m_\mu + m_\tau)/(\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2 = 0.666661$, against $2/3$, to within the uncertainty of the tau's mass. No one knows why. A weaker curiosity, noted in 1951, is that $6\pi^5 = 1836.118$, within two parts in 10^5 of the proton-to-electron ratio; it may mean nothing. If the masses are thresholds and excited windings of knots, relations of this kind are what the geometry of the knots should produce.

17.10 Light, Neutrinos and the Massless

Light is the medium's travelling wave; matter is its confined circulation. A travelling wave has only a front: there is no back for it to deliver to, and no frame in which it is at rest. Its energy and momentum are related by $E = pc$, and its mass is zero. Matter has a front and a back, a frame in which it is at rest, and mass E/c^2. The first edition said photons are "fully aligned" with the flow and have "zero impedance"; that is restated: light is not confined.

Neutrinos have mass. They change from one type to another as they travel, which they could not do without mass,[20] and direct measurement puts the mass of the electron's neutrino below about half an electronvolt, less than a millionth of the electron's.[21] The differences between the squares of the three masses, measured from the changes, are 7.4×10^{-5} and 2.5×10^{-3} eV^2, so the heaviest neutrino is at least 0.05 eV. What makes a neutrino's pattern carry so little energy is open. ◦

[20] Fukuda et al., 1998.
[21] KATRIN Collaboration, 2025.

Particle	In GPT	Measured
Photon	The medium's travelling wave: not confined	No mass; $E = pc$
Neutrino	A confined pattern holding very little energy	Mass below about 0.45 eV; three types that change into one another
Electron	A single knot of circulation at c, one handedness	0.511 MeV; charge -1; looks like a point to 10^{-18} m
Proton	A composite knot; almost all confined motion	938 MeV; charge $+1$; radius 0.84 fm

Table 17.5: Particles as patterns of the medium. HYPOTHESIS

17.11 What Remains to Be Derived

- The numbers of Table 17.4, from the medium's non-linear response, which sets the thresholds at which knots form (Part 19). ◦
- The size of the take per turn, and the re-phasing rate that closes the loop. ◦
- The electron's magnetic moment, $g = 2$ and its correction, from the full pattern and its field. ◦
- The coefficients of the liquid-drop formula from the overlap of lobes. ◦
- Whether a body changing its order weighs differently while it changes. ◦

Conclusion

Mass is energy, and matter is energy slowed down: motion of the medium at the speed of light, confined so that it goes nowhere. Light trapped in a box has exactly the energy and momentum of a mass E/c^2; the proton is almost all confined motion; Dirac's electron already contains a loop at c of radius $\hbar/2mc$, turning 2.5×10^{20} times a second. The medium pressed into matter gives up its in-phase energy to the circulation and leaves as the neutral counterflow, to be re-phased far away and return: that turnover keeps matter in being, its balance gives every electron the same mass, and the inflow that supplies it is gravity. A pattern of the medium is carried by the medium, which is why every body falls alike; a fixed take per turn makes a body's draw follow its energy, and the smallness of that take is why gravity is 10^{36} times weaker than charge. Every change of energy changes mass by $\Delta E/c^2$, from a charged battery to the binding of nuclei and the light of the Sun. The numbers of mass, the electron's 0.511 MeV, the ratio 1836, the strength $1/137$, are derived by no theory yet; they are GPT's target.

Changes in the Second Edition

This list records what changed from the first edition and why. Changes to the argument alter what the paper claims or how it supports a claim; clarifications improve wording, sources or presentation and leave the argument as it was.

Changes to the argument

- Mass is defined as the energy of a confined circulation of the medium at c, divided by c^2, in place of an impedance profile $\rho_{\text{imp}} = f(\text{coherence, geometry, phase})$. The evidence that matter is confined motion is added: the box of light, the proton's mass, Dirac's electron, pairs from confinement, matter made from light.
- The scale of that motion is set out: the electron's loop, frequency, acceleration and force, and the conclusion that rest is the most extreme motion there is.
- Inertia's measured shape at high speed (γ across, γ^3 along) is given. "No universal tendency to continue motion" is withdrawn; motion is kept as a relation with the medium.
- The effects of heat, binding, stress, speed and melting on mass are kept, with their directions, and given their measured size, $\Delta E/c^2$. Depth in a field is corrected: locally unchanged, reduced as seen from far away.
- "Equal matter, different mass" and mass-tuning devices are restated: mass tracks energy to about 10^{-12}; changes beyond $\Delta E/c^2$ are bounded and open. Effective mass in crystals is added as a measured case of inertia from exchange with a surrounding medium.
- Photons are restated as the medium's unconfined travelling wave. Neutrinos are given their measured mass. The classification table is replaced.
- The turnover is added, joining the first edition's "perpetually refreshed" mass and its absorption of gravitons, and set out as a full cycle in three dimensions, with its return through re-phasing, its energy route, every electron's identical mass as a balance (tested by the constancy of the proton-to-electron mass ratio), a fixed take per turn giving a body's draw in proportion to its energy, the transparency of matter, and identity as the invariants of the pattern.
- Inertial, passive and active gravitational mass are separated. Passive mass follows from matter being made of the medium it falls in; an earlier wording of this edition, that a body is carried at a rate set by its own intake, is corrected.
- New: the numbers of mass. The arithmetic of the box of light, giving γE and $\gamma(E/c^2)v$; the proton's mass budget; a table of loop radius, turns, acceleration and holding force for five particles; the match between the loop and Dirac's trembling; the magnetic moment and its correction; mass in the hydrogen atom and the meaning of $1/137$ as the ratio of the electron's orbital speed to its circulation; nuclear binding, with each term of the liquid-drop formula given a mechanism; mass and the stars; why gravity is weak; and a table of the numbers still to be derived, with Koide's relation.
- The formation of knots, their three dimensions, charge, the fragments of forced dissolution, and the path from knots to atoms, drafted for this paper, are moved to Part 19, *The Pathway to Matter*. Mass is how much; matter is how it is arranged.
- The energy equation $E = \int \Delta P_g \, dV$ and the reinterpretation of redshift through variable mass are withdrawn.

Clarifications

- Table 17.3 (11 November): the volume, balance and pairing terms are reworded to the two meanings of phase (Part 15): overlapping patterns reinforce in spatial phase; identical knots of the same circulation push each other out.

- The subtitle changes from *Impedance, Coherence, and the Graviton Field* to *Energy Slowed Down: Matter as Sustained Circulation in the Medium*.
- "Massless, self-repulsive" gravitons are described as in Part 15. References to corridors and to "reflection" of gravitons are replaced by the inflow, the counterflow and the turnover.
- Four figures, five tables, status marks and this list are added; the principle that everything is a later phase of what came before is stated in the introduction.

References

Carl D. Anderson. The positive electron. *Physical Review*, 43:491–494, 1933.

J. Bagdonaite et al. A stringent limit on a drifting proton-to-electron mass ratio from alcohol in the early universe. *Science*, 339:46–48, 2013.

D. F. Bartlett and D. Van Buren. Equivalence of active and passive gravitational mass using the Moon. *Physical Review Letters*, 57:21–24, 1986.

Enrico Calloni et al. Towards weighing the condensation energy to ascertain the Archimedes force of vacuum. *Physical Review D*, 90:022002, 2014.

J. E. Faller, W. J. Hollander, P. G. Nelson, and M. P. McHugh. Gyroscope-weighing experiment with a null result. *Physical Review Letters*, 64:825–826, 1990.

Y. Fukuda et al. Evidence for oscillation of atmospheric neutrinos. *Physical Review Letters*, 81: 1562–1567, 1998.

R. Gerritsma et al. Quantum simulation of the Dirac equation. *Nature*, 463:68–71, 2010.

David Hestenes. The zitterbewegung interpretation of quantum mechanics. *Foundations of Physics*, 20:1213–1232, 1990.

KATRIN Collaboration. Direct neutrino-mass measurement based on 259 days of KATRIN data. *Science*, 388:180–185, 2025.

Shareef Ali Rashada. From revelation to foundation (part 14, second edition). https://peopleoftheworld.net/graviton-pressure-theory/full-framework/part-14-from-revelation-to-foundation/, 2026a.

Shareef Ali Rashada. Gravitons (part 15, second edition). https://peopleoftheworld.net/graviton-pressure-theory/full-framework/part-15-gravitons/, 2026b.

Shareef Ali Rashada. The properties of gravitational fields (part 16, second edition). https://peopleoftheworld.net/graviton-pressure-theory/full-framework/part-16-the-properties-of-gravitational-fields/, 2026c.

Shareef Ali Rashada. The mathematical foundation (part 23, second edition). https://peopleoftheworld.net/graviton-pressure-theory/full-framework/part-23-the-mathematical-foundation/, 2026d.

E. Schrödinger. Über die kräftefreie bewegung in der relativistischen quantenmechanik. *Sitzungsberichte der Preussischen Akademie der Wissenschaften, Physikalisch-mathematische Klasse*, 24:418–428, 1930.

T. H. R. Skyrme. A unified field theory of mesons and baryons. *Nuclear Physics*, 31:556–569, 1962.

STAR Collaboration. Measurement of e^+e^- momentum and angular distributions from linearly polarized photon collisions. *Physical Review Letters*, 127:052302, 2021.

G. R. Stewart. Heavy-fermion systems. *Reviews of Modern Physics*, 56:755–787, 1984.

William Thomson. On vortex atoms. *Proceedings of the Royal Society of Edinburgh*, 6:94–105, 1867.

Pierre Touboul et al. Microscope mission: Final results of the test of the equivalence principle. *Physical Review Letters*, 129:121102, 2022.

Yi-Bo Yang et al. Proton mass decomposition from the QCD energy momentum tensor. *Physical Review Letters*, 121:212001, 2018.