

Formalization of the Proof of Fodor's Lemma

Theofanis Chatzidiamantis Christoforidis

January 2024

Basic Definitions

Let α be a limit ordinal and $C \subseteq \alpha$ a set.

Definition

- ▶ C is called **unbounded** in α if $\sup(C \cap \alpha) = \alpha$.

Basic Definitions

Let o be a limit ordinal and $C \subseteq o$ a set.

Definition

- ▶ C is called **unbounded** in o if $\sup(C \cap o) = o$.
- ▶ C is called **closed** in o if for every $\alpha < o$, if $C \cap \alpha \neq \emptyset$, then $\sup(C \cap \alpha) \in C$.

Basic Definitions

Let o be a limit ordinal and $C \subseteq o$ a set.

Definition

- ▶ C is called **unbounded** in o if $\sup(C \cap o) = o$.
- ▶ C is called **closed** in o if for every $\alpha < o$, if $C \cap \alpha \neq \emptyset$, then $\sup(C \cap \alpha) \in C$.
- ▶ C is called **club** in o if it is closed and unbounded in o .

Basic Definitions

Let o be a limit ordinal and $C \subseteq o$ a set.

Definition

- ▶ C is called **unbounded** in o if $\sup(C \cap o) = o$.
- ▶ C is called **closed** in o if for every $\alpha < o$, if $C \cap \alpha \neq \emptyset$, then $\sup(C \cap \alpha) \in C$.
- ▶ C is called **club** in o if it is closed and unbounded in o .

Remark

Equivalently, C is unbounded in o if for every $\alpha < o$, there is a $\beta \in C$ such that $\alpha < \beta$.

Basic Definitions

Definition

Let α be a limit ordinal and $S \subseteq \alpha$ a set. S is called **stationary** in α if for every club set $C \subseteq \alpha$, $S \cap C \neq \emptyset$.

Basic Definitions

Definition

Let α be a limit ordinal and $S \subseteq \alpha$ a set. S is called **stationary** in α if for every club set $C \subseteq \alpha$, $S \cap C \neq \emptyset$.

Definition

Let $f : C \rightarrow D$ be a function, where C and D are sets of ordinals. f is called **regressive** if for every $0 < \alpha \in C$, $f(\alpha) < \alpha$.

Basic Definitions

Definition

Let κ be a cardinal.

- ▶ The **cofinality** $\text{cof}(\kappa)$ of κ is the smallest possible cardinality of an unbounded set in κ

Basic Definitions

Definition

Let κ be a cardinal.

- ▶ The **cofinality** $\text{cof}(\kappa)$ of κ is the smallest possible cardinality of an unbounded set in κ
- ▶ A cardinal is **regular** if $\text{cof}(\kappa) = \kappa$

Basic Definitions

Definition

Let κ be a cardinal.

- ▶ The **cofinality** $\text{cof}(\kappa)$ of κ is the smallest possible cardinality of an unbounded set in κ
- ▶ A cardinal is **regular** if $\text{cof}(\kappa) = \kappa$

Remark

Cofinality generalizes to ordinals and this is what Lean actually does.

The Theorem

Our goal is to prove the following:

Theorem (Fodor's Lemma)

Let κ be an uncountable regular cardinal, $S \subseteq \kappa$ a stationary set and let $f : S \rightarrow \kappa$ be a regressive function. Then there is an ordinal $\theta < \kappa$ such that $f^{-1}(\{\theta\})$ is stationary in κ .

Sets and Ordinals are different types.

- ▶ In set theory, everything is a set, including cardinals and ordinals.
This is not true in the type-theoretic framework we are working with.

Sets and Ordinals are different types.

- ▶ In set theory, everything is a set, including cardinals and ordinals. This is not true in the type-theoretic framework we are working with.
- ▶ With this in mind, our main objects of study here are of type ***Set*** ***Ordinal***.

Sets and Ordinals are different types.

- ▶ In set theory, everything is a set, including cardinals and ordinals. This is not true in the type-theoretic framework we are working with.
- ▶ With this in mind, our main objects of study here are of type ***Set*** ***Ordinal***.
- ▶ This makes sense: recall that in ZFC, an ordinal contains every ordinal strictly smaller than itself.

Sets and Ordinals are different types.

In Lean, for $C : \text{Set } \text{Ordinal}$ and $o : \text{Ordinal}$, we define:

► $\text{strict_Ordinal_res } C \ o := \{c \in C \mid c < o\}$, corresponding to $C \cap o$.

Sets and Ordinals are different types.

In Lean, for $C : \text{Set } \text{Ordinal}$ and $o : \text{Ordinal}$, we define:

- ▶ $\text{strict_Ordinal_res } C \ o := \{c \in C \mid c < o\}$, corresponding to $C \cap o$.
- ▶ $\text{Ordinal_res } C \ o := \{c \in C \mid c \leq o\}$

Sets and Ordinals are different types.

In Lean, for $C : \text{Set } \text{Ordinal}$ and $o : \text{Ordinal}$, we define:

- ▶ $\text{strict_Ordinal_res } C \ o := \{c \in C \mid c < o\}$, corresponding to $C \cap o$.
- ▶ $\text{Ordinal_res } C \ o := \{c \in C \mid c \leq o\}$

"strict" is the correct notion, but Ordinal_res has nicer properties and we use it whenever they are equal (i.e., when $o \notin C$)

We need one last definition :

We need one last definition :

Definition

Let κ be a cardinal and let $(C_\alpha)_{\alpha < \kappa}$ be a sequence of subsets of κ .
The **diagonal intersection** of $(C_\alpha)_{\alpha < \kappa}$ is defined to be

$$\Delta_{\alpha < \kappa} C_\alpha := \{\beta < \kappa \mid \beta \in C_\theta \ \forall \theta < \beta\}$$

General Structure:

Basic properties of supremums + Definitions + regularity of κ



The intersection of two club sets is club



The intersection of less than κ club sets is club



The diagonal intersection of κ club sets is club



Fodor's Lemma

Proving Closedness:

Proving Closedness:

This came down to manipulating supremums.

Proving Unboundedness:

For each of the main theorems about club sets, we construct a "good" increasing sequence $\mathbb{N} \rightarrow \textit{Ordinal}$ recursively.

Proving Unboundedness:

For each of the main theorems about club sets, we construct a "good" increasing sequence $\mathbb{N} \rightarrow \text{Ordinal}$ recursively.

We then use our closedness assumptions to prove that its limit is in our intersection.

These arguments use both closedness and unboundedness assumptions.

On Cofinality

Why is it useful to work with sequences of length ω ?

On Cofinality

Why is it useful to work with sequences of length ω ?

If κ is uncountable and regular, the limit of such an ascending sequence will always be less than κ by regularity of κ , since \mathbb{N} is countable.

On Cofinality

Why is it useful to work with sequences of length ω ?

If κ is uncountable and regular, the limit of such an ascending sequence will always be less than κ by regularity of κ , since \mathbb{N} is countable.

With this in mind, a very easy generalisation for the properties of club sets is apparent:

Remark

We can replace our uncountable regular cardinal with any ordinal α satisfying $\text{cof}(\alpha) > \aleph_0$.