

# Riemann Hypothesis

## A Computational Research Framework · Mathematics/Number Theory

Lois-Kleinner Alpasan · Anticloud FZ LLE, UAE Free Zone · research@0-1.gg

PAX L5 Narrow L2 General 27B · 2026-10-02

### ABSTRACT

The central conjecture posits that there exists a self-adjoint operator  $H$  acting on a specific Hilbert space  $H_{\text{arith}}$  such that:  $\text{Spec}(H) = \{ \gamma_n : \zeta(1/2 + i\gamma_n) = 0 \}$  where  $\gamma_n$  are the imaginary parts of the non-trivial zeros of the Riemann zeta function  $\zeta(s)$ . **Refinement:** Unlike previous failed attempts that relied on ad-hoc Hamiltonians, this proposal hypothesizes that  $H$  arises naturally from the **quantum geometry of the adèle ring**  $A_{\mathbb{Q}}$ , specifically through the action of the Laplacian on the space of automorphic forms over the field of rational numbers, regularized via the **Selberg trace formula** for the modular group  $SL_2(\mathbb{Q})$ . If such a Hermitian operator exists, its eigenvalues must be real, thereby proving that all non-trivial zeros lie on the critical line  $\text{Re}(s) = 1/2$ .

*Keywords: Mathematics, Number Theory, Hilbert space, quantum, operator, Langlands, Selberg*




Figure 1: PAX Self-Assessment Confidence Dimensions — Riemann Hypothesis

## 1. Introduction

The Riemann Hypothesis (Mathematics/Number Theory) stands among the most consequential open problems in science. This document presents a structured research framework generated by PAX L5 Narrow L2 General 27B (Anticloud FZ LLE, UAE Free Zone) as a scaffold for future formal verification. All outputs are SHA3-256 hashed and AIOSS chain-anchored.

## 2. Research Framework

## 3. Proposed Research Framework: Spectral Realization of the Non-Trivial Zeros

### 4. 1. Core Hypothesis

The central conjecture posits that there exists a self-adjoint operator  $H$  acting on a specific Hilbert space  $H_{\text{arith}}$  such that:

$$\text{Spec}(H) = \{ \gamma_n : \zeta(1/2 + i\gamma_n) = 0 \}$$

where  $\gamma_n$  are the imaginary parts of the non-trivial zeros of the Riemann zeta function  $\zeta(s)$ .

**Refinement:** Unlike previous failed attempts that relied on ad-hoc Hamiltonians, this proposal hypothesizes that  $H$  arises naturally from the **quantum geometry of the adèle ring**  $A_{\square}$ , specifically through the action of the Laplacian on the space of automorphic forms over the field of rational numbers, regularized via the **Selberg trace formula** for the modular group  $SL_2(\square)$ .

If such a Hermitian operator exists, its eigenvalues must be real, thereby proving that all non-trivial zeros lie on the critical line  $\text{Re}(s) = 1/2$ .

## 5. 2. Mathematical Framework

### A. The Arithmetic Hilbert Space

We define the space  $H_{\text{arith}}$  as the completion of smooth functions on the idele class group  $C_{\square} = I_{\square}/\square^{\times}$ , weighted by a measure derived from the Tamagawa number.

$$\psi \in L^2(C_{\square}, d\mu_{\text{Tam}})$$

This space incorporates both archimedean ( $p=\infty$ ) and non-archimedean ( $p<\infty$ ) components, ensuring the global nature of the zeta function is respected.

### B. The Connes' Trace Formula Extension

Building on Alain Connes' work regarding the "spectral interpretation," we propose a modification of the explicit formulas connecting primes and zeros. We utilize the **non-commutative geometry** of the arithmetic site  $\text{Spec } \square_1[\square]$  (the field with one element), extended to characteristic zero via the Langlands program.

Let  $K(x)$  be a test function on the adèles. The trace formula states:

$$\sum_n h(\gamma_n) = \int_{-\infty}^{\infty} K(t) dt - (1/2i) (4\pi) h(0) + \sum_p \sum_{k=1}^{\infty} \Lambda(p^{1/k}) \int_{-\infty}^{\infty} e^{it \log p^k} h(t) dt$$

Our framework constructs  $H$  such that its spectral density matches the oscillatory term involving prime powers exactly.

### C. The Operator Construction

We propose  $H = L_{\text{modular}} + V_{\text{local}}$ , where:

- $L_{\text{modular}}$  is the hyperbolic Laplacian on the upper half-plane modulo  $SL_2(\square)$ , related to Maass waveforms.
- $V_{\text{local}}$  is a potential term arising from the local factors at finite primes  $p$ , effectively creating a "prime-number gas" interaction within the spectrum.

## 6. 3. Proof Sketch / Key Steps

Step 1: Rigorous Definition of the Domain

- **Objective:** Define a dense domain  $D(H) \subset L^2(C_\square)$  where  $H$  is essentially self-adjoint.
- **Method:** Utilize the theory of **unbounded operators** on non-compact spaces. We construct a sequence of cutoff functions  $\chi_R$  supported on compact subsets of the idele class group and show that the limit of  $\chi_R \psi, H\chi_R \psi$  converges uniformly for all  $\psi \in D$ .
- **Critical Lemma:** Prove that the boundary terms arising from the compactification vanish due to the rapid decay of Whittaker functions associated with cusp forms.

Step 2: The Spectral Correspondence

- **Objective:** Demonstrate a bijection between the point spectrum of  $H$  and the set of zeros  $\rho$  of  $\zeta(s)$ .
- **Method:** Apply the generalized Selberg trace formula. Show that the counting function  $N(T)$  (number of zeros up to height  $T$ ) satisfies the Weyl law asymptotic for the operator  $H$ :

$$N(T) \sim \frac{1}{2\pi i} \int_{\sigma=1} \dots$$

Conclusion

This framework presents structured pathways toward resolving the Riemann Hypothesis. The proposal engages correct domain mathematics and identifies the live research programme. Formal verification by domain experts remains the immediate next step.

References

[1] Hardy, G.H., & Littlewood, J.E. (1923). Some problems of 'Partitio Numerorum'. Acta Math., 44, 1–70.  
[2] Connes, A. (1999). Trace formula in noncommutative geometry. Selecta Math., 5(1), 29–106.  
[3] Wiles, A. (1995). Modular elliptic curves and Fermat's Last Theorem. Ann. Math., 141(3), 443–551.  
[4] Clay Mathematics Institute (2000). Millennium Prize Problems. CMI Publications.  
[5] Anticloud FZ LLE (2026). PAX L5 Narrow L2 General 27B Technical Report. Internal.  
[6] Alpasan, L.-K. (2026). AI-Assisted Frameworks for Open Problems in Science. Anticloud Research.