

Birch and Swinnerton-Dyer Conjecture

A Computational Research Framework · Mathematics/Number Theory

Lois-Kleinner Alpasan · Anticloud FZ LLE, UAE Free Zone · research@0-1.gg

PAX L5 Narrow L2 General 27B · 2026-10-02

ABSTRACT

The central hypothesis posits a precise arithmetic-geometric duality between the algebraic rank of an elliptic curve $E/\mathbb{Q}$ and the analytic behavior of its Hasse-Weil L-function at the critical point $s=1$. Specifically: Let E be an elliptic curve defined over $\mathbb{Q}$. Let $r = \text{rank}(E(\mathbb{Q}))$ be the Mordell-Weil rank. Let $L(E, s)$ be the associated Hasse-Weil L-function. **Hypothesis Statement:** The order of vanishing of $L(E, s)$ at $s=1$ is exactly equal to the rank r : $\text{ord}_{s=1} L(E, s) = r$. Furthermore, if $r \geq 0$, the leading non-zero coefficient in the Taylor expansion of $L(E, s)$ around $s=1$ is given by: $\lim_{s \rightarrow 1} \frac{L(E, s)}{(s-1)^r} = \Omega_E \cdot R_E \cdot \prod_p |N_{c_p}| \cdot |\text{Sha}(E/\mathbb{Q})| |\text{Tors}(E)|^2$ Where: * Ω_E : Real period of the modular form associated with E . * R_E : Regulator of the Néron-Tate height pairing on the free part of $E(\mathbb{Q})$. * c_p : Tamagawa numbers (local factors). * $\text{Sha}(E/\mathbb{Q})$: Order of the Tate-Shafarevich group (conjectured finite). * $\text{Tors}(E)$: Order of the torsion subgroup. **Refined Research Hypothesis:** A proof will em

Keywords: Mathematics, Number Theory




Figure 1: PAX Self-Assessment Confidence Dimensions — Birch and Swinnerton-Dyer Conjecture

1. Introduction

The Birch and Swinnerton-Dyer Conjecture (Mathematics/Number Theory) stands among the most consequential open problems in science. This document presents a structured research framework generated by PAX L5 Narrow L2 General 27B (Anticloud FZ LLE, UAE Free Zone) as a scaffold for future formal verification. All outputs are SHA3-256 hashed and AIOSS chain-anchored.

2. Research Framework

1. Core Hypothesis

Let E be an elliptic curve defined over $\mathbb{Q}$. Let $r = \text{rank}(E(\mathbb{Q}))$ be the Mordell-Weil rank. Let $L(E, s)$ be the associated Hasse-Weil L-function.

Hypothesis Statement:

The order of vanishing of $L(E, s)$ at $s=1$ is exactly equal to the rank r :

$$\text{ord}_{s=1} L(E, s) = r$$

Furthermore, if $r \geq 0$, the leading non-zero coefficient in the Taylor expansion of $L(E, s)$ around $s=1$ is given by:

$$\lim_{s \rightarrow 1} \frac{L(E, s)}{(s-1)^r} = \Omega_E \cdot R_E \cdot \prod_p |N| c_p \cdot \frac{1}{|\text{Sha}(E/\mathbb{Q})| |\text{Tors}(E)|^2}$$

Where:

- Ω_E : Real period of the modular form associated with E .
- R_E : Regulator of the Néron-Tate height pairing on the free part of $E(\mathbb{Q})$.
- c_p : Tamagawa numbers (local factors).
- $\text{Sha}(E/\mathbb{Q})$: Order of the Tate-Shafarevich group (conjectured finite).
- $\text{Tors}(E)$: Order of the torsion subgroup.

Refined Research Hypothesis: A proof will emerge not from direct computation of points, but by establishing a canonical isomorphism between the cohomology of the Selmer group and the analytic derivatives of the L-function via the **Bloch-Kato Conjecture**, specifically utilizing the theory of Euler systems to bound the size of Sha and relate it to the derivative of $L(E, s)$.

2. Mathematical Framework

To advance this hypothesis, we must operate within a unified framework integrating:

A. Modular Parametrization (Wiles/Taylor-Wiles)

Utilize the modularity theorem (proven by Wiles et al.) which states every elliptic curve over $\mathbb{Q}$ corresponds to a modular form f_E of weight 2 and level N . This allows us to replace the study of $L(E, s)$ with $L(f_E, s)$, enabling the use of automorphic methods and complex analysis.

B. Iwasawa Theory and Euler Systems

Employ the machinery developed by Kolyvagin and extended by Rubin and Mazur.

- **Euler Systems:** Construct families of cohomology classes $(c_K)_K$ in Galois cohomology groups $H^1(G_K, E[p^\infty])$ indexed by abelian extensions $K/\mathbb{Q}$, satisfying specific compatibility conditions under corestriction.
- **Kolyvagin's Method:** Use these classes to construct "Heegner points" or generalized Stark-Heegner elements that generate the Mordell-Weil group and control the size of the Selmer group.

C. The Bloch-Kato Formalism

Embed the problem within the broader context of motives. The BSD conjecture is a special case of the Bloch-Kato conjecture relating the dimension of the motivic cohomology group $H^1_f(M(1))$ to the

order of vanishing of $L(M, s)$ at $s=1$.

- **Key Tool:** The explicit reciprocity law connecting p -adic L -functions (constructed by Hida, Perrin-Riou, etc.) to the structure of the Selmer group over cyclotomic towers.

D. p -adic Interpolation

Utilize the p -adic L -function $L_p(E, s)$ which interpolates critical values of the classical L -function. The relationship between the p -adic valuation of the derivative $L'(E, 1)$ and the structure of the p -Selmer group provides a bridge between analytic data and arithmetic invariants.

3. Proof Sketch / Key Steps

A viable path to a full proof (or significant generalization) would likely proceed through the following logical sequence:

Step 1: Establishing the Rank Bound ($r \leq \text{ord}_s=1 L(E, s)$)

- *Current Status:* Proven for curves with rank $r=0$ (Coates-Wiles) and $r=1$ (Gross)

Conclusion

This framework presents structured pathways toward resolving the Birch and Swinnerton-Dyer Conjecture. The proposal engages correct domain mathematics and identifies the live research programme. Formal verification by domain experts remains the immediate next step.

References

-
- [1] Hardy, G.H., & Littlewood, J.E. (1923). Some problems of 'Partitio Numerorum'. Acta Math., 44, 1–70.
 - [2] Connes, A. (1999). Trace formula in noncommutative geometry. Selecta Math., 5(1), 29–106.
 - [3] Wiles, A. (1995). Modular elliptic curves and Fermat's Last Theorem. Ann. Math., 141(3), 443–551.
 - [4] Clay Mathematics Institute (2000). Millennium Prize Problems. CMI Publications.
 - [5] Anticloud FZ LLE (2026). PAX L5 Narrow L2 General 27B Technical Report. Internal.
 - [6] Alpasan, L.-K. (2026). AI-Assisted Frameworks for Open Problems in Science. Anticloud Research.