

1. Plain T_EXnology

Theorem T. *All things are not necessarily the same**

2. Permutations

TAoCP in chapter 1.2.5 gives two methods to generate all permutations of a given ordered set. Quantities of permutations are considered with relevance to computing efficiencies.

3. The Wide-Awake example Group

We re-think*, re-word, and re-start with a set of attributes, elements or objects, $W = \{ \text{woozy, vacuous, sleepy, wide-awake} \}$. These elements are used to generate all possible arrangements η which are ordered n -tuples with $1 \leq n \leq 4$. For example, $\eta = (\text{woozy, wide-awake})$ is a 2-tuple. Now the set *Woozy* is the set of all permutations that jumble such elements like η .

Let $(Woozy, \circ, 0, -)$ be the group with the set *Woozy*, a binary operation \circ , a neutral element 0, and for each element $\pi \in Woozy$ there is an inverse element $-\pi \in Woozy$ such that $\pi \circ -\pi = 0$.

For now, here, we call this group's binary operation *composition*. Given two elements $\pi, \eta \in Woozy$, then $\pi \circ \eta \in Woozy$ and $\eta \circ \pi \in Woozy$.

* *T_EXbook*, *texbook.tex*, <https://www.ctan.org/tex-archive/systems/knuth/dist/tex>

* The Mathematics of the Rubiks Cube, <http://web.mit.edu/sp.268/www/rubik.pdf>

4. Creating the Woozy set

Theorem X. *An ordered set of n elements has $n!$ arrangements.*

This had a little consideration. Here, we convey our understanding of the Permutations and Factorials section.*

Given a set of objects $W = \{a_1, a_2, \dots, a_n\}$. P_n is the set of arrangements given n objects $a_1, \dots, a_n \in W$, such as $\{(a_1, a_2, \dots, a_n), (a_2, a_1, \dots), \dots\}$. For example, with $W = \{1, 2, 3\}$, we have

$$P_3 = \{(123), (231), (312), (132), (321), (213)\}.$$

Method 1, now, moves from $n = 3$ to $n = 4$ as follows. For each element in $P_{n-1} = P_3$, place element a_n in each possible vacuous position to arrive at $P_n = P_4$, that is

$$P_4 = \{(a_n a_1 a_2 a_3), (a_1 a_n a_2 a_3), (a_1 a_2 a_n a_3), (a_1 a_2 a_3 a_n), \dots, (a_n a_2 a_1 a_3), (a_2 a_n a_1 a_3), (a_2 a_1 a_n a_3), (a_2 a_1 a_3 a_n)\}$$

* TAoCP chapter 1.2.5, <https://www-cs-faculty.stanford.edu/%7Eknuth/taocp.html>

5. Accounting for these Arrangements

Adding up all permutations that are so generated we have p_n the number of all elements in P_n

And again, after some re-view, we sense a need to re-word. P_{nn} is the set of permuted n -tuples, and P_n is the, probably bigger, set of all the k -tuples with $k \in \{1, 2, \dots, n\}$. In other words, P_n may mean different things, or sets of things. This also applies to quantities that could be denoted like p_{nk} , and p_{nn} , and in case of our big wide-awake bean bag, which we sum up to p_n ; probably.

First, we started with $p_n = \sum_{k=1}^n k!$ to be the quantity p_n that accounts for all the elements of arrangements in set P_n , with $p_k = k!$ for $1 \leq k \leq n$.

However, on the back of some scrap paper, we jotted down $\{(1), (2), (3), (4)\}$ and saw that $\{(2), (3), (4)\}$ are not included in our sum, and $\{(12), (21), (13), (31), (14), (41), (23), (32), (24), (42), (34), (43)\}$ has 10 2-tuples unaccounted for, etc.)

So, for now, given that $p_{nk} = n(n-1)\dots(n-k+1)^*$, combined with $p_n = \sum_{k=1}^n p_{nk}$, we count the number of arrangements of n objects to be $p_n = \sum_{k=1}^n \frac{n!}{(n-k)!}$ or some such like.

* TAOCP chapter 1.2.5, <https://www-cs-faculty.stanford.edu/%7Eknuth/taocp.html>

6. Making concrete Space

We now look at the set W that we enumerated above and apply method 1 to arrange things.

Given W as above, we have

$$W_{oozy_{41}} = \{(\text{ woozy }), (\text{ vacuus }), (\text{ sleepy }), (\text{ wide-awake })\}$$

Then, taking one step at a time and applying method 1, given the set

$W_{oozy_{11}} = \{(\text{ sleepy })\}$ together with another element, wide-awake $\in W$, and we get

$$W_{oozy_{22}} = \{(\text{ wide-awake, sleepy }), (\text{ sleepy, wide-awake })\}.$$