

ANALYSIS 3

Contents

I	Sequence of Functions	1
1	Pointwise convergence	1
2	Uniform convergence	1
II	Series of Functions	3
3	Convergence of a Numerical Series	3
4	Finite Expansion	4
III	Power Series	5
5	Radius of Convergence	5
IV	Integrals Depending on a Parameter	7
6	Improper Integral	7
V	Fourier Series	8
7	Trigonometric Coefficients	8
7.1	Even Functions	8
7.2	Odd Functions	8
VI	Laplace Transforms	9
8	Transforms of some functions	9

8.1	Unit step function	9
8.2	Dirac Delta Function	9
8.3	Usual Elementary functions	9
9	Properties of the Transform	10
10	Translation	11
VII	Systems of Differential Equations	12
11	Solving The System of DEs	12
11.1	Change of Basis	13
11.2	Solving Matrix Formula	13
12	Fundamental Solutions	14
12.1	Wronskian of vector functions	14
13	Solving n-th order Homogeneous Linear DE	15
13.1	DE from a Set of Fundamental Solutions	16
13.2	Method of Variation of constants	16

Sequence of Functions

SECTION 1

Pointwise convergence

This is a very natural way of proving convergence since all you have to do is fix f_n to a point x then the sequence just becomes an ordinary sequence of numbers, and if they all converge to a number we can define a limit function f and say that they converge to f pointwisely.

Definition 1 We say that a sequence of functions f_n where $f_n : I \rightarrow \mathbb{R}, I \subset \mathbb{R}$, converges pointwise to function $f : I \rightarrow \mathbb{R}$ on the interval I if:

$$\forall x \in I \forall \epsilon > 0 \exists n \in \mathbb{N} \forall n \geq N : |f_n(x) - f(x)| < \epsilon.$$

You either prove convergence using the definition or by doing:

1. Let $x = 0$ then find $\lim_{n \rightarrow \infty} f_n(0) = \text{some } f(x)$
2. Then let $x \neq 0$ and again find $\lim_{n \rightarrow \infty} f_n(x) = f(x)$
3. If neither of the results are unbounded $\pm\infty$ then we say $f_n(x)$ is convergent to some $f(x)$

Remark if the result of step 1 is $g(x)$ and step 2 results in $h(x)$ where $g(x) \neq h(x)$ then we define the limit function:

$$f(x) = \begin{cases} g(x) & x = 0 \\ h(x) & x \in]0, 1] \end{cases}.$$

SECTION 2

Uniform convergence

The idea of uniform convergence is that the sequence always approaches it's limit function as the value of n increases.

Definition 2 We say that a sequence of functions f_n where $f_n : I \rightarrow \mathbb{R}, I \subset \mathbb{R}$, converges uniformly to function $f : I \rightarrow \mathbb{R}$ on the interval I if:

$$\forall \epsilon > 0 \exists N \in \mathbb{N} \forall n \geq N \forall x \in I : \sup_{x \in I} |f_n(x) - f(x)| < \epsilon.$$

Remark We can also prove uniform convergence by proving

$$\lim_{n \rightarrow \infty} \sup_{x \in I} |f_n(x) - f(x)| = 0.$$

There is also an easy way to prove uniform convergence of a function by

1. Prove that the sequence of functions $f_n(x)$ is pointwise convergent to a function $f(x)$ ¹
2. Define a function $g(x) = |f_n(x) - f(x)|$ and find the maxima of that function at a point x_0 (usually by doing $dg/dx = 0$)
3. If $\lim_{n \rightarrow \infty} g(x_0) = 0$ then the sequence converges uniformly to $f(x)$

¹if the f is continuous then the convergence is uniform

Series of Functions

Definition 3

Let $f_n(x)$ be sequence of functions defined on $I \subset \mathbb{R}$, we define the series $S(x)$ to be

$$S(x) = \sum_{n=0}^{\infty} f_n(x).$$

SECTION 3

Convergence of a Numerical Series

In order to prove a series of functions converge we have to prove that it converges for all fixed x .

Theorem 1

Suppose there exists a sequence a_n such that $\forall x, n \ |f_n| \leq a_n$. The Weierstrass test states that if $\sum a_n$ converges then $\sum f_n(x)$ converges uniformly and absolutely

Theorem 2

Let a_n be a sequence of numbers, if $\left| \frac{a_{n+1}}{a_n} \right| = l$ then the sequence is a geometric Series

$$\sum_{n=0}^{\infty} a_n \begin{cases} \text{converges} & \text{if } |l| < 1 \\ \text{diverges} & \text{if } |l| \geq 1 \end{cases}.$$

Theorem 3

A harmonic series is defined to be $a_n = \frac{1}{n^p}$

$$\sum_{n=0}^{\infty} \frac{1}{n^p} \begin{cases} \text{converges} & \text{if } p > 1 \\ \text{diverges} & \text{if } p \leq 1 \end{cases}.$$

Theorem 4

Let a_n be a sequence of numbers. The 2 series $\sum_{n=0}^{\infty} a_n$ and $\sum_{n=0}^{\infty} 2^n a_n$ are simultaneously convergent/divergent.

Theorem 5

The sequence $\sum_{n=0}^{\infty} (-1)^n a_n$ is convergent if a_n is decreasing and $\lim_{n \rightarrow \infty} a_n = 0$.

Theorem 6

Consider the series $S = \sum_{n=0}^{\infty} a_n$

$$\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} = l \quad \text{such that} \quad \begin{cases} l < 1 & \text{if } S \text{ converges} \\ l > 1 & \text{if } S \text{ diverges} \\ l = 1 & \text{this test cannot help us} \end{cases}.$$

SECTION 4

Finite Expansion

The general formula for the finite expansion (Taylor-young formula) is

$$f(x) = f(x-a) + \frac{x}{1!} f'(x-a) + \frac{x^2}{2!} f''(x-a) + \cdots + \frac{x^n}{n!} f^{(n)}(x-a) + x^n o(1) \quad x \rightarrow a.$$

Some important expansions to keep in mind are

a) $e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$

b) $\sin(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}$

c) $\cos(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$

d) $\frac{1}{1-x} = \sum_{n=0}^{\infty} x^n$

e) $\sinh(x) = \sum_{n=0}^{\infty} \frac{x^{2n+1}}{(2n+1)!}$

f) $\cosh(x) = \sum_{n=0}^{\infty} \frac{x^{2n}}{(2n)!}$

g) $\ln(1+x) = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n}$

h) $\ln(1-x) = \sum_{n=1}^{\infty} -\frac{x^n}{n}$

i) $(1+x)^\alpha = \sum_{n=0}^{\infty} \frac{\prod_{k=0}^{n-1} (\alpha - k)}{n!} x^n$

Power Series

A power series is just a series in the following formula

$$\begin{aligned} f(x) &= \sum_{n=0}^{\infty} a_n(x - x_0)^n \\ &= \sum_{n=0}^{\infty} U_n. \end{aligned}$$

SECTION 5

Radius of Convergence

For some values of x a power series can either diverge or converge, to determine the interval of convergence we employ the ratio test

Theorem 7 Let r be the radius of convergence and I be the domain of convergence. If we compute the limit

$$\Gamma = \lim_{n \rightarrow \infty} \left| \frac{U_{n+1}}{U_n} \right|.$$

The ratio test states that

$$\begin{cases} \text{if } \Gamma = 0 & \text{then } r = \infty \text{ and } I = \mathbb{R} \\ \text{if } \Gamma = \infty & \text{then } r = 0 \text{ and } I = \{0\} \end{cases}.$$

In the case that Γ isn't 0 or ∞ we set $\Gamma < 1$ and then we find $|x| < R$, finally we can say that $r = R$ and $I =]-R, R[$. A special case need to be done for the points $-R$ and R to determine if they belong in I .

Remark The power series $f(x)$ is continuous and will always uniformly converge in the interval of convergence I

Theorem 8 If $\sum a_n x^n$ and $\sum b_n x^n$ be 2 power series with radii R_1 and R_2 . For the power series $\sum (a_n + b_n) x^n$ the radius of convergence R

$$R = \min\{R_1, R_2\}.$$

Theorem 9 If $S(x) = \sum_{n=0}^{\infty} a_n(x - x_0)^n$ is a power series with radius R then $S'(x) = \sum_{n=0}^{\infty} n a_n(x - x_0)^{n-1}$ as well as $\int_{x_0}^x S(t) dt$ both a radius of R

The general term a_k of a power series $S(x) = \sum_{n=0}^{\infty} a_n(x - x_0)^n$ is equal to

$$a_k = \frac{S^{(k)}(x_0)}{k!}.$$

$$\begin{aligned}y &= a_0 + a_1x + a_2x^2 + \cdots + a_nx^n \\y' &= a_1 + 2a_2x + 3a_3x^2 + \cdots + na_nx^{n-1} \\y'' &= 2a_2 + 6a_3x + \cdots + n(n-1)a_nx^{n-2} + n(n+1)a_{n+1}x^{n-1}\end{aligned}$$

Integrals Depending on a Parameter

We define an integratable function

$$\begin{aligned} f : I \times U &\longrightarrow \mathbb{R} \times \mathbb{R} \\ t, x &\longmapsto f(t, x). \end{aligned}$$

where t is a parameter.

Let

$$F(x) = \int_a^b f(t, x) dt.$$

Theorem 10 If f is of class C^p then F is also of class C^n and

$$\frac{\partial^p F}{\partial x^p} = \int_a^b \frac{\partial^p f}{\partial x^p} dt.$$

Remark F is differentiable if $\frac{\partial f}{\partial x}$ is convergent.

SECTION 6

Improper Integral

Theorem 11 If $\int_a^b |f(t)| dt$ is convergent then $\int_a^b f(t) dt$ is also convergent and

$$\left| \int_a^b f(t) dt \right| \leq \int_a^b |f(t)| dt$$

Theorem 12 The Weierstrass test states that $\int_a^b f(t, x) dt$ is convergent if there exists a function $g(t)$ such that $|f(t, x)| \leq g(t)$ and $\int_a^b g(t) dt$

Fourier Series

Definition 4 A Fourier series is a series of functions of general term

$$u_n(x) = a_0 + a_n \cos(nx) + b_n \sin(nx).$$

SECTION 7

Trigonometric Coefficients

The Fourier coefficients of a function defined on an interval $F \subset \mathbb{R}$ of period $T = \frac{2\pi}{\omega} \implies \omega = \frac{2\pi}{T}$ are

$$\begin{aligned} a_0 &= \frac{1}{T} \int_F f(x) \, dx \\ a_n &= \frac{2}{T} \int_F f(x) \cos \omega n x \, dx \\ b_n &= \frac{2}{T} \int_F f(x) \sin \omega n x \, dx \end{aligned}$$

1. $\sin n \pi = 0$
2. $\sin n \pi/2 = (-1)^n$
3. $\cos n \pi = (-1)^n$
4. $\cos n \pi/2 = 0$

SUBSECTION 7.1

Even Functions

If a function has a domain $F = [-\ell; \ell]$ and $f(x) = f(-x) \, \forall x \in \mathbb{R}$ the Fourier coefficients ($T = |F|$) become

$$\begin{aligned} a_0 &= \frac{1}{\ell} \int_0^\ell f(x) \, dx \\ a_n &= \frac{2}{\ell} \int_0^\ell f(x) \cos nx \, dx \\ b_n &= 0 \end{aligned}$$

SUBSECTION 7.2

Odd Functions

If a function has a domain $F = [-\ell; \ell]$ and $-f(x) = f(-x) \, \forall x \in \mathbb{R}$ the Fourier coefficients ($T = |F|$) become

$$\begin{aligned} a_0 &= 0 \\ a_n &= 0 \\ b_n &= \frac{2}{\ell} \int_0^\ell f(x) \sin nx \, dx \end{aligned}$$

Laplace Transforms

The Laplace transform of a function is defined as

$$F(p) = \mathcal{L}\{f(t)\} = \int_0^{\infty} e^{-pt} f(t) dt.$$

It only exists if the integral above converges.

SECTION 8

Transforms of some functions

SUBSECTION 8.1

Unit step function

Also known as Heaviside's unit step function, it is defined as

$$u(t) = \begin{cases} 1 & \text{if } t \geq 0 \\ 0 & \text{if } t < 0 \end{cases}.$$

$$\mathcal{L}\{u(t)\} = \frac{1}{p} \quad \text{for } \operatorname{Re}(p) > 0.$$

SUBSECTION 8.2

Dirac Delta Function

The Dirac Delta function

$$\delta(t) = \begin{cases} \infty & \text{if } t = 0 \\ 0 & \text{if } t \neq 0 \end{cases}.$$

$$\mathcal{L}\{\delta(t)\} = 1.$$

SUBSECTION 8.3

Usual Elementary functions

$$\text{a) } \mathcal{L}\{1\} = \frac{1}{p}$$

$$\text{b) } \mathcal{L}\{t\} = \frac{1}{p^2}$$

$$\text{c) } \mathcal{L}\{t^n\} = \frac{n!}{p^{n+1}}$$

$$\text{d) } \mathcal{L}\{\sin \omega t\} = \frac{\omega}{p^2 + \omega^2}$$

$$\begin{array}{ll} \text{e) } \mathcal{L}\{\cos \omega t\} = \frac{p}{p^2 + \omega^2} & \text{f) } \mathcal{L}\{\sinh \omega t\} = \frac{\omega}{p^2 - \omega^2} \\ \text{g) } \mathcal{L}\{\cosh \omega t\} = \frac{p}{p^2 - \omega^2} & \text{h) } \mathcal{L}\{e^{at}\} = \frac{1}{p - a} \end{array}$$

SECTION 9

Properties of the Transform

1. Linearity:

$$\mathcal{L}\{\lambda f + \mu g\} = \lambda \mathcal{L}\{f\} + \mu \mathcal{L}\{g\}.$$

2. Homothety:

$$\mathcal{L}\{f(kt)\} = \frac{1}{k} F\left(\frac{p}{k}\right).$$

3. Derivation:

$$\begin{aligned} \mathcal{L}\{f'(t)\} &= p \mathcal{L}\{f(t)\} - f(0^+) \\ \mathcal{L}\{f''(t)\} &= p^2 \mathcal{L}\{f(t)\} - pf(0^+) - f'(0^+) \\ \mathcal{L}\{f^{(n)}(t)\} &= p^n \mathcal{L}\{f(t)\} - \sum_{k=1}^n p^{n-k} f^{(k-1)}(0^+) \end{aligned}$$

4. Integration:

$$\mathcal{L}\left\{\int_0^t f(u) \, du\right\} = \frac{F(p)}{p}.$$

5. Initial value theorem:

$$f(0^+) = \lim_{p \rightarrow \infty} p \mathcal{L}\{f(t)\}.$$

6. Final value theorem:

$$f(\infty) = \lim_{p \rightarrow 0} p \mathcal{L}\{f(t)\}.$$

Remark

$$\begin{aligned} \mathcal{L}\{tf(t)\} &= -\frac{d}{dp} F(p) \\ \mathcal{L}\{t^2 f(t)\} &= \frac{d^2}{dp^2} F(p) \\ \mathcal{L}\{t^n f(t)\} &= (-1)^n \frac{d^n}{dp^n} F(p) \end{aligned}$$

Remark Convolution over a domain $I \subset \mathbb{R}$ is defined as

$$f(t) * g(t) = \int_I f(\tau) g(t - \tau) \, d\tau = \int_I f(t - \tau) g(\tau) \, d\tau.$$

and it's transform is

$$\mathcal{L}\{f(t) * g(t)\} = F(p) \cdot G(p).$$

SECTION 10

Translation

In the time domain:

$$\mathcal{L}\{f(t - a)\} = e^{-ap}F(p).$$

In the p -domain:

$$\mathcal{L}\{e^{at}f(t)\} = F(p + a).$$

Systems of Differential Equations

Consider a system of first order differential equations (S)

$$\begin{cases} \frac{dx_1}{dt} = a_{11}(t)x_1 + a_{12}(t)x_2 + \cdots + a_{1n}(t)x_n + b_1(t) \\ \frac{dx_2}{dt} = a_{21}(t)x_1 + a_{22}(t)x_2 + \cdots + a_{2n}(t)x_n + b_2(t) \\ \vdots \\ \frac{dx_n}{dt} = a_{n1}(t)x_1 + a_{n2}(t)x_2 + \cdots + a_{nn}(t)x_n + b_n(t) \end{cases}.$$

in matrix form the equation can be written as

$$\frac{d\vec{x}}{dt} = A\vec{x} + \vec{b}.$$

and the initial condition can be written as

$$\vec{x}_0(t_0) = \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{pmatrix}.$$

SECTION 11

Solving The System of DEs

There 3 main ways of solving systems of DEs:

1. Laplace Transform
2. Change of Basis
3. Solving Matrix Formula

Remark Let A be a diagonalizable matrix

$$A = PDP^{-1}.$$

so we define that

$$e^{At} = Pe^{Dt}P^{-1}.$$

or in other words

$$e^{At} = P \begin{pmatrix} e^{\lambda_1 t} & 0 & 0 & 0 \\ 0 & e^{\lambda_2 t} & 0 & 0 \\ 0 & 0 & \ddots & 0 \\ 0 & 0 & 0 & e^{\lambda_n t} \end{pmatrix} P^{-1}.$$

SUBSECTION 11.1

Change of Basis

We consider a new system of DEs to be

$$\frac{d\vec{y}}{dt} = P^{-1}AP\vec{y} + P^{-1}\vec{b}.$$

which simplifies to

$$\frac{d\vec{y}}{dt} = D\vec{y} + \vec{B}.$$

in this new system we can solve for \vec{y}

$$\begin{cases} \frac{dy_1}{dt} = \lambda_1 y_1 + B_1 \\ \frac{dy_2}{dt} = \lambda_2 y_2 + B_2 \\ \vdots \\ \frac{dy_n}{dt} = \lambda_n y_n + B_n \end{cases}.$$

Remark The solution to a differential equation of the form

$$\frac{dy}{dt} = \alpha y + \beta.$$

is

$$y = c_1 e^{\alpha t} - \frac{\beta}{\alpha}.$$

after we find the solution to the new system, we can simply obtain the solution to the original system by

$$\vec{x} = P\vec{y}.$$

and by substituting t_0 in \vec{x} we can solve for the constant terms (c_1, c_2, \dots, c_n) using \vec{x}_0 .

SUBSECTION 11.2

Solving Matrix Formula

The formula for a system of first order equations is

$$\vec{x} = \vec{x}_h + \vec{x}_p.$$

Where

$$\begin{aligned}\vec{x}_h &= V(t, t_0)\vec{x}_0 \\ \vec{x}_p &= \int_{t_0}^t V(t, u)\vec{b}(u) \, du\end{aligned}$$

where

$$V(t, t_0) = X(t)X^{-1}(t_0).$$

if $t = 0$ then the formula becomes

$$\vec{x} = e^{At}\vec{x}_0 + \int_0^t e^{A(t-u)}\vec{b}(u) \, du.$$

SECTION 12

Fundamental Solutions

For any given system of homogeneous linear DEs there exists a set of n functions such they for a linearly independent basis for a general solution of said DEs, in other words for a given DE there exists a set of vector functions $(\vec{\zeta}_1, \vec{\zeta}_2, \dots, \vec{\zeta}_n)$ such that

$$\vec{x}(t) = c_1\vec{\zeta}_1(t) + c_2\vec{\zeta}_2(t) + \dots + c_n\vec{\zeta}_n(t) \quad \text{where} \quad c_1, c_2, \dots, c_n \in \mathbb{R}.$$

We define the fundamental matrix of the system

$$X = \begin{pmatrix} \vec{\zeta}_1 & \vec{\zeta}_2 & \dots & \vec{\zeta}_n \end{pmatrix} = \begin{pmatrix} \zeta_{11} & \zeta_{12} & \dots & \zeta_{1n} \\ \zeta_{21} & \zeta_{22} & \dots & \zeta_{2n} \\ \vdots & & \ddots & \\ \zeta_{n1} & \zeta_{n2} & \dots & \zeta_{nn} \end{pmatrix}.$$

The system can be written in terms of X as

$$\frac{dX}{dt} = AX.$$

Remark The fundamental solutions are linearly independent $\implies \det(X) \neq 0$

SUBSECTION 12.1

Wronskian of vector functions

Consider the vector functions:

$$\vec{\phi}_1(t) = \begin{pmatrix} \phi_{11}(t) \\ \phi_{21}(t) \\ \vdots \\ \phi_{n1}(t) \end{pmatrix} \quad \dots \quad \vec{\phi}_n(t) = \begin{pmatrix} \phi_{1n}(t) \\ \phi_{2n}(t) \\ \vdots \\ \phi_{nn}(t) \end{pmatrix}.$$

The Wronskian is defined to the determinant:

$$W(\vec{\phi}_1, \vec{\phi}_2, \dots, \vec{\phi}_n) = \begin{vmatrix} \phi_{11} & \phi_{12} & \cdots & \phi_{1n} \\ \phi_{21} & \phi_{22} & \cdots & \phi_{2n} \\ \vdots & & \ddots & \\ \phi_{n1} & \phi_{n2} & \cdots & \phi_{nn} \end{vmatrix}.$$

If the Wronskian = 0 then the functions $(\vec{\phi}_1, \vec{\phi}_2, \dots, \vec{\phi}_n)$ are said to be linearly independent.

When dealing with DEs the concept of a Wronskian can be applied to *non-vector functions* as follows

$$W(\phi_1, \phi_2, \dots, \phi_n) = \begin{vmatrix} \phi_1 & \phi_2 & \cdots & \phi_n \\ \phi'_1 & \phi'_2 & \cdots & \phi'_n \\ \vdots & & \ddots & \\ \phi_1^{(n-1)} & \phi_2^{(n-1)} & \cdots & \phi_n^{(n-1)} \end{vmatrix}.$$

SECTION 13

Solving n -th order Homogeneous Linear DE

We define the notation $D^n x = \frac{d^n x}{dt^n}$.

An n -th order linear DE is any equation of the form:

$$D^n x + a_1(t)D^{n-1}x + \cdots + a_{n-1}(t)Dx + a_n(t)x = 0.$$

We can then write the equation in vector form

$$\begin{aligned} x &= x_1 \\ Dx &= x_2 \\ D^2x &= x_3 \\ &\vdots \\ D^{n-1}x &= x_n \\ D^n x &= -a_n x_1 - a_{n-1} x_2 - \cdots - a_1 x_n \end{aligned}$$

and we take

$$\vec{x} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = \begin{pmatrix} x \\ Dx \\ \vdots \\ D^{n-1}x \end{pmatrix}.$$

then we can write the system as

$$\frac{d\vec{x}}{dt} = A\vec{x} \quad \text{where} \quad A = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & & & \ddots & \\ 0 & 0 & 0 & \cdots & 1 \\ -a_n & -a_{n-1} & \cdots & \cdots & -a_1 \end{pmatrix}.$$

SUBSECTION 13.1

DE from a Set of Fundamental Solutions

Given a set of fundamental solutions $(\zeta_1, \zeta_2, \dots, \zeta_n)$, due to the uniqueness theorem those solutions only satisfy one DE. To find that DE we simply compute

$$W(x, \zeta_1, \zeta_2, \dots, \zeta_n) = \begin{vmatrix} x & \zeta_1 & \cdots & \zeta_n \\ D x & D \zeta_1 & \cdots & D \zeta_n \\ \vdots & & \ddots & \\ D^n x & D^n \zeta_1 & \cdots & D^n \zeta_n \end{vmatrix} = 0.$$

Example | Given the fundamental set of solutions $(e^{\omega t}, e^{-\omega t})$, find the second order homogeneous equation for that set of solutions:

$$\begin{aligned} W(x, e^{\omega t}, e^{-\omega t}) &= 0 \\ \implies \begin{vmatrix} x & e^{\omega t} & e^{-\omega t} \\ x' & \omega e^{\omega t} & -\omega e^{-\omega t} \\ x'' & \omega^2 e^{\omega t} & \omega^2 e^{-\omega t} \end{vmatrix} &= 0 \\ \implies x'' - \omega^2 x &= 0. \end{aligned}$$

SUBSECTION 13.2

Method of Variation of constants
