

1      The importance of higher moments in  
2                      VaR and cVaR estimation.



3

4                      Faes E.<sup>1</sup>    Mertens de Wilmars S.<sup>2</sup>    Pratesi F.<sup>3</sup>

5                                      Antwerp Management School

6                      Prof. dr. Annaert    Prof. dr. De Ceuster    Prof. dr. Zhang

7                                      *Master in Finance*

8                                      June 2021

<sup>1</sup>Enjo.Faes@student.ams.ac.be

<sup>2</sup>Stephane.MertensdeWilmars@student.ams.ac.be

<sup>3</sup>Filippo.Pratesi@student.ams.ac.be

For our families and loved ones

# Acknowledgements

First of all, many thanks to our families and loved ones that supported us during the writing of this thesis. Secondly, thank you professors Zhang, Annaert and De Ceuster for the valuable insights you have given us in preparation of this thesis and the many questions answered. We must be grateful for the classes of R programming by prof Zhang.

Secondly, we have to thank the developer of the software we used for our thesis. A profuse thanks to Allaire, the founder and CEO of RStudio. Thanks for making data science easier, more accessible and fun. We must also be grateful to Gruber for inventing “Markdown”, to MacFarlane for creating “Pandoc” which converts Markdown to a large number of output formats, and to Xie for creating “knitr” which introduced R Markdown as a way of embedding code in Markdown documents, and “bookdown” which added tools for technical and longer-form writing. Special thanks to Ismay, who created the “thesisdown” package that helped many PhD students write their theses in R Markdown. And a very special thanks to McManigle, whose adaption of Evans’ adaptation of Gillow’s original maths template for writing an Oxford University DPhil thesis in “LaTeX” provided the template that Ulrik Lyngs in turn adapted for R Markdown, which we also owe a big thank you. Without which this thesis could not have been written in this format (Lyngs 2019).

Finally, we thank Ghalanos (2020b) for making the implementation of GARCH models integrated in R via his package “Rugarch”. By doing this, he facilitated the process of understanding the whole process and doing the analysis for our thesis.

Enjo Faes,  
Stephane Mertens de Wilmars,  
Filippo Pratesi  
Antwerp Management School, Antwerp  
27 June 2021

# Abstract

<sup>41</sup> The greatest abstract all times

# Contents

43	<b>List of Figures</b>	<b>vii</b>
44	<b>List of Tables</b>	<b>viii</b>
45	<b>List of Abbreviations</b>	<b>ix</b>
46	<b>Introduction</b>	<b>1</b>
47	<b>1 Literature review</b>	<b>4</b>
48	1.1 Stylized facts of returns . . . . .	4
49	1.2 SGT (Skewed Generalized t-distribution) . . . . .	5
50	1.3 Volatility modeling . . . . .	6
51	1.3.1 Rolling volatility . . . . .	6
52	1.3.2 ARCH model . . . . .	6
53	1.3.3 Univariate GARCH models . . . . .	10
54	1.4 ACD models . . . . .	12
55	1.5 Value at Risk . . . . .	13
56	1.6 Conditional Value at Risk . . . . .	13
57	1.7 Past literature on the consequences of higher moments for VaR	
58	determination . . . . .	15
59	<b>2 Data and methodology</b>	<b>16</b>
60	2.1 Data . . . . .	16
61	2.1.1 Descriptives . . . . .	17
62	2.2 Methodology . . . . .	24
63	2.2.1 Garch models . . . . .	24
64	2.2.2 ACD models . . . . .	25
65	2.2.3 Analysis Tests VaR and cVaR . . . . .	26

66	<b>3 Empirical Findings</b>	<b>28</b>
67	3.1 Density of the returns . . . . .	28
68	3.1.1 MLE distribution parameters . . . . .	28
69	3.2 Results of GARCH with constant higher moments . . . . .	30
70	3.3 Results of GARCH with time-varying higher moments . . . . .	36
71	<b>4 Robustness Analysis</b>	<b>38</b>
72	4.1 Specification checks . . . . .	38
73	4.1.1 Figures control tests . . . . .	38
74	4.1.2 Residual heteroscedasticity . . . . .	38
75	<b>5 Conclusion</b>	<b>39</b>
76	<b>Appendices</b>	
77	<b>A Appendix</b>	<b>42</b>
78	A.1 Alternative distributions than the normal . . . . .	42
79	A.1.1 Student's t-distribution . . . . .	42
80	A.1.2 Generalized Error Distribution . . . . .	43
81	A.1.3 Skewed t-distribution . . . . .	43
82	A.1.4 Skewed Generalized Error Distribution . . . . .	44
83	A.2 GARCH models . . . . .	44
84	A.2.1 GARCH model . . . . .	44
85	A.2.2 IGARCH model . . . . .	45
86	A.2.3 EGARCH model . . . . .	46
87	A.2.4 GJRARCH model . . . . .	46
88	A.2.5 NAGARCH model . . . . .	46
89	A.2.6 TGARCH model . . . . .	47
90	A.2.7 TSGARCH model . . . . .	47
91	A.2.8 EWMA . . . . .	48
92	<b>Works Cited</b>	<b>49</b>

# List of Figures

94	2.1	Eurostoxx 50 Price Index prices . . . . .	20
95	2.2	Eurostoxx 50 Price Index log returns . . . . .	21
96	2.3	Eurostoxx 50 rolling volatility (22 days, calculated over 252 days) .	22
97	2.4	Density vs. Normal Eurostoxx 50 log returns) . . . . .	22
98	2.5	Absolute prediction errors . . . . .	23

# List of Tables

100	1.1	GARCH models, the founders . . . . .	11
101	1.2	Higher moments and VaR . . . . .	15
102	2.1	Summary statistics of the returns . . . . .	19
103	3.1	Maximum likelihood estimates of unconditional distribution functions	29



# List of Abbreviations

105	<b>ACD</b>	. . . . .	Autoregressive Conditional Density models (Hansen, 1994)
106	<b>ARCH</b>	. . . . .	Autoregressive Conditional Heteroscedasticity model (Bollerslev,
107			1986)
108	<b>GARCH</b>	. . . .	Generalized Autoregressive Conditional Heteroscedasticity model
109			(Bollerslev, 1986)
110	<b>IGARCH</b>	. . . .	Integrated GARCH (Bollerslev, 1986)
111	<b>EGARCH</b>	. . . .	Exponential GARCH (Nelson, 1991)
112	<b>GJRARCH</b>		Glosten-Jagannathan-Runkle GARCH model (Glosten et al.
113			1993)
114	<b>NAGARCH</b>	. . . .	Nonlinear asymmetric GARCH (Engle and Ng, 1993)
115	<b>TGARCH</b>	. . . .	Threshold GARCH (Zakoian, 1994)
116	<b>TSGARCH</b>	. . . .	Also called Absolute Value GARCH or AVGARCH referring to
117			Taylor (1986) and Schwert (1989)
118	<b>EWMA</b>	. . . . .	Exponentially Weighted Moving Average model
119	<b>i.i.d, iid</b>	. . . . .	Independent and identically distributed
120	<b>T</b>	. . . . .	Student's T-distribution
121	<b>ST</b>	. . . . .	Skewed Student's T-distribution
122	<b>SGT</b>	. . . . .	Skewed Generalized T-distribution
123	<b>GED</b>	. . . . .	Generalized Error Distribution
124	<b>SGED</b>	. . . . .	Skewed Generalized Error Distribution
125	<b>NORM</b>	. . . . .	Normal distribution
126	<b>VaR</b>	. . . . .	Value-at-Risk
127	<b>cVaR</b>	. . . . .	Expected shortfall or conditional Value-at-Risk

## Introduction

129 A general assumption in finance is that stock returns are normally distributed.  
130 However, various authors have shown that this assumption does not hold in practice:  
131 stock returns are not normally distributed (Officer 1972). For example, Theodossiou  
132 (2000) mentions that “empirical distributions of log-returns of several financial assets  
133 exhibit strong higher-order moment dependencies which exist mainly in daily and  
134 weekly log-returns and prevent monthly, bimonthly and quarterly log-returns from  
135 obeying the normality law implied by the central limit theorem. As a consequence,  
136 price changes do not follow the geometric Brownian motion.” So in reality, stock  
137 returns exhibit fat-tails and peakedness (Officer 1972), these are some of the so-  
138 called stylized facts of returns.

139

140 Additionally, a point of interest is the predictability of stock prices. Fama (1965)  
141 explains that the question in academic and business circles is: “To what extent can  
142 the past history of a common stock’s price be used to make meaningful predictions  
143 concerning the future price of the stock?”. There are two viewpoints towards the  
144 predictability of stock prices. Firstly, some argue that stock prices are unpredictable  
145 or very difficult to predict by their past returns (i.e. have very little serial correlation)  
146 because they simply follow a Random Walk process (Fama 1970). On the other hand,  
147 Lo & MacKinlay mention that “financial markets *are* predictable to some extent  
148 but far from being a symptom of inefficiency or irrationality, predictability is the oil  
149 that lubricates the gears of capitalism”. Furthermore, there is also no real robust  
150 evidence for the predictability of returns themselves, let alone be out-of-sample  
151 (Welch and Goyal 2008). This makes it difficult for corporations to manage market

152 risk, i.e. the variability of stock prices.

153  
154 Risk, in general, can be defined as the volatility of unexpected outcomes  
155 (Jorion 2007). The measure Value at Risk (VaR), developed in response to the  
156 financial disaster events of the early 1990s, has been very important in the financial  
157 world. Corporations have to manage their risks and thereby include a future risk  
158 measurement. The tool of VaR has now become a standard measure of risk for many  
159 financial institutions going from banks, that use VaR to calculate the adequacy of  
160 their capital structure, to other financial services companies to assess the exposure  
161 of their positions and portfolios. The 5% VaR can be informally defined as the  
162 maximum loss of a portfolio, during a time horizon, excluding all the negative events  
163 with a combined probability lower than 5% while the Conditional Value at Risk  
164 (CVaR) can be informally defined as the average of the events that are lower than  
165 the VaR. Bali, Mo, et al. (2008) explains that many implementations of the CVaR  
166 have the assumption that asset and portfolio's returns are normally distributed but  
167 that it is an inconsistency with the evidence empirically available which outlines  
168 a more skewed distribution with fatter tails than the normal. This lead to the  
169 conclusion that the assumption of normality, which simplifies the computation of  
170 VaR, can bring to incorrect numbers, underestimating the probability of extreme  
171 events happening.

172  
173 This paper has the aim to replicate and update the research made by Bali, Mo,  
174 et al. (2008) on US indexes, analyzing the dynamics proposed with a European  
175 outlook. The main contribution of the research is to provide the industry with a  
176 new approach to calculating VaR with a flexible tool for modeling the empirical  
177 distribution of returns with higher accuracy and characterization of the tails.

178  
179 The paper is organized as follows. Chapter 1 discusses at first the alternative  
180 distribution than the normal that we are going to evaluate during the analysis  
181 (Student's t-distribution, Generalized Error Distribution, Skewed t-distribution,

## *Introduction*

182 Skewed Generalized Error Distribution, Skewed Generalized t-distribution), then the  
183 discrete time GARCH models used (GARCH, IGARCH, EGARCH, GJRGARCH,  
184 NAGARCH, TGARCH, TSGARCH or AVGARCH and EWMA) are presented as  
185 extensions of the Engle (1982) 's ARCH model. Chapter 2 describes the dataset  
186 used and the methodology followed in modeling the volatility with the GARCH  
187 model by Bollerslev (1986) and with its refinements using Maximum likelihood  
188 estimation to find the distribution parameters. Then a description is given of how  
189 are performed the control tests (un- and conditional coverage test, dynamic quantile  
190 test) used in the paper to evaluate the performances of the different GARCH models  
191 and underlying distributions. In chapter 3, findings are presented and discussed,  
192 in chapter 4 the findings of the performed tests are shown and interpreted and in  
193 chapter 5 the investigation and the results are summarized.

# 1

## Literature review

### 1.1 Stylized facts of returns

When analyzing returns as a time-series, we look at log returns. The log returns are similar to simple returns so the stylized facts of returns apply to both. One assumption that is made often in financial applications is that returns are iid, or independently and identically distributed, another is that they are normally distribution. Are these valid assumptions? Below the stylized facts<sup>1</sup> following Annaert (2021) for returns are given.

- Returns are *small and volatile* (with the standard deviation being larger than the mean on average)
- Returns have very little serial correlation as mentioned by for example Bollerslev (1987).
- Returns exhibit conditional heteroskedasticity, or *volatility clustering*. There is no constant variance, but it is time-varying (homoskedasticity). Bollerslev (1987) describes it as “rates of return data are characterized by volatile and tranquil periods”.

---

<sup>1</sup>Stylized facts are the statistical properties that appear to be present in many empirical asset returns (across time and markets)

## 1. Literature review

- Returns also exhibit *asymmetric volatility*, in that sense volatility increases more after a negative return shock than after a large positive return shock. This is also called the *leverage effect*.
- Returns are *not normally distributed* which is also one of the conclusions by Fama (1965). Returns have tails fatter than a normal distribution (leptokurtosis) and thus are riskier than under the normal distribution. Log returns **can** be assumed to be normally distributed. However, this will be examined in our empirical analysis if this is appropriate. This makes that simple returns follow a log-normal distribution, which is a skewed density distribution.

Firms holding a portfolio have a lot of things to consider: expected return of a portfolio, the probability to get a return lower than some threshold, the probability that an asset in the portfolio drops in value when the market crashes. All the previous requires information about the return distribution or the density function. What we know from the stylized facts of returns that the normal distribution is not appropriate for returns. In appendix we summarize some alternative distributions (T, GED, ST, SGED) that could be a better approximation of returns than the normal one. Below

## 1.2 SGT (Skewed Generalized t-distribution)

The SGT distribution of introduced by Theodossiou (1998) and applied by Bali and Theodossiou (2007) and Bali, Mo, et al. (2008). According to Bali, Mo, et al. (2008) the proposed solutions (use of historical simulation, student's t-distribution, generalized error distribution or a mixture of two normal distributions) to the non-normality of standardized financial returns only partially solved the issues of skewness and leptokurtosis. The density of the generalized t-distribution of McDonald and Newey (1988) is given by equation (1.1) (Bollerslev et al. 1994).

$$f(\varepsilon_t \sigma_t^{-1}; p, \psi) = \frac{p}{2\sigma_t \cdot \psi^{1/p} B(1/p, \psi) \cdot [1 + |\varepsilon_t|^p / (\psi b^p \sigma_t^p)]^{\psi+1/p}} \quad (1.1)$$

where  $B(1/\eta, \psi)$  is the beta function ( $=\Gamma(1/\eta)\Gamma(\psi)\Gamma(1/\eta + \psi)$ ),  $\psi\eta > 2$ ,  $\eta > 0$  and  $\psi > 0$ ,  $\beta = [\Gamma(\psi)\Gamma(1/\eta)/\Gamma(3/\eta)\Gamma(\psi - 2/\eta)]^{1/2}$ , the scale factor and one shape parameter  $p$ .

Again the skewed variant is given by equation (A.4) of appendix but with  $f_1(\cdot)$  equal to equation (1.1) following Trottier and Ardia (2015).

## 1.3 Volatility modeling

### 1.3.1 Rolling volatility

When volatility needs to be estimated on a specific trading day, the method used as a descriptive tool would be to use rolling standard deviations. Engle (2001) explains the calculation of rolling standard deviations, as the standard deviation over a fixed number of the most recent observations. For example, for the past month it would then be calculated as the equally weighted average of the squared deviations from the mean (i.e. residuals) from the last 22 observations (the average amount of trading or business days in a month). All these deviations are thus given an equal weight. Also, only a fixed number of past recent observations is examined. Engle regards this formulation as the first ARCH model.

### 1.3.2 ARCH model

Autoregressive Conditional Heteroscedasticity (ARCH) models, proposed by Engle (1982), was in the first case not used in financial markets but on inflation. Since then, it has been used as one of the workhorses of volatility modeling. To fully capture the logic behind GARCH models, the building blocks are examined in the first place. There are three building blocks of the ARCH model: returns, the innovation process and the variance process (or volatility function), written out in respectively equation (1.2), (1.3) and (1.4). Returns are written as a constant part ( $\mu$ ) and an unexpected part, called noise or the innovation process. The innovation process is the volatility ( $\sigma_t$ ) times  $z_t$ , which is an independent identically distributed random variable with a mean of 0 (zero-mean) and a variance of 1 (unit-variance).

## 1. Literature review

263 The independent from iid, notes the fact that the  $z$ -values are not correlated, but  
264 completely independent of each other. The distribution is not yet assumed. The  
265 third component is the variance process or the expression for the volatility. The  
266 variance is given by a constant  $\omega$ , plus the random part which depends on the return  
267 shock of the previous period squared ( $\varepsilon_{t-1}^2$ ). In that sense when the uncertainty  
268 or surprise in the last period increases, then the variance becomes larger in the  
269 next period. The element  $\sigma_t^2$  is thus known at time  $t - 1$ , while it is a deterministic  
270 function of a random variable observed at time  $t - 1$  (i.e.  $\varepsilon_{t-1}^2$ ).

$$y_t = \mu + \varepsilon_t \tag{1.2}$$

$$\varepsilon_t = \sigma_t \times z_t, \text{ where } z_t \stackrel{iid}{\sim} (0, 1) \tag{1.3}$$

$$\sigma_t^2 = \omega + \alpha_1 \times \varepsilon_{t-1}^2 \tag{1.4}$$



### 1.3. Volatility modeling

From these components we could look at the conditional moments (or expected returns and variance). We can plug in the component  $\sigma_t$  into the conditional mean innovation  $\varepsilon_t$  and use the conditional mean innovation to examine the conditional mean return. In equation (1.5) and (1.6) they are derived. Because the random variable  $z_t$  is distributed with a zero-mean, the conditional expectation is 0. As a consequence, the conditional mean return in equation (1.6) is equal to the unconditional mean in the most simple case. But variations are possible using ARMA (eg. AR(1)) processes.

$$\mathbb{E}_{t-1}(\varepsilon_t) = \mathbb{E}_{t-1}(\sqrt{\omega + \alpha_1 \times \varepsilon_{t-1}^2} \times z_t) = \sigma_t \mathbb{E}_{t-1}(z_t) = 0 \quad (1.5)$$

$$\mathbb{E}_{t-1}(y_t) = \mu + \mathbb{E}_{t-1}(\varepsilon_t) = \mu \quad (1.6)$$

For the conditional variance, knowing everything that happened until and including period  $t - 1$  the conditional innovation variance is given by equation (1.7). This is equal to  $\sigma_t^2$ , while the variance of  $z_t$  is equal to 1. Then it is easy to derive the conditional variance of returns in equation (1.8), that is why equation (1.4) is called the variance equation.

$$var_{t-1}(\varepsilon_t) = \mathbb{E}_{t-1}(\varepsilon_t^2) = \mathbb{E}_{t-1}(\sigma_t^2 \times z_t^2) = \sigma_t^2 \mathbb{E}_{t-1}(z_t^2) = \sigma_t^2 \quad (1.7)$$

$$var_{t-1}(y_t) = var_{t-1}(\varepsilon_t) = \sigma_t^2 \quad (1.8)$$

The unconditional variance is also interesting to derive, while this is the long-run variance, which will be derived in (1.12). After deriving this using the law of iterated expectations and assuming stationarity for the variance process, one would get (1.9) for the unconditional variance, equal to the constant  $c$  and divided by  $1 - \alpha_1$ , the slope of the variance equation.

## 1. Literature review

$$\sigma^2 = \frac{\omega}{1 - \alpha_1} \quad (1.9)$$

289 This leads to the properties of ARCH models. - Stationarity condition for variance:  
290  $\omega > 0$  and  $0 \leq \alpha_1 < 1$ .

- 291 • Zero-mean innovations
- 292 • Uncorrelated innovations

293 Thus a weak white noise process  $\varepsilon_t$ .

294 Stationarity implies that the series on which the ARCH model is used does  
295 not have any trend and has a constant expected mean. Only the conditional  
296 variance is changing.

297 The unconditional 4th moment, kurtosis  $\mathbb{E}(\varepsilon_t^4)/\sigma^4$  of an ARCH model is given  
298 by equation (1.10). This term is larger than 3, which implicates that the fat-  
299 tails (a stylized fact of returns).

$$3 \frac{1 - \alpha_1^2}{1 - 3\alpha_1^2} \quad (1.10)$$

300 Another property of ARCH models is that it takes into account volatility clustering.  
301 Because we know that  $\text{var}(\varepsilon_t) = \mathbb{E}(\varepsilon_t^2) = \sigma^2 = \omega/(1 - \alpha_1)$ , we can plug in  $\omega$   
302 for the conditional variance  $\text{var}_t(\varepsilon_{t+1}) = \mathbb{E}(\varepsilon_{t+1}^2) = \sigma_{t+1}^2 = c + \alpha_1 \times \varepsilon_t^2$ . Thus it  
303 follows that equation (1.11) displays volatility clustering. If we examine the RHS,  
304 as  $\alpha_1 > 0$  (condition for stationarity), when shock  $\varepsilon_t^2$  is larger than what you  
305 expect it to be on average  $\sigma^2$  the LHS will also be positive. Then the conditional  
306 variance will be larger than the unconditional variance. Briefly, large shocks will  
307 be followed by more large shocks.

$$\sigma_{t+1}^2 - \sigma^2 = \alpha_1 \times (\varepsilon_t^2 - \sigma^2) \quad (1.11)$$

Excess kurtosis can be modeled, even when the conditional distribution is assumed to be normally distributed. The third moment, skewness, can be introduced using a skewed conditional distribution as we saw in part A.1. The serial correlation for squared innovations is positive if fourth moment exists (equation (1.10), this is volatility clustering once again.

The estimation of ARCH model and in a next step GARCH models will be explained in the methodology. However how will then the variance be forecasted? Well, the conditional variance for the  $k$ -periods ahead, denoted as period  $T + k$ , is given by equation (1.12). This can already be simplified, while we know that  $\sigma_{T+1}^2 = \omega + \alpha_1 \times \varepsilon_T^2$  from equation (1.4).

$$\begin{aligned}\mathbb{E}_T(\varepsilon_{T+k}^2) &= \omega \times (1 + \alpha_1 + \dots + \alpha^{k-2}) + \alpha^{k-1} \times \sigma_{T+1}^2 \\ &= \omega \times (1 + \alpha_1 + \dots + \alpha^{k-1}) + \alpha^k \times \sigma_T^2\end{aligned}\tag{1.12}$$

It can be shown that then the conditional variance in period  $T+k$  is equal to equation (1.13). The LHS is the predicted conditional variance  $k$ -periods ahead above its unconditional variance,  $\sigma^2$ . The RHS is the difference current last-observed return residual  $\varepsilon_T^2$  above the unconditional average multiplied by  $\alpha_1^k$ , a decreasing function of  $k$  (given that  $0 \leq \alpha_1 < 1$ ). The further ahead predicting the variance, the closer  $\alpha_1^k$  comes to zero, the closer to the unconditional variance, i.e. the long-run variance.

$$\mathbb{E}_T(\varepsilon_{T+k}^2) - \sigma^2 = \alpha_1^k \times (\varepsilon_T^2 - \sigma^2)\tag{1.13}$$

### 1.3.3 Univariate GARCH models

An improvement on the ARCH model is the Generalised Autoregressive Conditional Heteroscedasticity (GARCH). This model and its variants come in to play because of the fact that calculating standard deviations through rolling periods, gives an equal weight to distant and nearby periods, by such not taking into account empirical evidence of volatility clustering, which can be identified as positive autocorrelation in the absolute returns. GARCH models are an extension to ARCH models, as they

## 1. Literature review

incorporate both a novel moving average term (not included in ARCH) and the autoregressive component.

An overview (of a selection) of investigated GARCH models is given in the following table.

**Table 1.1:** GARCH models, the founders

Author(s)/user(s)	Model
Engle (1982)	ARCH model
Bollerslev (1986)	GARCH model
Bollerslev (1986)	IGARCH model
Nelson (1991)	EGARCH model
Glosten et al. (1993)	GJRGARCH model
Engle and Ng (1993)	NAGARCH model
Zakoian (1994)	TGARCH model
Taylor (1986) and Schwert (1989)	TSGARCH (or AVGARCH) model
Morgan Guaranty Trust Company (1996)	EWMA or RiskMetrics model

## 1.4 ACD models

An extension to GARCH models was proposed by Hansen (1994), the autoregressive conditional density estimation model (referred to as ACD models, sometimes ARCD). It focuses on time variation in higher moments (skewness and kurtosis), because the degree and frequency of extreme events seem to be not expected by traditional models. Some GARCH models are already able to capture the dynamics by relying on a different unconditional distribution than the normal distribution (for example skewed distributions like the SGED, SGT), or a model that allows to model these higher moments. However, Ghalanos (2016) mentions that these models also assume the shape and skewness parameters to be constant (not time varying). As Ghalanos mentions: “the research on time varying higher moments has mostly explored different parameterizations in terms of dynamics and distributions with little attention to the performance of the models out-of-sample and ability to outperform a GARCH model with respect to VaR.” Also one could question the marginal benefits of the ACD, while the estimation procedure is not simple (nonlinear bounding specification of higher moment distribution parameters and interaction). So, are skewness (skewness parameter) and kurtosis (shape parameters) time varying? The literature investigating higher moments has arguments for and against this statement. In part 2.2.2 the specification is given.

## 355 1.5 Value at Risk

356 Value-at-Risk (VaR) is a risk metric developed simultaneously by [Markowitz1952]  
 357 and [Roy1952] to calculate how much money an investment, portfolio, department  
 358 or institution such as a bank could lose in a market downturn, though in this period  
 359 it remained mostly a theoretical discussion due to lacking processing power and  
 360 industry demand for risk management measures. According to [Holton2002] VaR  
 361 gained traction in the last decade of the 20th century when financial institutions  
 362 started using it to determine their regulatory capital requirements. A  $VaR_{99}$  finds  
 363 the amount that would be the greatest possible loss in 99% of cases. It can be  
 364 defined as the threshold value  $\theta_t$ . Put differently, in 1% of cases the loss would  
 365 be greater than this amount. It is specified as in (1.14). [Christofferson2001] puts  
 366 forth a general framework for specifying VaR models and comparing between  
 367 two alternatives models.

$$Pr(y_t \leq \theta_t | \Omega_{t-1}) \equiv \phi \quad (1.14)$$

368 With  $y_t$  expected returns in period  $t$ ,  $\Omega_{t-1}$  the information set available in the  
 369 previous period and  $\phi$  the chosen confidence level.

## 370 1.6 Conditional Value at Risk

371 One major shortcoming of the VaR is that it does not provide information on  
 372 the probability distribution of losses beyond the threshold amount. As VaR lacks  
 373 subadditivity of different percentile outcomes, [Artzner1998] reject it as a coherent  
 374 measure of risk. This is problematic, as losses beyond this amount would be more  
 375 problematic if there is a large probability distribution of extreme losses, than if  
 376 losses follow say a normal distribution. To solve this issue, they provide a conceptual  
 377 idea of a conditional VaR (cVaR) which quantifies the average loss one would expect  
 378 if the threshold is breached, thereby taking the distribution of the tail into account.  
 379 Mathematically, a  $cVaR_{99}$  is the average of all the  $VaR$  with a confidence level

## 1.6. Conditional Value at Risk

equal to or higher than 99. It is commonly referred to as expected shortfall (ES) sometimes and was written out in the form it is used by today by (Bertsimas et al. 2004). It is specified as in (1.15).

To calculate  $\theta_t$ , VaR and cVaR require information on the expected distribution mean, variance and other parameters, to be calculated using the previously discussed GARCH models and distributions.

$$Pr(y_t \leq \theta_t | \Omega_{t-1}) \equiv \int_{-\infty}^{\theta_t} f(y_t | \Omega_{t-1}) dy_t = \phi \quad (1.15)$$

With the same notations as before, and  $f$  the (conditional) probability density function of  $y_t$ .

According to the BIS framework, banks need to calculate both  $VaR_{99}$  and  $VaR_{97.5}$  daily to determine capital requirements for equity, using a minimum of one year of daily observations (Basel Committee on Banking Supervision 2016). Whenever a daily loss is recorded, this has to be registered as an exception. Banks can use an internal model to calculate their VaRs, but if they have more than 12 exceptions for their  $VaR_{99}$  or 30 exceptions for their  $VaR_{97.5}$  they have to follow a standardized approach. Similarly, banks must calculate  $cVaR_{97.5}$ .

## 395 1.7 Past literature on the consequences of higher 396 moments for VaR determination

397 Here comes the discussion about studies that have looked at higher moments and  
398 VaR determination. Also a summary of studies that discusses time-varying higher  
399 moments, but not a big part, while it is also only a small part of the empirical  
400 findings (couple of GARCH-ACD models).

**Table 1.2:** Higher moments and VaR

Author	Higher moments
Hansen (1994)	
\@harvey1999	

Brooks et al. (2005)



# 2

## Data and methodology

### 2.1 Data

We worked with daily returns on the EURO STOXX 50 Index denoted in EUR from 31 December, 1986 to 27 April, 2021. It is the leading blue-chip index of the Eurozone, was founded in 1999 and covers 50 of the most liquid and largest (in terms of free-float market capitalization) stocks. Its composition is reviewed annually in September, from each of the 19 EURO STOXX Supersector indices the biggest stocks are selected until the coverage is at 60% of the free-float market cap of each of the EURO STOXX Supersector index then all the current EURO STOXX 50 stocks are used in the selection list from which the largest 40 in terms of free-float market cap are selected and the remaining 10 stocks are chosen among those ranked between 41 and 60 (*Calculation guide STOXX* ® 2020). The calculation of the index is made with the (2.1), that measures the changes in price of the index for fixed weights.

$$\text{Index}_t = \frac{\sum_{i=1}^n (p_{it} \cdot s_{it} \cdot f_{it} \cdot c_{it} \cdot x_{it})}{D_t} = \frac{M_t}{D_t} \quad (2.1)$$

where:  $t$  = Time the index is computed  $n$  = Number of companies in the index  
 $p_{it}$  = Price of company ( $i$ ) at time ( $t$ )  $s_{it}$  = Number of shares of company ( $i$ ) at

## 2. Data and methodology

time (t)  $ff_{it}$  = Free float factor of company (i) at time (t)  $cf_{it}$  = Weighting cap  
factor of company (i) at time (t)  $x_{it}$  = Exchange rate from local currency into  
index currency for company (i) at time (t)  $M_t$  = Free-float market capitalization  
of the index at time (t)  $D_t$  = Divisor of the index at time (t)  
Changes in weights caused by corporate actions are proportionally distributed  
across the components of the index and the index Divisor is computed with  
the (2.2) formula.

$$D_{t+1} = D_t \cdot \frac{\sum_{i=1}^n (p_{it} \cdot s_{it} \cdot ff_{it} \cdot cf_{it} \cdot x_{it}) \pm \Delta MC_{t+1}}{\sum_{i=1}^n (p_{it} \cdot s_{it} \cdot ff_{it} \cdot cf_{it} \cdot x_{it})} \quad (2.2)$$

where:  $\Delta MC_{t+1}$  = Difference between the closing market capitalization of the index  
and the adjusted closing market capitalization of the index  
(Optional)

The same analysis has been performed for the INDEX 1, INDEX 2, INDEX  
3 and the INDEX 4 indexes with not different conclusions. The findings of these  
researches are available upon requests.

### 2.1.1 Descriptives

#### Table of summary statistics

Equation 2.1 provides the main statistics describing the return series analyzed.  
Returns are computed with equation (2.3).

$$R_t = 100 (\ln(I_t) - \ln(I_{t-1})) \quad (2.3)$$

where  $I_t$  is the index price at time  $t$  and  $I_{t-1}$  is the index price at  $t - 1$ .  
The arithmetic mean of the series is 0.017% with a standard deviation of 1.307%  
and a median of 0.036 which translate to an annualized mean of 4.208% and  
an annualized standard deviation of 20.748%. The skewness statistic is highly  
significant and negative at -0.31 and the excess kurtosis is also highly significant  
and positive at 7.208. These 2 statistics give an overview of the distribution of the

returns which has thicker tails than the normal distribution with a higher presence of left tail observations. A formal test such as the Jarque-Bera one with its statistic at 19528.62 and a high statistical significance, confirms the non normality feeling given by the Skewness and Kurtosis.

The right column of table 2.1 displays the same descriptive statistics but for the standardized residuals obtained from a simple GARCH model as mentioned in table 2.1 in Note 2\*. Again, Skewness statistic at -0.633 with a high statistical significance level and the excess Kurtosis at 5.134 also with a high statistical significance, suggest a non normal distribution of the standardized residuals and the Jarque-Bera statistic at NA, given its high significance, confirms the rejection of the normality assumption.

## Descriptive figures

### Stylized facts

As can be seen in figure 2.1 the Euro area equity and later, since 1999 the EuroStoxx 50 went up during the tech (“dot com”) bubble reaching an ATH of €5464.43. Then, there was a correction to boom again until the burst of the 2008 financial crisis. After which it decreased significantly. With an ATL at 09 March, 2009 of €1809.98. There is an improvement, but then the European debt crisis, with its peak in 2010-2012, occurred. From then there was some improvement until the “health crisis”, which arrived in Europe, February 2020. This crisis recovered very quickly reaching already values higher than the pre-COVID crisis level.

## 2. Data and methodology

**Table 2.1:** Summary statistics of the returns

Statistics	Eurostoxx.50	Standardized.Residuals
Arithmetic Mean	0.0167	-0.0409
Median	0.0357	-0.0193
Maximum	10.4376	5.7126
Minimum	-13.2404	-11.7732
Stdev	1.307	0.9992
Skewness	-0.31 (0***)	-0.6327 (0***)
Excess Kurtosis	7.2083 (0***)	5.134 (0***)
Jarque-Bera	19528.6196***	10431.0514***

Notes

<sup>1</sup> This table shows the descriptive statistics of the daily percentage returns of EURO STOXX 50 over the period 1987-01-01 to 2021-04-27 (8954 observations). Including arithmetic mean, median, maximum, minimum, standard deviation, skewness, excess kurtosis and the Jarque-Bera test.

<sup>2</sup> The standardized residual is derived from a maximum likelihood estimation (simple GARCH model) as follows:


$$R_t = \alpha_0 + \alpha_1 R_{t-1} + z_t \sigma_t$$

$$\sigma_t^2 = \beta_0 + \beta_1 \sigma_{t-1}^2 z_{t-1}^2 + \beta_2 \sigma_{t-1}^2,$$

Where  $z$  is the standard residual (assumed to have a normal distribution).

<sup>3</sup> \*, \*\*, \*\*\* represent significance levels at the 5


463 In figure 2.2 the daily log-returns are visualized. A stylized fact that is observable  
464 is the volatility clustering. As can be seen: periods of large volatility are mostly  
465 followed by large volatility and small volatility by small volatility.



**Figure 2.1:** Eurostoxx 50 Price Index prices

466 In figure 2.4 the density distribution of the log returns are examined. As can  
 467 be seen, as already mentioned in part 1.1, log returns are not really normally  
 468 distributed. So


## 2. Data and methodology




**Figure 2.2:** Eurostoxx 50 Price Index log returns

469

ACF plots: to do...




**Figure 2.3:** Eurostoxx 50 rolling volatility (22 days, calculated over 252 days)



**Figure 2.4:** Density vs. Normal Eurostoxx 50 log returns)

## 2. Data and methodology



**Figure 2.5:** Absolute prediction errors



## 2.2 Methodology

### 2.2.1 Garch models

As already mentioned in part 1.3.3, GARCH models GARCH, EGARCH, IGARCH, GJRGARCH, NGARCH, TGARCH and NAGARCH (or TSGARCH) will be estimated. Additionally the distributions will be examined as well, including the normal, student-t distribution, skewed student-t distribution, generalized error distribution, skewed generalized error distribution and the skewed generalized t distribution.

They will be estimated using maximum likelihood. As already mentioned, fortunately, Ghalanos (2020b) has made it easy for us to implement this methodology in the R language (v.3.6.1) with the package “rugarch” v.1.4-4 (*R univariate garch*), which gives us a bit more time to focus on the results and the interpretation.

Maximum likelihood estimation is a method to find the distribution parameters that best fit the observed data, through maximization of the likelihood function, or the computationally more efficient log-likelihood function (by taking the natural logarithm). It is assumed that the return data is i.i.d. and that there is some underlying parametrized density function  $f$  with one or more parameters that generate the data, defined as a vector  $\theta$  (equation (2.5)). These functions are based on the joint probability distribution of the observed data (equation (2.7)). Subsequently, the (log)likelihood function is maximized using an optimization algorithm (equation (2.9)).

$$y_1, y_2, \dots, y_N \sim i.i.d \quad (2.4)$$

$$y_i \sim f(y|\theta) \quad (2.5)$$

## 2. Data and methodology

$$L(\theta) = \prod_{i=1}^N f(y_i|\theta) \quad (2.6)$$

$$\log(L(\theta)) = \sum_{i=1}^N \log f(y_i|\theta) \quad (2.7)$$

$$\theta^* = \arg \max_{\theta} [L] \quad (2.8)$$

$$\theta^* = \arg \max_{\theta} [\log(L)] \quad (2.9)$$

### 493 2.2.2 ACD models

494 Following Ghalanos (2016), arguments of ACD models are specified as in Hansen  
 495 (1994). The density function  $f(y|\alpha)$  has parameters  $\alpha_t = (\mu_t, \sigma_t, \nu_t)$ , with equation  
 496 (2.10), the conditional mean equation. Equation (2.11) as the conditional variance.  
 497 And  $\nu_t = \nu(\theta, x_t)$  the remaining parameters of the distribution like the skewness  
 498 and kurtosis (shape) parameters.

$$\mu_t = \mu(\theta, x_t) = E(y_t | x_t) \quad (2.10)$$

$$\sigma_t^2 = \sigma^2(\theta, x_t) = E((y_t - \mu_t)^2 | x_t) \quad (2.11)$$

499 To further explain the difference between GARCH and ACD. The scaled  
 500 innovations are given by equation (2.12). The conditional density is given by  
 501 equation (2.13) and related to the density function  $f(y|\alpha)$  as in equation (2.2.2).

$$z_t(\theta) = \frac{y_t - \mu(\theta, x_t)}{\sigma(\theta, x_t)} \quad (2.12)$$

$$g(z | \eta_t) = \frac{d}{dz} P(z_t < z | \eta_t) \quad (2.13)$$

$$f(y_t | \mu_t, \sigma_t^2, \eta_t) = \frac{1}{\sigma_t} g(z_t | \eta_t) \quad (2.14)$$

502

503 Again Ghalanos (2016) makes it easier to implement the somewhat complex  
504 ACD models using the R language with package “racd”.

### 505 **2.2.3 Analysis Tests VaR and cVaR**

#### 506 **Unconditional coverage test of Kupiec (1995)**

507 A number of tests are computed to see if the value-at-risk estimations capture  
508 the actual losses well. A first one is the unconditional coverage test by Kupiec (1995).  
509 The unconditional coverage or proportion of failures method tests if the actual  
510 value-at-risk exceedances are consistent with the expected exceedances (a chosen  
511 percentile, e.g. 1% percentile) of the VaR model. Following Kupiec (1995) and  
512 Ghalanos (2020a), the number of exceedence follow a binomial distribution (with  
513 thus probability equal to the significance level or expected proportion) under the  
514 null hypothesis of a correct VaR model. The test is conducted as a likelihood ratio  
515 test with statistic like in equation (2.15), with  $p$  the probability of an exceedence  
516 for a confidence level,  $N$  the sample size and  $X$  the number of exceedence. The  
517 null hypothesis states that the test statistic  $LR^{uc}$  is  $\chi^2$ -distributed with one degree  
518 of freedom or that the probability of failure  $\hat{p}$  is equal to the chosen percentile  $\alpha$ .

$$LR^{uc} = -2 \ln \left( \frac{(1-p)^{N-X} p^X}{\left(1 - \frac{X}{N}\right)^{N-X} \left(\frac{X}{N}\right)^X} \right) \quad (2.15)$$

#### 519 **Conditional coverage test of Christoffersen et al. (2001)**

520 Christoffersen et al. (2001) proposed the conditional coverage test. It is tests for  
521 unconditional covrage and serial independence. The serial independence is important  
522 while the  $LR^{uc}$  can give a false picture while at any point in time it classifies

## 2. Data and methodology

523 inaccurate VaR estimates as “acceptably accurate” (Bali and Theodossiou 2007). For  
524 a certain VaR estimate an indicator variable,  $I_t(\alpha)$ , is computed as equation (2.16).

$$I_t(\alpha) = \begin{cases} 1 & \text{if exceedence occurs} \\ 0 & \text{if no exceedence occurs} \end{cases} . \quad (2.16)$$

525 It involves a likelihood ratio test’s null hypothesis is that the statistic is  $\chi^2$ -  
526 distributed with two degrees of freedom or that the probability of violation  $\hat{p}$   
527 (unconditional coverage) as well as the conditional coverage (independence) is  
528 equal to the chosen percentile  $\alpha$ .

### 529 **Dynamic quantile test**

530 Engle and Manganelli (1999) with the aim to provide completeness to the  
531 conditional coverage test of Christoffersen et al. (2001) developed the Dynamic  
532 quantile test. It consists in testing some restriction in a ... (work-in-progress).

# 3

## Empirical Findings


### 3.1 Density of the returns

#### 3.1.1 MLE distribution parameters

In table 3.1 we can see... Additionally, for every distribution fitted with MLE,

plots are generated to compare the theoretical distribution with the observed returns.

### 3. Empirical Findings



539

**Table 3.1:** Maximum likelihood estimates of unconditional distribution functions

	$\mu$	$\sigma$	$\lambda$	$p$	$q$	$\nu$	$L$
SGT	0.02 (0.013)	1.321 (0.026)**	-0.04 (0.012)**	1.381 (0.071)**	3.317 (0.534)**		-13973.01 27
SGED	0.02 (0.01)	1.274 (0.016)**	-0.018 (0.008)*	0.918 (0.016)**	Inf		-14008.18 27
GED	0.032 (0.005)**	1.276 (0.016)**	0	0.913 (0.016)**	Inf		-14009.09 28
ST	0.019 (0.014)**	1.487 (0.056)**	0.949 (0.013)**			2.785 (0.1)**	-13997.35 28
T	0.056 (0.01)**	1.494 (0.056)**				2.765 (0.097)**	-14005.14 28
Normal	0.017 (0.014)	1.304 (0.015)**	0	2	Inf		-15093.32 30

Notes

## 3.2 Results of GARCH with constant higher moments

```
distributions <- c("norm", "std", "sstd", "ged", "sged")

#garchspec <- garchfit <- garchforecast <- stdret <- vector(mode = "list", length =
#names(garchspec) <- names(garchfit) <- names(garchforecast) <- names(stdret) <- di
Models.garch <- c("sGARCH", "eGARCH", "fGARCH.AVGARCH", "fGARCH.NAGARCH", "gjrGARCH", "

for(i in 1:length(Models.garch)){
  assign(paste0("garchspec.", Models.garch[i]), vector(mode = "list", length = length(dis
  assign(paste0("garchfit.", Models.garch[i]), vector(mode = "list", length = length(dist
  assign(paste0("stdret.", Models.garch[i]), vector(mode = "list", length = length(distri
}

# ls(pattern = "garchspec.")
# sapply(ls(pattern = "garchspec."), FUN = setNames, distributions)

# .sGARCH-----
for(i in 1:length(distributions)){
  # Specify a GARCH model with constant mean
  garchspec.sGARCH[[i]] <- ugarchspec(mean.model = list(armaOrder = c(1,0)),
                                     variance.model = list(model = "sGARCH", garchOrder = c(1,1), var
                                     distribution.model = distributions[i])

  # Estimate the model
  garchfit.sGARCH[[i]] <- ugarchfit(data = R, spec = garchspec.sGARCH[[i]])
  # Compute stdret using residuals()
  stdret.sGARCH[[i]] <- residuals(garchfit.sGARCH[[i]], standardize = TRUE)
}

# .eGARCH-----
```

### 3. Empirical Findings

```
for(i in 1:length(distributions)){
  # Specify a GARCH model with constant mean
  garchspec.eGARCH[[i]] <- ugarchspec(mean.model = list(armaOrder = c(1,0)),
                                     variance.model = list(model = "eGARCH", variance.targeting = TRUE),
                                     distribution.model = distributions[i])
  # Estimate the model
  garchfit.eGARCH[[i]] <- ugarchfit(data = R, spec = garchspec.eGARCH[[i]])
  # Compute stdret using residuals()
  stdret.eGARCH[[i]] <- residuals(garchfit.eGARCH[[i]], standardize = TRUE)
}

# .fGARCH.NAGARCH-----
for(i in 1:length(distributions)){
  # Specify a GARCH model with constant mean
  garchspec.fGARCH.NAGARCH[[i]] <- ugarchspec(mean.model = list(armaOrder = c(1,0)),
                                               variance.model = list(model = "fGARCH", submodel = "NAGARCH"),
                                               distribution.model = distributions[i])
  # Estimate the model
  garchfit.fGARCH.NAGARCH[[i]] <- ugarchfit(data = R, spec = garchspec.fGARCH.NAGARCH[[i]])
  # Compute stdret using residuals()
  stdret.fGARCH.NAGARCH[[i]] <- residuals(garchfit.fGARCH.NAGARCH[[i]], standardize = TRUE)
}

# .fGARCH.AVGARCH-----
for(i in 1:length(distributions)){
  # Specify a GARCH model with constant mean
  garchspec.fGARCH.AVGARCH[[i]] <- ugarchspec(mean.model = list(armaOrder = c(1,0)),
                                               variance.model = list(model = "fGARCH", submodel = "AVGARCH"),
                                               distribution.model = distributions[i])
  # Estimate the model
```



### 3.2. Results of GARCH with constant higher moments

```
garchfit.fGARCH.AVGARCH[[i]] <- ugarchfit(data = R, spec = garchspec.fGARCH.AVGARCH[[i]]
# Compute stdret using residuals()
stdret.fGARCH.AVGARCH[[i]] <- residuals(garchfit.fGARCH.AVGARCH[[i]], standardize = T
}

# .gjrGARCH-----
for(i in 1:length(distributions)){
# Specify a GARCH model with constant mean
garchspec.gjrGARCH[[i]] <- ugarchspec(mean.model = list(armaOrder = c(1,0)),
      variance.model = list(model = "gjrGARCH", variance.targeting = F
      distribution.model = distributions[i])
# Estimate the model
garchfit.gjrGARCH[[i]] <- ugarchfit(data = R, spec = garchspec.gjrGARCH[[i]])
# Compute stdret using residuals()
stdret.gjrGARCH[[i]] <- residuals(garchfit.gjrGARCH[[i]], standardize = TRUE)
}

# fGARCH.TGARCH-----
for(i in 1:length(distributions)){
# Specify a GARCH model with constant mean
garchspec.fGARCH.TGARCH[[i]] <- ugarchspec(mean.model = list(armaOrder = c(1,0)),
      variance.model = list(model = "fGARCH", submodel = "TGARCH", var
      distribution.model = distributions[i])
# Estimate the model
garchfit.fGARCH.TGARCH[[i]] <- ugarchfit(data = R, spec = garchspec.fGARCH.TGARCH[[i]]
# Compute stdret using residuals()
stdret.fGARCH.TGARCH[[i]] <- residuals(garchfit.fGARCH.TGARCH[[i]], standardize = TRU
}

# .iGARCH-----
```

### 3. Empirical Findings

```
for(i in 1:length(distributions)){
# Specify a GARCH model with constant mean
garchspec.iGARCH[[i]] <- ugarchspec(mean.model = list(armaOrder = c(1,0)),
                                     variance.model = list(model = "iGARCH", variance.targeting = TRUE),
                                     distribution.model = distributions[i])
# Estimate the model
garchfit.iGARCH[[i]] <- ugarchfit(data = R, spec = garchspec.iGARCH[[i]])
# Compute stdret using residuals()
stdret.iGARCH[[i]] <- residuals(garchfit.iGARCH[[i]], standardize = TRUE)
}

# .csGARCH-----
# for(i in 1:length(distributions)){
# # Specify a GARCH model with constant mean
# garchspec.csGARCH[[i]] <- ugarchspec(mean.model = list(armaOrder = c(1,0)),
#                                       variance.model = list(model = "csGARCH", variance.targeting = TRUE),
#                                       distribution.model = distributions[i])
# # Estimate the model
# garchfit.csGARCH[[i]] <- ugarchfit(data = R, spec = garchspec.csGARCH[[i]])
# # Compute stdret using residuals()
# stdret.csGARCH[[i]] <- residuals(garchfit.csGARCH[[i]], standardize = TRUE)
# }

# we need EWMA
for(i in 1:length(distributions)){
# Specify a GARCH model with constant mean
garchspec.EWMA[[i]] <- ugarchspec(mean.model = list(armaOrder = c(1,0)),
                                   variance.model = list(model = "iGARCH", variance.targeting = TRUE),
                                   distribution.model = distributions[i], fixed.pars = list(omega = 0.000001, alpha1 = 0.999999, alpha2 = 0.000001, alpha3 = 0.000001, alpha4 = 0.000001, alpha5 = 0.000001, alpha6 = 0.000001, alpha7 = 0.000001, alpha8 = 0.000001, alpha9 = 0.000001, alpha10 = 0.000001, alpha11 = 0.000001, alpha12 = 0.000001, alpha13 = 0.000001, alpha14 = 0.000001, alpha15 = 0.000001, alpha16 = 0.000001, alpha17 = 0.000001, alpha18 = 0.000001, alpha19 = 0.000001, alpha20 = 0.000001, alpha21 = 0.000001, alpha22 = 0.000001, alpha23 = 0.000001, alpha24 = 0.000001, alpha25 = 0.000001, alpha26 = 0.000001, alpha27 = 0.000001, alpha28 = 0.000001, alpha29 = 0.000001, alpha30 = 0.000001, alpha31 = 0.000001, alpha32 = 0.000001, alpha33 = 0.000001, alpha34 = 0.000001, alpha35 = 0.000001, alpha36 = 0.000001, alpha37 = 0.000001, alpha38 = 0.000001, alpha39 = 0.000001, alpha40 = 0.000001, alpha41 = 0.000001, alpha42 = 0.000001, alpha43 = 0.000001, alpha44 = 0.000001, alpha45 = 0.000001, alpha46 = 0.000001, alpha47 = 0.000001, alpha48 = 0.000001, alpha49 = 0.000001, alpha50 = 0.000001, alpha51 = 0.000001, alpha52 = 0.000001, alpha53 = 0.000001, alpha54 = 0.000001, alpha55 = 0.000001, alpha56 = 0.000001, alpha57 = 0.000001, alpha58 = 0.000001, alpha59 = 0.000001, alpha60 = 0.000001, alpha61 = 0.000001, alpha62 = 0.000001, alpha63 = 0.000001, alpha64 = 0.000001, alpha65 = 0.000001, alpha66 = 0.000001, alpha67 = 0.000001, alpha68 = 0.000001, alpha69 = 0.000001, alpha70 = 0.000001, alpha71 = 0.000001, alpha72 = 0.000001, alpha73 = 0.000001, alpha74 = 0.000001, alpha75 = 0.000001, alpha76 = 0.000001, alpha77 = 0.000001, alpha78 = 0.000001, alpha79 = 0.000001, alpha80 = 0.000001, alpha81 = 0.000001, alpha82 = 0.000001, alpha83 = 0.000001, alpha84 = 0.000001, alpha85 = 0.000001, alpha86 = 0.000001, alpha87 = 0.000001, alpha88 = 0.000001, alpha89 = 0.000001, alpha90 = 0.000001, alpha91 = 0.000001, alpha92 = 0.000001, alpha93 = 0.000001, alpha94 = 0.000001, alpha95 = 0.000001, alpha96 = 0.000001, alpha97 = 0.000001, alpha98 = 0.000001, alpha99 = 0.000001, alpha100 = 0.000001, alpha101 = 0.000001, alpha102 = 0.000001, alpha103 = 0.000001, alpha104 = 0.000001, alpha105 = 0.000001, alpha106 = 0.000001, alpha107 = 0.000001, alpha108 = 0.000001, alpha109 = 0.000001, alpha110 = 0.000001, alpha111 = 0.000001, alpha112 = 0.000001, alpha113 = 0.000001, alpha114 = 0.000001, alpha115 = 0.000001, alpha116 = 0.000001, alpha117 = 0.000001, alpha118 = 0.000001, alpha119 = 0.000001, alpha120 = 0.000001, alpha121 = 0.000001, alpha122 = 0.000001, alpha123 = 0.000001, alpha124 = 0.000001, alpha125 = 0.000001, alpha126 = 0.000001, alpha127 = 0.000001, alpha128 = 0.000001, alpha129 = 0.000001, alpha130 = 0.000001, alpha131 = 0.000001, alpha132 = 0.000001, alpha133 = 0.000001, alpha134 = 0.000001, alpha135 = 0.000001, alpha136 = 0.000001, alpha137 = 0.000001, alpha138 = 0.000001, alpha139 = 0.000001, alpha140 = 0.000001, alpha141 = 0.000001, alpha142 = 0.000001, alpha143 = 0.000001, alpha144 = 0.000001, alpha145 = 0.000001, alpha146 = 0.000001, alpha147 = 0.000001, alpha148 = 0.000001, alpha149 = 0.000001, alpha150 = 0.000001, alpha151 = 0.000001, alpha152 = 0.000001, alpha153 = 0.000001, alpha154 = 0.000001, alpha155 = 0.000001, alpha156 = 0.000001, alpha157 = 0.000001, alpha158 = 0.000001, alpha159 = 0.000001, alpha160 = 0.000001, alpha161 = 0.000001, alpha162 = 0.000001, alpha163 = 0.000001, alpha164 = 0.000001, alpha165 = 0.000001, alpha166 = 0.000001, alpha167 = 0.000001, alpha168 = 0.000001, alpha169 = 0.000001, alpha170 = 0.000001, alpha171 = 0.000001, alpha172 = 0.000001, alpha173 = 0.000001, alpha174 = 0.000001, alpha175 = 0.000001, alpha176 = 0.000001, alpha177 = 0.000001, alpha178 = 0.000001, alpha179 = 0.000001, alpha180 = 0.000001, alpha181 = 0.000001, alpha182 = 0.000001, alpha183 = 0.000001, alpha184 = 0.000001, alpha185 = 0.000001, alpha186 = 0.000001, alpha187 = 0.000001, alpha188 = 0.000001, alpha189 = 0.000001, alpha190 = 0.000001, alpha191 = 0.000001, alpha192 = 0.000001, alpha193 = 0.000001, alpha194 = 0.000001, alpha195 = 0.000001, alpha196 = 0.000001, alpha197 = 0.000001, alpha198 = 0.000001, alpha199 = 0.000001, alpha200 = 0.000001, alpha201 = 0.000001, alpha202 = 0.000001, alpha203 = 0.000001, alpha204 = 0.000001, alpha205 = 0.000001, alpha206 = 0.000001, alpha207 = 0.000001, alpha208 = 0.000001, alpha209 = 0.000001, alpha210 = 0.000001, alpha211 = 0.000001, alpha212 = 0.000001, alpha213 = 0.000001, alpha214 = 0.000001, alpha215 = 0.000001, alpha216 = 0.000001, alpha217 = 0.000001, alpha218 = 0.000001, alpha219 = 0.000001, alpha220 = 0.000001, alpha221 = 0.000001, alpha222 = 0.000001, alpha223 = 0.000001, alpha224 = 0.000001, alpha225 = 0.000001, alpha226 = 0.000001, alpha227 = 0.000001, alpha228 = 0.000001, alpha229 = 0.000001, alpha230 = 0.000001, alpha231 = 0.000001, alpha232 = 0.000001, alpha233 = 0.000001, alpha234 = 0.000001, alpha235 = 0.000001, alpha236 = 0.000001, alpha237 = 0.000001, alpha238 = 0.000001, alpha239 = 0.000001, alpha240 = 0.000001, alpha241 = 0.000001, alpha242 = 0.000001, alpha243 = 0.000001, alpha244 = 0.000001, alpha245 = 0.000001, alpha246 = 0.000001, alpha247 = 0.000001, alpha248 = 0.000001, alpha249 = 0.000001, alpha250 = 0.000001, alpha251 = 0.000001, alpha252 = 0.000001, alpha253 =
```

### 3.2. Results of GARCH with constant higher moments

```
# Estimate the model
garchfit.EWMA[[i]] <- ugarchfit(data = R, spec = garchspec.EWMA[[i]])
# Compute stdret using residuals()
stdret.EWMA[[i]] <- residuals(garchfit.EWMA[[i]], standardize = TRUE)
}

# make the histogram
#
# chart.Histogram(stdret.iGARCH[[1]], methods = c("add.normal", "add.density" ),
#               colorset = c("gray", "red", "blue"))

table3 <- matrix(nrow = 12, ncol = 5)
colnames(table3) <- distributions

#trying a loop, maybe you can solve that @filippo?
## column loop i = normal distribution, std, sstd, ged, sged
table3[1,1] <- garchfit.sGARCH[[1]]@fit$coef[1] #first parameter estimate
table3[2,1] <- garchfit.sGARCH[[1]]@fit$se.coef[1] #first standard error
table3[3,1] <- garchfit.sGARCH[[1]]@fit$coef[2] #second parameter estimate
table3[4,1] <- garchfit.sGARCH[[1]]@fit$se.coef[2]

#...
table3 <- round(table3, 3)

# for (i in length(distributions)) {
#   for (j in nrow(table3)) {
#     table3[j,i] <- garchfit.sGARCH[[i]]@fit$coef
```

### 3. Empirical Findings

```
#      table3[j+1,i] <-garchfit.sGARCH[[i]]@fit$se.coef  
#    }  
# }
```

```
print("sGARCH")  
garchfit.sGARCH[[1]]@fit$coef  
garchfit.sGARCH[[1]]@fit$se.coef
```

```
print("iGARCH")  
garchfit.iGARCH[[1]]@fit$coef  
garchfit.iGARCH[[1]]@fit$se.coef
```

```
print("EWMA")  
garchfit.EWMA[[1]]@fit$coef  
c(garchfit.EWMA[[1]]@fit$se.coef[1:2],NA,garchfit.EWMA[[1]]@fit$se.coef[3], NA)
```

```
print("eGARCH")  
garchfit.eGARCH[[1]]@fit$coef  
garchfit.eGARCH[[1]]@fit$se.coef
```

```
print("gjrGARCH")  
garchfit.gjrGARCH[[1]]@fit$coef  
garchfit.gjrGARCH[[1]]@fit$se.coef
```

```
print("NAGARCH")  
garchfit.fGARCH.NAGARCH[[1]]@fit$coef  
garchfit.fGARCH.NAGARCH[[1]]@fit$se.coef
```

```
print("TGARCH")  
garchfit.fGARCH.TGARCH[[1]]@fit$coef  
garchfit.fGARCH.TGARCH[[1]]@fit$se.coef
```

### 3.3. Results of GARCH with time-varying higher moments

```
print("TSGARCH/AVGARCH")
garchfit.fGARCH.AVGARCH[[1]]@fit$coef
garchfit.fGARCH.AVGARCH[[1]]@fit$se.coef
```

## 3.3 Results of GARCH with time-varying higher moments

```
require(racd)
require(rugarch)
require(parallel)
require(xts)
# ACD specification
sGARCH_ACDspec = acdspec(mean.model = list(armaOrder = c(1, 0)), variance.model = list(
distribution.model = list(model = 'jsu', skewOrder = c(1, 1, 1), shapeOrder = c(1,1,1)

# sGARCH
cl = makePSOCKcluster(10)
fit = acdfit(sGARCH_ACDspec, as.data.frame(R), solver = 'msoptim', solver.control = 1

# plotxts comes from implementing https://stackoverflow.com/a/50051183/271616
# par(mfrow = c(2, 2), mai = c(0.75, 0.75, 0.3, 0.3))
# cm <- plot.zoo(xts(fit@model$modeldata$data, fit@model$modeldata$index), auto.gri
# cm <- lines(fitted(fit), col = 2)
# cm
# cs <- plot(xts(abs(fit@model$modeldata$data), fit@model$modeldata$index), auto.gri
# minor.ticks = FALSE, main = 'Conditional Sigma', yaxis.right = F, col = 'grey')
# cs <- lines(sigma(fit), col = 'steelblue')
# cs
# plot(racd::skewness(fit), col = 'steelblue', yaxis.right = F, main = 'Conditional
# plot(racd::kurtosis(fit), col = 'steelblue', yaxis.right = F, main = 'Conditional
```

### 3. Empirical Findings

```
# pnl <- function(fitted(fit),xts(fit@model$modeldata$data, fit@model$modeldata$index),
#   panel.number <- parent.frame()$panel.number
#   if (panel.number == 1) lines(fitted(fit), xts(fit@model$modeldata$data, fit@model$modeldata$index),
#   lines(fitted(fit),xts(fit@model$modeldata$data, fit@model$modeldata$index),
# }
# plot(xts(fit@model$modeldata$data, fit@model$modeldata$index), auto.grid = T,
# # lines(fitted(fit), col = 2) + grid()
#
# plot(xts(fit@model$modeldata$data, fit@model$modeldata$index), auto.grid = T,
```

# 4

## Robustness Analysis

### 4.1 Specification checks

In order to check if the models are specified correctly, some specification checks have to be performed. The specification checks have to be done on the standardized residuals of the estimated GARCH model given by the following equation:

$$\hat{Z}_t = \frac{\hat{\varepsilon}_t}{\hat{\sigma}_t} = \frac{R_t - \hat{\mu}}{\hat{\sigma}_t}$$

#### 4.1.1 Figures control tests

Autocorrelation function of the standardized residuals and autocorrelation function of the squared standardized residuals.

Then the density can be examined standardized residuals and compared with the normal distribution.

Also the QQ-plot can be examined.

#### 4.1.2 Residual heteroscedasticity

Ljung-Box test on the squared or absolute standardized residuals.

zero-mean unit-variance not skewed no excess kurtosis no serial correlation in the squares no serial correlation in the cubes no serial correlation in the squares

560

561

# 5

## Conclusion



# Appendices



# A

## Appendix

### A.1 Alternative distributions than the normal

#### A.1.1 Student's t-distribution

A common alternative for the normal distribution is the Student t distribution. Similarly to the normal distribution, it is also symmetric (skewness is equal to zero). The probability density function (pdf), consistent with Ghalanos (2020a), is given by equation (A.1). As will be seen in 1.3, GARCH models are used for volatility modeling in practice. Bollerslev (1987) examined the use of the GARCH-Student or GARCH-t model as an alternative to the standard Normal distribution, which relaxes the assumption of conditional normality by assuming the standardized innovation to follow a standardized Student t-distribution (Bollerslev 2008).

$$f(x) = \frac{\Gamma(\frac{\nu+1}{2})}{\Gamma(\frac{\nu}{2})\sqrt{\beta\pi\nu}} \left(1 + \frac{(x-\alpha)^2}{\beta\beta}\right)^{-(\nu+1)/2} \quad (\text{A.1})$$

where  $\alpha, \beta$  and  $\nu$  are respectively the location, scale and shape (tail-thickness) parameters. The symbol  $\Gamma$  is the Gamma function.

Unlike the normal distribution, which depends entirely on two moments only, the student t distribution has fatter tails (thus it has a kurtosis coefficient), if the

## A. Appendix

degrees of freedom are finite. This kurtosis coefficient is given by equation (A.2). This is useful while as already mentioned, the standardized residuals appear to have fatter tails than the normal distribution following Bollerslev (2008).

$$kurt = 3 + \frac{6}{\nu - 4} \quad (\text{A.2})$$

### A.1.2 Generalized Error Distribution

The GED distribution is nested in the generalized t distribution by McDonald and Newey (1988) is used in the GED-GARCH model by Nelson (1991) to model stock market returns. This model replaced the assumption of conditional normally distributed error terms by standardized innovations that following a generalized error distribution. It is a symmetric, unimodal distribution (location parameter is the mode, median and mean). This is also sometimes called the exponential power distribution (Bollerslev 2008). The conditional density (pdf) is given by equation (A.3) following Ghalanos (2020a).

$$f(x) = \frac{pe^{\left|\frac{x - \mu}{\sigma}\right|^p}}{2^{1+p(-1)}\sigma\Gamma(p^{-1})} \quad (\text{A.3})$$

where  $\mu, \sigma$  and  $p$  are respectively the location, scale and shape parameters .

### A.1.3 Skewed t-distribution

The density function can be derived following Fernández and Steel (1998) who showed how to introduce skewness into uni-modal standardized distributions (Trottier and Ardia 2015). The first equation from Trottier and Ardia (2015), here equation (A.4) presents the skewed t-distribution.

$$f_{\xi}(z) \equiv \frac{2\sigma_{\xi}}{\xi + \xi^{-1}} f_1(z_{\xi}), \quad z_{\xi} \equiv \begin{cases} \xi^{-1}(\sigma_{\xi}z + \mu_{\xi}) & \text{if } z \geq -\mu_{\xi}/\sigma_{\xi} \\ \xi(\sigma_{\xi}z + \mu_{\xi}) & \text{if } z < -\mu_{\xi}/\sigma_{\xi} \end{cases} \quad (\text{A.4})$$

where  $\mu_\xi \equiv M_1 (\xi - \xi^{-1})$ ,  $\sigma_\xi^2 \equiv (1 - M_1^2) (\xi^2 + \xi^{-2}) + 2M_1^2 - 1$ ,  $M_1 \equiv 2 \int_0^\infty u f_1(u) du$  and  $\xi$  between 0 and  $\infty$ .  $f_1(\cdot)$  is in this case equation (A.1), the pdf of the student t distribution.

According to Giot and Laurent (2003); Giot and Laurent (2004), the skewed t-distribution outperforms the symmetric density distributions.

#### A.1.4 Skewed Generalized Error Distribution

What also will be interesting to examine is the SGED distribution of Theodossiou (2000) in GARCH models, as in the work of Lee et al. (2008). The SGED distribution extends the Generalized Error Distribution (GED) to allow for skewness and leptokurtosis. The density function can be derived following Fernández and Steel (1998) who showed how to introduce skewness into uni-modal standardized distributions (Trottier and Ardia 2015). It can also be found in Theodossiou (2000). The pdf is then given by the same equation (A.4) as the skewed t-distribution but with  $f_1(\cdot)$  equal to equation (A.3).

## A.2 GARCH models

All the GARCH models are estimated using the package “rugarch” by Ghalanos (2020b). We use specifications similar to Ghalanos (2020a). Parameters have to be restricted so that the variance output always is positive, except for the EGARCH model, as this model does not mathematically allow for a negative output.

### A.2.1 GARCH model

The standard GARCH model (Bollerslev 1986) is written consistent with Ghalanos (2020a) as in equation (A.5) without external regressors.

$$\sigma_t^2 = \omega + \sum_{j=1}^q \alpha_j \varepsilon_{t-j}^2 + \sum_{j=1}^p \beta_j \sigma_{t-j}^2 \quad (\text{A.5})$$

## A. Appendix

619 where  $\sigma_t^2$  denotes the conditional variance,  $\omega$  the intercept and  $\varepsilon_t^2$  the residuals from  
620 the used mean process. The GARCH order is defined by  $(q, p)$  (ARCH, GARCH).  
621 As Ghalanos (2020a) describes: “one of the key features of the observed behavior of  
622 financial data which GARCH models capture is volatility clustering which may be  
623 quantified in the persistence parameter  $\hat{P}$ ” specified as in equation (A.6).

$$\hat{P} = \sum_{j=1}^q \alpha_j + \sum_{j=1}^p \beta_j. \quad (\text{A.6})$$

624 The unconditional variance of the standard GARCH model of Bollerslev is very  
625 similar to the ARCH model, but with the Garch parameters ( $\beta$ 's) included as  
626 in equation (A.7).

$$\begin{aligned} \hat{\sigma}^2 &= \frac{\hat{\omega}}{1 - \hat{P}} \\ &= \frac{\hat{\omega}}{1 - \alpha - \beta} \end{aligned} \quad (\text{A.7})$$

### 627 A.2.2 IGARCH model

628 Following Ghalanos (2020a), the integrated GARCH model (Bollerslev 1986) can  
629 also be estimated. This model assumes the persistence  $\hat{P} = 1$ . This is done by  
630 Ghalanos, by setting the sum of the ARCH and GARCH parameters to 1. Because  
631 of this unit-persistence, the unconditional variance cannot be calculated.

### A.2.3 EGARCH model

The EGARCH model or exponential GARCH model (Nelson 1991) is defined as in equation (A.8). The advantage of the EGARCH model is that there are no parameter restrictions, since the output is log variance (which cannot be negative mathematically), instead of variance.

$$\log_e(\sigma_t^2) = \omega + \sum_{j=1}^q (\alpha_j z_{t-j} + \gamma_j (|z_{t-j}| - E|z_{t-j}|)) + \sum_{j=1}^p \beta_j \log_e(\sigma_{t-j}^2) \quad (\text{A.8})$$

where  $\alpha_j$  captures the sign effect and  $\gamma_j$  the size effect.

### A.2.4 GJRARCH model

The GJRARCH model (Glosten et al. 1993) models both positive as negative shocks on the conditional variance asymmetrically by using an indicator variable  $I$ , it is specified as in equation (A.9).

$$\sigma_t^2 = \omega + \sum_{j=1}^q (\alpha_j \varepsilon_{t-j}^2 + \gamma_j I_{t-j} \varepsilon_{t-j}^2) + \sum_{j=1}^p \beta_j \sigma_{t-j}^2 \quad (\text{A.9})$$

where  $\gamma_j$  represents the *leverage* term. The indicator function  $I$  takes on value 1 for  $\varepsilon \leq 0$ , 0 otherwise. Because of the indicator function, persistence of the model now crucially depends on the asymmetry of the conditional distribution used according to Ghalanos (2020a).

### A.2.5 NAGARCH model

The NAGARCH or nonlinear asymmetric model (Engle and Ng 1993). It is specified as in equation (A.10). The model is *asymmetric* as it allows for positive and negative shocks to differently affect conditional variance and *nonlinear* because a large shock is not a linear transformation of a small shock.

## A. Appendix

$$\sigma_t^2 = \omega + \sum_{j=1}^q \alpha_j (\varepsilon_{t-j} + \gamma_j \sqrt{\sigma_{t-j}})^2 + \sum_{j=1}^p \beta_j \sigma_{t-j}^2 \quad (\text{A.10})$$

As before,  $\gamma_j$  represents the *leverage* term.

### A.2.6 TGARCH model

The TGarch or threshold model (Zakoian 1994) also models assymetries in volatility depending on the sign of the shock, but contrary to the GJRGARCH model it uses the conditional standard deviation instead of conditional variance. It is specified as in (A.11).

$$\sigma_t = \omega + \sum_{j=1}^q (\alpha_j^+ \varepsilon_{t-j}^+ \alpha_j^- + \varepsilon_{t-j}^-) + \sum_{j=1}^p \beta_j \sigma_{t-j} \quad (\text{A.11})$$

where  $\varepsilon_{t-j}^+$  is equal to  $\varepsilon_{t-j}$  if the term is negative and equal to 0 if the term is positive. The reverse applies to  $\varepsilon_{t-j}^-$ . They cite Davidian and Carroll (1987) who find that using volatility instead of variance as scaling input variable gives better variance estimates. This is due to absolute residuals (contrary to squared residuals with variance) more closely predicting variance for non-normal distributions.

### A.2.7 TSGARCH model

The absolute value Garch model or TS-Garch model, as named after Taylor (1986) and Schwert (1989), models the conditional standard deviation and is intuitively specified like a normal GARCH model, but with the absolute value of the shock term. It is specified as in (A.12).

$$\sigma_t = \omega + \sum_{j=1}^q (\alpha_j |\varepsilon_{t-j}|) + \sum_{j=1}^p \beta_j \sigma_{t-j} \quad (\text{A.12})$$



### 667 A.2.8 EWMA

668 A alternative to the series of GARCH models is the exponentially weighted moving  
 669 average or EWMA model. This model calculates conditional variance based on the  
 670 shocks from previous periods. The idea is that by including a smoothing parameter  
 671  $\lambda$  more weight is assigned to recent periods than distant periods. The  $\lambda$  must  
 672 be less than 1. It is specified as in (A.13).

$$\sigma_t^2 = (1 - \lambda) \sum_{j=1}^{\infty} (\lambda^j \varepsilon_{t-j}^2) \quad (\text{A.13})$$

673 In practice a  $\lambda$  of 0.94 is often used, such as by the financial risk management com-  
 674 pany RiskMetrics<sup>TM</sup> model of J.P. Morgan (Morgan Guaranty Trust Company 1996).

## Works Cited

- 676 Annaert, Jan (Jan. 2021). *Quantitative Methods in Finance*. Version 0.2.1. Antwerp  
677 Management School.
- 678 Bali, Turan G., Hengyong Mo, and Yi Tang (Feb. 2008). “The role of autoregressive  
679 conditional skewness and kurtosis in the estimation of conditional VaR”. In: *Journal*  
680 *of Banking and Finance* 32.2. Publisher: North-Holland, pp. 269–282. DOI:  
681 10.1016/j.jbankfin.2007.03.009.
- 682 Bali, Turan G. and Panayiotis Theodossiou (Feb. 22, 2007). “A conditional-SGT-VaR  
683 approach with alternative GARCH models”. In: *Annals of Operations Research* 151.1,  
684 pp. 241–267. DOI: 10.1007/s10479-006-0118-4. URL:  
685 <http://link.springer.com/10.1007/s10479-006-0118-4>.
- 686 Basel Committee on Banking Supervision (2016). *Minimum capital requirements for*  
687 *market risk*. Tech. rep. Issue: January Publication Title: Bank for International  
688 Settlements, pp. 92–92. URL: [https://www.bis.org/basel\\_framework/](https://www.bis.org/basel_framework/).
- 689 Bertsimas, Dimitris, Geoorey J Lauprete, and Alexander Samarov (2004). “Shortfall as a  
690 risk measure: properties, optimization and applications”. In: *Journal of Economic*  
691 *Dynamics and Control* 28, pp. 1353–1381. DOI: 10.1016/S0165-1889(03)00109-X.
- 692 Bollerslev, Tim (1986). “Generalized Autoregressive Conditional Heteroskedasticity”. In:  
693 *Journal of Econometrics* 31, pp. 307–327.
- 694 — (1987). “A Conditionally Heteroskedastic Time Series Model for Speculative Prices  
695 and Rates of Return”. In: *The Review of Economics and Statistics* 69.3. Publisher:  
696 The MIT Press, pp. 542–547. DOI: 10.2307/1925546. URL:  
697 <https://www.jstor.org/stable/1925546>.
- 698 — (Sept. 4, 2008). “Glossary to ARCH (GARCH)”. In: p. 46. DOI:  
699 10.2139/ssrn.1263250. URL: <https://ssrn.com/abstract=1263250>.
- 700 Bollerslev, Tim, Robert F. Engle, and Daniel B. Nelson (Jan. 1994). “Chapter 49 Arch  
701 models”. In: *Handbook of Econometrics* 4. Publisher: Elsevier, pp. 2959–3038. DOI:  
702 10.1016/S1573-4412(05)80018-2.
- 703 Brooks, Chris et al. (2005). “Autoregressive conditional kurtosis”. In: *Journal of*  
704 *Financial Econometrics* 3.3, pp. 399–421. DOI: 10.1093/jjfinec/nbi018.
- 705 *Calculation guide STOXX®* (2020). Tech. rep.
- 706 Christoffersen, Peter, Jinyong Hahn, and Atsushi Inoue (July 2001). “Testing and  
707 comparing Value-at-Risk measures”. In: *Journal of Empirical Finance* 8.3,  
708 pp. 325–342. DOI: 10.1016/S0927-5398(01)00025-1.
- 709 Davidian, M. and R. J. Carroll (Dec. 1987). “Variance Function Estimation”. In: *Journal*  
710 *of the American Statistical Association* 82.400. Publisher: JSTOR, pp. 1079–1079.  
711 DOI: 10.2307/2289384.
- 712 Engle, R. F. (1982). “Autoregressive Conditional Heteroscedacity with Estimates of  
713 variance of United Kingdom Inflation,journal of Econometrica, Volume 50, Issue 4  
714 (Jul., 1982),987-1008.” In: *Econometrica* 50.4, pp. 987–1008.

- Engle, Robert (2001). “GARCH 101: The use of ARCH/GARCH models in applied econometrics”. In: *Journal of Economic Perspectives*. DOI: 10.1257/jep.15.4.157.
- Engle, Robert F. and S. Manganelli (1999). *CAViaR: Conditional Autoregressive Value at Risk by Regression Quantiles*. Tech. rep. San Diego: UC San Diego. URL: <http://www.jstor.org/stable/1392044>.
- Engle, Robert F. and Victor K. Ng (Dec. 1993). “Measuring and Testing the Impact of News on Volatility”. In: *The Journal of Finance* 48.5. Publisher: John Wiley and Sons, Ltd, pp. 1749–1778. DOI: 10.1111/j.1540-6261.1993.tb05127.x.
- Fama, Eugene (1970). *Efficient Capital Markets: A Review of Theory and Empirical Work*. Tech. rep. 2, pp. 383–417. DOI: 10.2307/2325486.
- Fama, Eugene F. (1965). “The Behavior of Stock-Market Prices”. In: *The Journal of Business* 38.1, pp. 34–105. URL: <http://www.jstor.org/stable/2350752>.
- Fernández, Carmen and Mark F. J. Steel (Mar. 1998). “On Bayesian Modeling of Fat Tails and Skewness”. In: *Journal of the American Statistical Association* 93.441, pp. 359–371. DOI: 10.1080/01621459.1998.10474117. URL: <http://www.tandfonline.com/doi/abs/10.1080/01621459.1998.10474117>.
- Ghalanos, Alexios (2016). *racd: Autoregressive Conditional Density Models*. <http://www.unstarched.net>, <https://bitbucket.org/alexiosg/>.
- (2020a). *Introduction to the rugarch package. (Version 1.4-3)*. URL: <http://cran.r-project.org/web/packages/rugarch/>.
- (2020b). *rugarch: Univariate GARCH models*. R package version 1.4-4.
- Giot, Pierre and Sébastien Laurent (Nov. 2003). “Value-at-risk for long and short trading positions”. In: *Journal of Applied Econometrics* 18.6, pp. 641–663. DOI: 10.1002/jae.710. URL: <http://doi.wiley.com/10.1002/jae.710>.
- (June 1, 2004). “Modelling daily Value-at-Risk using realized volatility and ARCH type models”. In: *Journal of Empirical Finance* 11.3, pp. 379–398. DOI: 10.1016/j.jempfin.2003.04.003. URL: <https://www.sciencedirect.com/science/article/pii/S092753980400012X>.
- Glosten, Lawrence R., Ravi Jagannathan, and David E. Runkle (Dec. 1993). “On the Relation between the Expected Value and the Volatility of the Nominal Excess Return on Stocks”. In: *The Journal of Finance* 48.5. Publisher: John Wiley and Sons, Ltd, pp. 1779–1801. DOI: 10.1111/j.1540-6261.1993.tb05128.x. URL: <http://doi.wiley.com/10.1111/j.1540-6261.1993.tb05128.x>.
- Hansen, Bruce E. (1994). “Autoregressive Conditional Density Estimation”. In: *International Economic Review* 35.3, pp. 705–730.
- Jorion, Philippe (2007). *Value at Risk: The New Benchmark For Managing Financial Risk*. 3rd ed. McGraw-Hill.
- Kupiec, P.H. (1995). “Techniques for Verifying the Accuracy of Risk Measurement Models”. In: *Journal of Derivatives* 3.2, pp. 73–84. DOI: 10.3905/jod.1995.407942.
- Lee, Ming-Chih, Jung-Bin Su, and Hung-Chun Liu (Oct. 22, 2008). “Value-at-risk in US stock indices with skewed generalized error distribution”. In: *Applied Financial Economics Letters* 4.6, pp. 425–431. DOI: 10.1080/17446540701765274. URL: <http://www.tandfonline.com/doi/abs/10.1080/17446540701765274>.
- Lyngs, Ulrik (2019). *oxforddown: An Oxford University Thesis Template for R Markdown*. <https://github.com/ulyngs/oxforddown>. DOI: 10.5281/zenodo.3484682.
- McDonald, James B. and Whitney K. Newey (Dec. 1988). “Partially Adaptive Estimation of Regression Models via the Generalized  $T$  Distribution”. In: *Econometric Theory*

## Works Cited

- 4.3, pp. 428–457. DOI: 10.1017/S0266466600013384. URL: [https://www.cambridge.org/core/product/identifier/S0266466600013384/type/journal\\_article](https://www.cambridge.org/core/product/identifier/S0266466600013384/type/journal_article).
- Morgan Guaranty Trust Company (1996). *RiskMetrics<sup>TM</sup>—Technical Document*. Tech. rep.
- Nelson, Daniel B. (Mar. 1991). “Conditional Heteroskedasticity in Asset Returns: A New Approach”. In: *Econometrica* 59.2. Publisher: JSTOR, pp. 347–347. DOI: 10.2307/2938260.
- Officer, R. R. (1972). *The Distribution of Stock Returns*. Tech. rep. 340, pp. 807–812.
- Schwert, G. William (1989). “Why Does Stock Market Volatility Change Over Time?” In: *The Journal of Finance* 44.5, pp. 1115–1153. DOI: 10.1111/j.1540-6261.1989.tb02647.x.
- Taylor, Stephen J. (1986). *Modelling financial time series*. Chichester: John Wiley and Sons, Ltd.
- Theodossiou, Panayiotis (1998). “Financial data and the skewed generalized t distribution”. In: *Management Science* 44.12 part 1. Publisher: INFORMS Inst.for Operations Res.and the Management Sciences, pp. 1650–1661. DOI: 10.1287/mnsc.44.12.1650.
- Theodossiou, Peter (2000). “Skewed Generalized Error Distribution of Financial Assets and Option Pricing”. In: *SSRN Electronic Journal*. DOI: 10.2139/ssrn.219679. URL: <http://www.ssrn.com/abstract=219679>.
- Trottier, Denis-Alexandre and David Ardia (Sept. 4, 2015). “Moments of standardized Fernandez-Steel skewed distributions: Applications to the estimation of GARCH-type models”. In: *Finance Research Letters* 18, pp. 311–316. DOI: 10.2139/ssrn.2656377. URL: <https://ssrn.com/abstract=2656377>.
- Welch, Ivo and Amit Goyal (July 2008). “A Comprehensive Look at The Empirical Performance of Equity Premium Prediction”. In: *Review of Financial Studies* 21.4, pp. 1455–1508. DOI: 10.1093/rfs/hhm014. URL: <https://academic.oup.com/rfs/article-lookup/doi/10.1093/rfs/hhm014>.
- Zakoian, Jean Michel (1994). “Threshold heteroskedastic models”. In: *Journal of Economic Dynamics and Control* 18.5, pp. 931–955. DOI: 10.1016/0165-1889(94)90039-6.