



The Spectrometer/Telescope for Imaging X-rays on-board Solar Orbiter: from photon to electron visibilities.

Università di Genova
DIMA | Dipartimento di Matematica

Anna Volpara, Paolo Massa, Andrea Francesco Battaglia,
Säm Krucker, Gordon Hurford, Gordon Emslie, Anna Maria Massone, Michele Piana

SPHERE Workshop

June 22, 2023



Università
di Genova



Outline

1. From photon to electron visibilities & Visibility inversion algorithm
2. Photon maps \rightarrow Electron maps \rightarrow Regularized photon maps
3. Physics inferred from electron maps
4. Conclusions and future works

Outline

- 1. From photon to electron visibilities & Visibility inversion algorithm**
2. Photon maps → Electron maps → Regularized photon maps
3. Physics inferred from electron maps
4. Conclusions and future works

From photon to electron visibilities

Photon visibilities:

$$V(u, v; \epsilon) = \mathcal{F}(I(x, y; \epsilon)) = \int \int I(x, y; \epsilon) e^{2\pi i(xu + yv)} dx dy \tag{1}$$

Electron visibilities:

$$W(u, v; E) = \frac{a}{4\pi R^2} \int \int N(x, y) \bar{F}(x, y; E) e^{2\pi i(xu + yv)} dx dy \tag{2}$$

Bremsstrahlung equation for visibilities:

$$V(u, v; \epsilon) = \int_{\epsilon}^{\infty} W(u, v; E) Q(\epsilon, E) dE \tag{3}$$

Prato et al., *A Regularized Visibility-Based Approach to Astronomical Imaging Spectroscopy*, SIAM Journal on Imaging Sciences, (2009)
Piana et al., *Electron flux spectral imaging of solar flares through regularized analysis of hard x-ray source visibilities*, The Astrophysical Journal, (2007)

From photon to electron visibilities

Photon visibilities:

$$V(u, v; \epsilon) = \mathcal{F}(I(x, y; \epsilon)) = \int \int I(x, y; \epsilon) e^{2\pi i(xu + yv)} dx dy \tag{1}$$

Electron visibilities:

$$W(u, v; E) = \frac{a}{4\pi R^2} \int \int \boxed{N(x, y) \bar{F}(x, y; E)} e^{2\pi i(xu + yv)} dx dy \tag{2}$$


Bremsstrahlung equation for visibilities:

$$V(u, v; \epsilon) = \int_{\epsilon}^{\infty} \boxed{W(u, v; E)} Q(\epsilon, E) dE \tag{3}$$

Prato et al., *A Regularized Visibility-Based Approach to Astronomical Imaging Spectroscopy*, SIAM Journal on Imaging Sciences, (2009)
Piana et al., *Electron flux spectral imaging of solar flares through regularized analysis of hard x-ray source visibilities*, The Astrophysical Journal, (2007)

Visibility inversion algorithm - Photon visibilities

Photon visibilities



Piana et al., *Electron flux spectral imaging of solar flares through regularized analysis of hard x-ray source visibilities*, The Astrophysical Journal, (2007)

Visibility inversion algorithm - Photon visibilities

Photon visibilities

	ε_1	ε_2	ε_N
(u_1, v_1)	$V_{1,1}$	$V_{1,2}$	$V_{1,N}$
(u_2, v_2)	$V_{2,1}$	$V_{2,2}$	$V_{2,N}$
\vdots	\vdots	\vdots	\ddots	\vdots
(u_{N_V}, v_{N_V})	$V_{N_V,1}$	$V_{N_V,2}$	$V_{N_V,N}$

Piana et al., *Electron flux spectral imaging of solar flares through regularized analysis of hard x-ray source visibilities*, The Astrophysical Journal, (2007)


Visibility inversion algorithm - Photon visibilities

Photon visibilities

	ε_1	ε_2	ε_N
(u_1, v_1)	$V_{1,1}$	$V_{1,2}$	$V_{1,N}$
(u_2, v_2)	$V_{2,1}$	$V_{2,2}$	$V_{2,N}$
\vdots	\vdots	\vdots	\ddots	\vdots
(u_{N_V}, v_{N_V})	$V_{N_V,1}$	$V_{N_V,2}$	$V_{N_V,N}$

Piana et al., *Electron flux spectral imaging of solar flares through regularized analysis of hard x-ray source visibilities*, The Astrophysical Journal, (2007)

Visibility inversion algorithm - Photon visibilities



Piana et al., *Electron flux spectral imaging of solar flares through regularized analysis of hard x-ray source visibilities*, The Astrophysical Journal, (2007)


Visibility inversion algorithm

Photon visibilities

	ε_1	ε_2	ε_N
(u_1, v_1)	$V_{1,1}$	$V_{1,2}$	$V_{1,N}$
(u_2, v_2)	$V_{2,1}$	$V_{2,2}$	$V_{2,N}$
...
(u_{N_V}, v_{N_V})	$V_{N_V,1}$	$V_{N_V,2}$	$V_{N_V,N}$

Prato et al., *A Regularized Visibility-Based Approach to Astronomical Imaging Spectroscopy*, SIAM Journal on Imaging Sciences, (2009)
Piana et al., *Electron flux spectral imaging of solar flares through regularized analysis of hard x-ray source visibilities*, The Astrophysical Journal, (2007)


Visibility inversion algorithm



Prato et al., *A Regularized Visibility-Based Approach to Astronomical Imaging Spectroscopy*, SIAM Journal on Imaging Sciences, (2009)


Piana et al., *Electron flux spectral imaging of solar flares through regularized analysis of hard x-ray source visibilities*, The Astrophysical Journal, (2007)

Visibility inversion algorithm



Prato et al., *A Regularized Visibility-Based Approach to Astronomical Imaging Spectroscopy*, SIAM Journal on Imaging Sciences, (2009)
Piana et al., *Electron flux spectral imaging of solar flares through regularized analysis of hard x-ray source visibilities*, The Astrophysical Journal, (2007)

Visibility inversion algorithm



Prato et al., *A Regularized Visibility-Based Approach to Astronomical Imaging Spectroscopy*, SIAM Journal on Imaging Sciences, (2009)
Piana et al., *Electron flux spectral imaging of solar flares through regularized analysis of hard x-ray source visibilities*, The Astrophysical Journal, (2007)

Visibility inversion algorithm




Figure: Reconstruction provided by MEM_GE, from electron visibilities.

Prato et al., *A Regularized Visibility-Based Approach to Astronomical Imaging Spectroscopy*, SIAM Journal on Imaging Sciences, (2009)


Piana et al., *Electron flux spectral imaging of solar flares through regularized analysis of hard x-ray source visibilities*, The Astrophysical Journal, (2007)

Outline

1. From photon to electron visibilities & Visibility inversion algorithm
- 2. Photon maps → Electron maps → Regularized photon maps**
3. Physics inferred from electron maps
4. Conclusions and future works

May 7, 2021

Photon images



Electron flux images






Figure: Photon images (*left panels*) for the energy intervals shown, compared with the electron flux images corresponding to the regularized electron visibilities (*right panels*) in the same energy range. The maps are produced using the MEM-GE algorithm.

May 7, 2021

Photon images



Regularized photon images



Electron flux images






Figure: Photon images (*left top panels*) for the energy intervals shown, compared with the electron flux images (*right panels*) and regularized photon maps (*left bottom panels*) in the same energy range. The maps are produced using the MEM-GE algorithm.

May 8, 2021

Photon images



Regularized photon images



Electron flux images




Figure: Photon images (*left top panels*) for the energy intervals shown, compared with the electron flux images (*right panels*) and regularized photon maps (*left bottom panels*) in the same energy range. The maps are produced using the MEM-GE algorithm.

Outline

1. From photon to electron visibilities & Visibility inversion algorithm
2. Photon maps \rightarrow Electron maps \rightarrow Regularized photon maps
- 3. Physics inferred from electron maps**
4. Conclusions and future works

May 8, 2021




Figure: Pixel-wise spectrum obtained from photon maps and regularized photon maps. Top left panel: selected pixels are indicated with colored crosses. The other panels show the pixel-wise spectrum obtained from photon maps (*solid line*) and regularized photon maps (*dotted line*). The pixels selected in the top left panel and their respective spectra are indicated with the same colour. Plots are logarithmic scaled on the y-axis.

May 8, 2021




Figure: Pixel-wise spectrum obtained from regularized electron maps. Left panel: selected pixels are indicated with colored crosses. Right panel shows the pixel-wise spectrum obtained from regularized electron maps. The pixels selected in the top left panel and their respective spectra are indicated with the same colour. Plots are logarithmic scaled on the y-axis.

May 8, 2021




Figure: Pixel-wise spectrum obtained from regularized electron maps. Left panel: selected pixels are indicated with colored crosses. Right panel shows the pixel-wise spectrum obtained from regularized electron maps. The pixels selected in the top left panel and their respective spectra are indicated with the same colour. Plots are logarithmic scaled on the y-axis.

May 8, 2021




Figure: Pixel-wise spectrum obtained from regularized electron maps. Left panel: selected pixels are indicated with colored crosses. Right panel shows the pixel-wise spectrum obtained from regularized electron maps. The pixels selected in the top left panel and their respective spectra are indicated with the same colour. Plots are logarithmic scaled on the y-axis.

May 8, 2021

The solution obtaining by inverting Equation (2)

$$W(u, v; E) = \frac{a}{4\pi R^2} \int \int N(x, y) \bar{F}(x, y; E) e^{2\pi i(xu + yv)} dx dy \quad (2)$$

provides the quantity:

$$REM_s(x, y; E) := N(x, y) \bar{F}(x, y; E)$$

May 8, 2021

From electron flux maps we have: $REM_s(x, y; E) \sim \bar{n}(x, y) \ell(x, y) \bar{F}(x, y; E)$ (4)

May 8, 2021

From electron flux maps we have: $REM_s(x, y; E) \sim \bar{n}(x, y) \ell(x, y) \bar{F}(x, y; E)$ (4)

From:

1. the solution of (5)

$$\frac{dE}{ds} = - \frac{K n}{E} \quad K = 2\pi e^4 \Lambda \quad (5)$$

2. the electron continuity equation in (6):

$$F(E) dE = F(E_o) dE_o \quad (6)$$

May 8, 2021

From electron flux maps we have: $REM_s(x, y; E) \sim \bar{n}(x, y) \ell(x, y) \bar{F}(x, y; E)$ (4)

From:

1. the solution of (5)

$$\frac{dE}{ds} = - \frac{K n}{E} \qquad K = 2\pi e^4 \Lambda \qquad (5)$$

2. the electron continuity equation in (6):


$$F(E) dE = F(E_o) dE_o \qquad (6)$$

$$REM_s(x, y; E) = A(x, y) E (E^2 + E_*^2(x, y))^{-\frac{\delta+1}{2}} \qquad (7)$$

where $E_*(x, y) = \sqrt{2K\tilde{N}(x, y)}$

Figure: The position of the maximum of $F(E) = E(E^2 + E_*^2)^{\frac{\delta+1}{2}}$ (in the plot $\delta = 5$) is linked to the value of E_* by:

$$E_{MAX} = \frac{E_*}{\sqrt{\delta}}$$



May 8, 2021





Figure: Top panel shows the selected pixels.

Table: E_* values for the selected pixels in the top panel. The further you move away from pixel 3, the more the value of E_* increases.

$$REM_s(x, y; E) = A(x, y) E (E^2 + E_*^2(x, y))^{-\frac{\delta+1}{2}} \tag{7}$$



Outline

1. From photon to electron visibilities & Visibility inversion algorithm
2. Photon maps → Electron maps → Regularized photon maps
3. Physics inferred from electron maps
4. **Conclusions and future works**

Conclusions and future works

- ☑ We have described a new approach to solar hard X-ray imaging spectroscopy:
 - ☑ two-dimensional Fourier transforms of the image in the photon domain are transformed into Fourier transforms of the electron flux maps.
 - ☑ This tool also provides regularized photon visibilities corresponding to the regularized electron visibilities.

Conclusions and future works

- ☒ We have described a new approach to solar hard X-ray imaging spectroscopy:
 - ☒ two-dimensional Fourier transforms of the image in the photon domain are transformed into Fourier transforms of the electron flux maps.
 - ☒ This tool also provides regularized photon visibilities corresponding to the regularized electron visibilities.

- ☐ We are working on electron spectra for analyzing the electron transport effects.

Conclusions and future works

- ☒ We have described a new approach to solar hard X-ray imaging spectroscopy:
 - ☒ two-dimensional Fourier transforms of the image in the photon domain are transformed into Fourier transforms of the electron flux maps.
 - ☒ This tool also provides regularized photon visibilities corresponding to the regularized electron visibilities.

- ☐ We are working on electron spectra for analyzing the electron transport effects.

- ☐ We are working to obtain:
 - the average density along the line of sight;
 - number spectrum of accelerated electrons.

Conclusions and future works

- ☑ We have described a new approach to solar hard X-ray imaging spectroscopy:
 - ☑ two-dimensional Fourier transforms of the image in the photon domain are transformed into Fourier transforms of the electron flux maps.
 - ☑ This tool also provides regularized photon visibilities corresponding to the regularized electron visibilities.
- ☐ We are working on electron spectra for analyzing the electron transport effects.
- ☐ We are working to obtain:
 - the average density along the line of sight;
 - number spectrum of accelerated electrons.

$$REM_s(x, y; E) = A(x, y)E(E^2 + E_*^2(x, y))^{-\frac{\delta+1}{2}}$$

Conclusions and future works

- ☑ We have described a new approach to solar hard X-ray imaging spectroscopy:
 - ☑ two-dimensional Fourier transforms of the image in the photon domain are transformed into Fourier transforms of the electron flux maps.
 - ☑ This tool also provides regularized photon visibilities corresponding to the regularized electron visibilities.
- ☐ We are working on electron spectra for analyzing the electron transport effects.
- ☐ We are working to obtain:
 - the average density along the line of sight;
 - number spectrum of accelerated electrons.

$$REM_s(x, y; E) = A(x, y)E(E^2 + E_*^2(x, y))^{-\frac{\delta+1}{2}} \longrightarrow \tilde{N}(x, y) = \frac{E_*^2(x, y)}{2K}$$

Conclusions and future works

- ✓ We have described a new approach to solar hard X-ray imaging spectroscopy:
 - ✓ two-dimensional Fourier transforms of the image in the photon domain are transformed into Fourier transforms of the electron flux maps.
 - ✓ This tool also provides regularized photon visibilities corresponding to the regularized electron visibilities.
- We are working on electron spectra for analyzing the electron transport effects.
- We are working to obtain:
 - the average density along the line of sight;
 - number spectrum of accelerated electrons.

$$REM_s(x, y; E) = A(x, y)E(E^2 + E_*^2(x, y))^{-\frac{\delta+1}{2}} \longrightarrow \tilde{N}(x, y) = \frac{E_*^2(x, y)}{2K} \longrightarrow \bar{n}(x, y) = \frac{\tilde{N}(x, y)}{s(x, y)}$$

Conclusions and future works

- ✓ We have described a new approach to solar hard X-ray imaging spectroscopy:
 - ✓ two-dimensional Fourier transforms of the image in the photon domain are transformed into Fourier transforms of the electron flux maps.
 - ✓ This tool also provides regularized photon visibilities corresponding to the regularized electron visibilities.
- We are working on electron spectra for analyzing the electron transport effects.
- We are working to obtain:
 - the average density along the line of sight;
 - number spectrum of accelerated electrons.

$$REM_s(x, y; E) = A(x, y)E(E^2 + E_*^2(x, y))^{-\frac{\delta+1}{2}} \longrightarrow \tilde{N}(x, y) = \frac{E_*^2(x, y)}{2K} \longrightarrow \bar{n}(x, y) = \frac{\tilde{N}(x, y)}{s(x, y)}$$

$$REM_s(x, y; E) \sim \bar{n}(x, y) \ell(x, y) \bar{F}(x, y; E)$$

Conclusions and future works

- ☑ We have described a new approach to solar hard X-ray imaging spectroscopy:
 - ☑ two-dimensional Fourier transforms of the image in the photon domain are transformed into Fourier transforms of the electron flux maps.
 - ☑ This tool also provides regularized photon visibilities corresponding to the regularized electron visibilities.
- ☐ We are working on electron spectra for analyzing the electron transport effects.
- ☐ We are working to obtain:
 - the average density along the line of sight;
 - number spectrum of accelerated electrons.

$$REM_s(x, y; E) = A(x, y)E(E^2 + E_*^2(x, y))^{-\frac{\delta+1}{2}} \longrightarrow \tilde{N}(x, y) = \frac{E_*^2(x, y)}{2K} \longrightarrow \bar{n}(x, y) = \frac{\tilde{N}(x, y)}{s(x, y)}$$

$$REM_s(x, y; E) \sim \bar{n}(x, y) \ell(x, y) \bar{F}(x, y; E) \longrightarrow \bar{F}(x, y, E)$$

Conclusions and future works

- ✓ We have described a new approach to solar hard X-ray imaging spectroscopy:
 - ✓ two-dimensional Fourier transforms of the image in the photon domain are transformed into Fourier transforms of the electron flux maps.
 - ✓ This tool also provides regularized photon visibilities corresponding to the regularized electron visibilities.
- We are working on electron spectra for analyzing the electron transport effects.
- We are working to obtain:
 - the average density along the line of sight;
 - number spectrum of accelerated electrons.

$$REM_s(x, y; E) = A(x, y)E(E^2 + E_*^2(x, y))^{-\frac{\delta+1}{2}} \longrightarrow \tilde{N}(x, y) = \frac{E_*^2(x, y)}{2K} \longrightarrow \tilde{n}(x, y) = \frac{\tilde{N}(x, y)}{s(x, y)}$$

$$REM_s(x, y; E) \sim \bar{n}(x, y) \ell(x, y) \bar{F}(x, y; E) \longrightarrow \bar{F}(x, y, E) \longrightarrow n_{acc}(x, y) = \bar{F}(x, y, E) \sqrt{\frac{m}{2E}}$$

References

- Brown et al., *Fast spectral fitting of hard X-ray bremsstrahlung from truncated power-law electron spectra*, Astronomy and Astrophysics, (2008)
- Emslie et al., *An empirical method to determine electron energy modification rates from spatially resolved hard X-ray data*, The Astrophysical Journal, (2001)
- Emslie et al., *RHESSI hard X-ray imaging spectroscopy of the large gamma-ray flare of 2002 July 23*, The Astrophysical Journal, (2003)
- Krucker et al., *The Spectrometer/Telescope for Imaging X-rays (STIX)*, Astronomy and Astrophysics, (2020)
- Massa et al., *First Hard X-Ray Imaging Results by Solar Orbiter STIX*, Solar Physics, (2022)
- Massa et al., *MEM_GE: A New Maximum Entropy Method for Image Reconstruction from Solar X-Ray Visibilities*, The Astrophysical Journal, (2020)
- Massa et al., *STIX imaging I – Concept*, arxiv, (2023)
- Massone et al., *Regularized solution of the solar Bremsstrahlung inverse problem: model dependence and implementation issues*, Inverse Problems in Science and Engineering, (2004)
- Piana et al., *Electron flux spectral imaging of solar flares through regularized analysis of hard x-ray source visibilities*, The Astrophysical Journal, (2007)
- Piana et al., *Hard X-rays imaging of Solar Flares*, Springer, (2022)
- Piana et al., *Regularized electron flux spectra in the 2002 July 23 solar flare*, The Astrophysical Journal, (2003)
- Prato et al., *A Regularized Visibility-Based Approach to Astronomical Imaging Spectroscopy*, SIAM Journal on Imaging Sciences, (2009)



THANK YOU FOR THE ATTENTION!

volpara@dima.unige.it

Università di Genova
DIMA | Dipartimento di Matematica
MIDA group



**Università
di Genova**

