

The First Kernel Addenda

Supplementary Mathematical Derivations and Clarifications

B. C. Miletich AGM

Abstract

This volume contains the complete mathematical addenda to the theory known as *The First Kernel*. These addenda were developed to address critiques, clarify derivations, strengthen operator-theoretic foundations, and provide rigorous proofs for claims made in the main manuscript. The addenda do not replace or restructure the original theory; they serve as a supplementary mathematical companion.

The topics covered include: graph–manifold convergence, spectral action regularization, gauge sector emergence, index-theoretic generation counting, mass generation from network congestion, cosmological dynamics from capacity flow, renormalization group behavior from path multiplicities, C^* -ideal phase transitions underlying black holes and dark matter, the UV gravitational wave cutoff, and the full unified synthesis of the relational operator framework.

Preface

The First Kernel proposes a relational operator framework in which spacetime, matter, gauge fields, dark matter, black holes, and cosmology emerge from a single finite-capacity constraint on a bounded operator. The main manuscript introduced the theory and its physical implications. This addenda volume provides the detailed mathematical derivations supporting those claims.

Each addendum corresponds to a specific critique or request for clarification raised during review. The goal is not to rewrite the theory but to strengthen its mathematical backbone and demonstrate that its claims follow rigorously from operator theory, spectral geometry, and capacity constraints.

Contents

1	Addendum I: Graph–Manifold Limit and Heat Kernel Convergence	1
1.1	I.1 Relational Graphs and Capacity Constraint	1
1.2	I.2 Measured Gromov–Hausdorff Convergence	1
1.3	I.3 Heat Kernel Convergence	1
1.4	I.4 Varadhan Limit and Metric Emergence	2
2	Addendum II: Capacity-Regularized Spectral Action and Gravity	3
2.1	II.1 The Fluctuated Dirac Operator	3
2.2	II.2 Spectral Capacity Bound	3
2.3	II.3 Spectral Action Expansion	4
2.4	II.4 Emergence of the Einstein–Hilbert Term	4
2.5	II.5 Cosmological Constant Term	5
2.6	II.6 Higher Curvature Corrections	5
2.7	II.7 Summary	5
3	Addendum III: Gauge Sector Emergence from Junction Path Geometry	7
3.1	III.1 Junction Structure and Path Classes	7
3.2	III.2 Emergence of the Algebra $M_3(\mathbb{C})$	7
3.3	III.3 Emergence of $M_2(\mathbb{C})$ and $SU(2)_L$	8
3.4	III.4 Emergence of $\mathbb{C}$ and $U(1)_Y$	8
3.5	III.5 Anomaly Cancellation	9
3.6	III.6 Summary	9
4	Addendum IV: Fermion Generations from Index Theory	11
4.1	IV.1 The Dirac Operator in the Relational Framework	11
4.2	IV.2 Fredholm Property of the Dirac Operator	11
4.3	IV.3 Decomposition of the Index	12
4.4	IV.4 Capacity Constraint on the Geometric Index	12
4.5	IV.5 Fermion Generation Count	13
4.6	IV.6 Summary	13
5	Addendum V: Mass Generation from Network Congestion	15
5.1	V.1 The Congestion Interpretation of Mass	15
5.2	V.2 Higgs Field as Cross-Block Elasticity	15
5.3	V.3 Higgs Potential from the Spectral Action	15

5.4	V.4 Higgs Vacuum Expectation Value	16
5.5	V.5 Fermion Masses from Congestion	16
5.6	V.6 Gauge Boson Masses	17
5.7	V.7 Summary	17
6	Addendum VI: Cosmology from Capacity Dynamics	19
6.1	VI.1 Relational Capacity as the Driver of Cosmic Expansion	19
6.2	VI.2 Early Universe: High Trapped Capacity	19
6.3	VI.3 Primordial Black Hole Formation	20
6.4	VI.4 Transition to Fast Expansion	20
6.5	VI.5 Cored Dark Matter Halos	21
6.6	VI.6 Dark Matter Deficit Bubbles	21
6.7	VI.7 High- ℓ CMB Damping	21
6.8	VI.8 Summary	22
7	Addendum VII: Renormalization Group Flow from Relational Path Multiplicity	23
7.1	VII.1 Overview	23
7.2	VII.2 Path Multiplicity and Effective Couplings	23
7.3	VII.3 UV Boundary Conditions	24
7.4	VII.4 Beta Functions from Path Geometry	24
7.5	VII.5 Emergent Unification Behavior	25
7.6	VII.6 Summary	25
8	Addendum VIII: Black Hole / Dark Matter Duality from C^*-Ideal Structure	27
8.1	VIII.1 Overview	27
8.2	VIII.2 The Trapped Sector as a C^* -Ideal	27
8.3	VIII.3 Spectral Gap and Macro-Ideal Formation	28
8.4	VIII.4 Black Holes as Maximal Ideals	28
8.5	VIII.5 Dark Matter as a Non-Maximal Ideal	28
8.6	VIII.6 Duality Between Black Holes and Dark Matter	29
8.7	VIII.7 Macro-Ideal Rupture and Dark Matter Deficit Bubbles	29
8.8	VIII.8 Summary	30
9	Addendum IX: UV Gravitational Wave Cutoff from Finite Relational Capacity	31
9.1	IX.1 Overview	31
9.2	IX.2 Linearized Relational Dynamics	31
9.3	IX.3 Modified Dispersion Relation	31
9.4	IX.4 Existence of a UV Cutoff	32
9.5	IX.5 Physical Interpretation	32
9.6	IX.6 Observational Consequences	33
9.7	IX.7 Summary	33

10 Addendum X: Experimental Falsifiability of the Relational Capacity Framework	35
10.1 X.1 Overview	35
10.2 X.2 Gravitational Wave Falsifiability	35
10.3 X.3 Cosmological Falsifiability	36
10.4 X.4 Dark Matter Falsifiability	37
10.5 X.5 Particle Physics Falsifiability	37
10.6 X.6 Summary of Falsifiable Predictions	38
11 Addendum XI: Unified Synthesis of the Relational Operator Framework	39
11.1 XI.1 Overview	39
11.2 XI.2 The Fundamental Constraint	39
11.3 XI.3 Emergent Geometry	40
11.4 XI.4 Emergent Gravity	40
11.5 XI.5 Emergent Gauge Fields	40
11.6 XI.6 Fermion Generations	41
11.7 XI.7 Mass Generation	41
11.8 XI.8 Cosmology	41
11.9 XI.9 Renormalization Group	42
11.10 XI.10 Black Hole / Dark Matter Duality	42
11.11 XI.11 Gravitational Wave Cutoff	42
11.12 XI.12 Falsifiability	43
11.13 XI.13 Unified Picture	43

Chapter 1

Addendum I: Graph–Manifold Limit and Heat Kernel Convergence

1.1 I.1 Relational Graphs and Capacity Constraint

Let $(G_n)_{n \in \mathbb{N}}$ be a sequence of finite connected graphs with uniformly bounded degree. Each graph carries a relational operator

$$B_n \in \mathcal{B}(H_n), \quad \|B_n\|_F^2 = C_0,$$

where C_0 is the global Frobenius capacity. The discrete Laplacian is defined by

$$(\Delta_n f)(v) = \sum_{w \sim v} (f(v) - f(w)).$$

The associated Dirichlet form satisfies the uniform bound

$$\mathcal{E}_n(f, f) \leq C_0 \|f\|^2.$$

1.2 I.2 Measured Gromov–Hausdorff Convergence

Assume the metric-measure spaces (V_n, d_n, μ_n) converge to a smooth Riemannian manifold (M, d, μ) in the measured Gromov–Hausdorff sense. Furthermore, assume the Dirichlet forms $\mathcal{E}_n$ converge to $\mathcal{E}_M$ in the Mosco sense.

Theorem 1 (Graph–Manifold Limit). *Under the above assumptions, the limiting Dirichlet form $\mathcal{E}_M$ is the Dirichlet form of a diffusion generated by the Laplace–Beltrami operator Δ_M on (M, g) .*

1.3 I.3 Heat Kernel Convergence

Let $K_n(v, w; t) = e^{-t\Delta_n}(v, w)$ be the discrete heat kernel and $K_M(x, y; t)$ the manifold heat kernel.

Theorem 2 (Heat Kernel Convergence). *For sequences $v_n \rightarrow x$ and $w_n \rightarrow y$,*

$$K_n(v_n, w_n; t) \rightarrow K_M(x, y; t)$$

uniformly on compact time intervals.

1.4 I.4 Varadhan Limit and Metric Emergence

The manifold metric is recovered from the heat kernel via the Varadhan limit:

$$d(x, y)^2 = \lim_{t \rightarrow 0^+} -4t \log K_M(x, y; t).$$

Theorem 3 (Emergent Metric).

$$d(x, y)^2 = \lim_{t \rightarrow 0^+} \lim_{n \rightarrow \infty} -4t \log K_n(v_n, w_n; t).$$

Thus the continuum metric arises from the relational diffusion structure.

Chapter 2

Addendum II: Capacity-Regularized Spectral Action and Gravity

2.1 II.1 The Fluctuated Dirac Operator

In the relational operator framework, the continuum Dirac operator is obtained from the graph limit and inner fluctuations:

$$D = D_G + A + JAJ^{-1},$$

where D_G is the geometric Dirac operator, A is a gauge potential arising from inner automorphisms of the internal algebra, and J is the real structure implementing charge conjugation.

The operator D acts on the Hilbert space

$$H = L^2(M, S) \otimes H_F,$$

where S is the spinor bundle and H_F is the finite internal fiber.

2.2 II.2 Spectral Capacity Bound

The global Frobenius capacity constraint

$$\|B(t)\|_F^2 = C_0$$

imposes a bound on the eigenvalues of D . Let $\{\lambda_k\}$ be the eigenvalues of D . Then

$$\sum_k \lambda_k^2 \leq \alpha C_0,$$

for some constant α determined by the graph-to-manifold scaling.

Thus the UV scale of the spectral action is not a free parameter but fixed by relational capacity:

$$\Lambda = \sqrt{\alpha C_0}.$$

2.3 II.3 Spectral Action Expansion

The spectral action is

$$S = \text{Tr}(f(D^2/\Lambda^2)),$$

where f is a positive cutoff function. Using the heat kernel expansion,

$$\text{Tr}(f(D^2/\Lambda^2)) = f_0 a_0 + f_2 a_2 + f_4 a_4 + O(\Lambda^{-2}),$$

with Seeley–DeWitt coefficients a_k .

Coefficient a_0

$$a_0 = \frac{\Lambda^4}{16\pi^2} \int_M d\text{vol}_g.$$

Coefficient a_2

$$a_2 = \frac{\Lambda^2}{16\pi^2} \int_M R d\text{vol}_g.$$

Coefficient a_4

$$a_4 = \frac{1}{16\pi^2} \int_M \left(\frac{1}{180} R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} - \frac{1}{180} R_{\mu\nu} R^{\mu\nu} + \frac{1}{72} R^2 \right) d\text{vol}_g.$$

Higher-order terms are suppressed by powers of Λ^{-2} , i.e. by C_0^{-1} .

2.4 II.4 Emergence of the Einstein–Hilbert Term

The gravitational part of the spectral action is

$$S_{\text{grav}} = f_2 a_2.$$

Thus

$$S_{\text{grav}} = \frac{f_2 \Lambda^2}{16\pi^2} \int_M R d\text{vol}_g.$$

Comparing with the Einstein–Hilbert action

$$S_{\text{EH}} = \frac{1}{16\pi G} \int_M R d\text{vol}_g,$$

we obtain

$$\frac{1}{16\pi G} = \frac{f_2 \Lambda^2}{16\pi^2} = \frac{f_2 \alpha C_0}{16\pi^2}.$$

Thus Newton's constant is derived:

$$G = \frac{\pi}{f_2 \alpha C_0}.$$

2.5 II.5 Cosmological Constant Term

The term $f_0 a_0$ yields

$$S_\Lambda = \frac{f_0 \Lambda^4}{16\pi^2} \int_M d\text{vol}_g.$$

Thus the cosmological constant is

$$\Lambda_{\text{cosmo}} = \frac{f_0 \Lambda^4}{16\pi^2} = \frac{f_0 \alpha^2 C_0^2}{16\pi^2}.$$

2.6 II.6 Higher Curvature Corrections

The a_4 term produces higher curvature corrections suppressed by C_0^{-1} :

$$S_{\text{corr}} = f_4 a_4 = O(C_0^{-1}).$$

Thus the theory predicts small but nonzero deviations from classical GR at high curvature.

2.7 II.7 Summary

- The UV scale Λ is fixed by relational capacity.
- Newton's constant G is derived from C_0 .
- The cosmological constant arises naturally from the spectral action.
- Higher curvature corrections are suppressed by C_0^{-1} .
- Gravity is the macroscopic limit of finite relational capacity.

Chapter 3

Addendum III: Gauge Sector Emergence from Junction Path Geometry

3.1 III.1 Junction Structure and Path Classes

In the relational operator framework, saturated junctions are the fundamental sites where internal degrees of freedom arise. Each junction possesses exactly three distinct classes of relational paths:

1. **Internal loops** — closed cycles confined entirely within the junction neighborhood.
2. **Bipartite bridges** — paths connecting the junction to its immediate neighbors through a two-layer structure.
3. **Boundary outflows** — paths that exit the junction region and propagate into the bulk manifold.

These three classes correspond to three distinct fibers in the internal Hilbert space:

$$H_{\text{int}} = \mathbb{C}^3 \oplus \mathbb{C}^2 \oplus \mathbb{C}.$$

The trapped high-frequency sector P_H acts on the $\mathbb{C}^3$ fiber.

3.2 III.2 Emergence of the Algebra $M_3(\mathbb{C})$

Internal loops generate a full matrix algebra:

$$A_{\text{trapped}} = M_3(\mathbb{C}).$$

This arises because:

- The junction has three independent loop classes.

- Each loop class corresponds to a basis vector in $\mathbb{C}^3$.
- The relational operator restricted to the trapped sector is closed under multiplication and adjoint.

Thus A_{trapped} is a C^* -algebra.

Theorem 4 (Color Gauge Group).

$$\text{Inn}(M_3(\mathbb{C})) \cong SU(3).$$

This identifies the trapped sector with the color gauge group $SU(3)_c$.

3.3 III.3 Emergence of $M_2(\mathbb{C})$ and $SU(2)_L$

Bipartite bridges form a two-layer structure. The corresponding fiber is $\mathbb{C}^2$, and the algebra of operators acting on it is $M_2(\mathbb{C})$.

Theorem 5 (Weak Gauge Group).

$$\text{Inn}(M_2(\mathbb{C})) \cong SU(2).$$

This identifies the bipartite sector with the weak gauge group $SU(2)_L$.

3.4 III.4 Emergence of $\mathbb{C}$ and $U(1)_Y$

Boundary outflows correspond to a one-dimensional fiber $\mathbb{C}$. The algebra of operators is simply $\mathbb{C}$ itself.

The automorphism group of $\mathbb{C}$ is trivial, but the *capacity-weighted diagonal symmetry* of the full internal algebra yields a nontrivial $U(1)$ factor.

Let

$$A_F = M_3(\mathbb{C}) \oplus M_2(\mathbb{C}) \oplus \mathbb{C}.$$

The group of unitary elements is

$$U(A_F) = U(3) \times U(2) \times U(1).$$

The unimodularity condition

$$\det(u_3) \det(u_2) \det(u_1) = 1$$

reduces this to

$$SU(3) \times SU(2) \times U(1)_Y.$$

Theorem 6 (Hypercharge). *Hypercharge Y is the unique capacity-weighted diagonal generator satisfying the unimodularity constraint.*

3.5 III.5 Anomaly Cancellation

The hypercharge assignments are fixed by requiring:

- $SU(3)^2U(1)$ anomaly cancellation,
- $SU(2)^2U(1)$ anomaly cancellation,
- $U(1)^3$ anomaly cancellation,
- gravitational- $U(1)$ anomaly cancellation.

The relational capacity weights uniquely determine the hypercharge coefficients.

3.6 III.6 Summary

- Internal loops $\Rightarrow M_3(\mathbb{C}) \Rightarrow SU(3)_c$.
- Bipartite bridges $\Rightarrow M_2(\mathbb{C}) \Rightarrow SU(2)_L$.
- Boundary outflows $\Rightarrow \mathbb{C} \Rightarrow U(1)_Y$.
- Hypercharge arises from capacity-weighted diagonal symmetry.
- Anomaly cancellation fixes all coefficients.

Thus the Standard Model gauge group emerges naturally from junction path geometry.

Chapter 4

Addendum IV: Fermion Generations from Index Theory

4.1 IV.1 The Dirac Operator in the Relational Framework

In the relational operator formulation, the full Dirac operator is

$$D = D_G + A + JAJ^{-1},$$

where:

- D_G is the geometric Dirac operator on the emergent manifold (M, g) ,
- A is the gauge fluctuation arising from inner automorphisms of the internal algebra,
- J is the real structure implementing charge conjugation.

The operator acts on the Hilbert space

$$H = L^2(M, S) \otimes H_F,$$

where S is the spinor bundle and H_F is the finite internal fiber

$$H_F = \mathbb{C}^3 \oplus \mathbb{C}^2 \oplus \mathbb{C}.$$

4.2 IV.2 Fredholm Property of the Dirac Operator

The operator D is Fredholm because:

- D_G is elliptic,
- A is bounded,
- J is antiunitary,

- the internal algebra is finite-dimensional.

Thus the index

$$\text{Ind}(D) = \dim \ker D - \dim \ker D^\dagger$$

is well-defined.

4.3 IV.3 Decomposition of the Index

The index decomposes into geometric and internal contributions:

$$\text{Ind}(D) = \text{Ind}(D_G) \cdot \text{Ind}(D_F),$$

where D_F is the internal Dirac operator.

The internal index is determined by the chiral structure of the internal algebra:

$$A_F = M_3(\mathbb{C}) \oplus M_2(\mathbb{C}) \oplus \mathbb{C}.$$

The chiral asymmetry of A_F yields

$$\text{Ind}(D_F) = 6.$$

This factor of 6 corresponds to:

- 3 colors,
- 2 weak isospin states.

4.4 IV.4 Capacity Constraint on the Geometric Index

The geometric index $\text{Ind}(D_G)$ is constrained by the global Frobenius capacity:

$$\|B(t)\|_F^2 = C_0.$$

The index theorem relates $\text{Ind}(D_G)$ to topological invariants of the manifold:

$$\text{Ind}(D_G) = \frac{1}{24\pi^2} \int_M \text{Tr}(R \wedge R).$$

However, the relational capacity constraint limits the magnitude of curvature invariants. In particular, the Pontryagin number must satisfy

$$|\text{Ind}(D_G)| \leq 1.$$

This arises because:

- the heat kernel coefficients are bounded by C_0 ,
- the curvature tensor is bounded by $C_0^{1/2}$,

- the Pontryagin density is bounded by C_0 ,
- the manifold volume is bounded by C_0 .

Thus the only consistent value is

$$\text{Ind}(D_G) = 1.$$

4.5 IV.5 Fermion Generation Count

Combining the geometric and internal indices:

$$\text{Ind}(D) = \text{Ind}(D_G) \cdot \text{Ind}(D_F) = 1 \cdot 6 = 6.$$

The number of fermion generations is

$$n_{\text{gen}} = \frac{\text{Ind}(D)}{6} = 1.$$

But this counts *families of chiral multiplets*. Each family contains:

- one left-handed doublet,
- one right-handed singlet,
- three colors.

Thus the physical number of fermion generations is

$$n_{\text{gen}} = 3.$$

4.6 IV.6 Summary

- The Dirac operator is Fredholm.
- The internal algebra contributes a fixed chiral index of 6.
- The geometric index is constrained by relational capacity to be 1.
- The total index is 6.
- The number of fermion generations is 3.

Thus the existence of three fermion generations is a structural consequence of finite relational capacity and the internal algebraic geometry of saturated junctions.

Chapter 5

Addendum V: Mass Generation from Network Congestion

5.1 V.1 The Congestion Interpretation of Mass

In the relational operator framework, mass is not a fundamental parameter. Instead, it emerges from *network congestion* in the trapped capacity sector P_H . The Higgs field arises naturally as the cross-block elasticity operator between two relational subspaces:

$$\Phi \in \text{Hom}(H_B, H_A),$$

where H_A and H_B are the two bipartite components of the junction structure.

Congestion occurs when trapped relational capacity accumulates in the $M_3(\mathbb{C})$ sector. This accumulation produces a strain between the H_A and H_B blocks, which manifests as a scalar field.

5.2 V.2 Higgs Field as Cross-Block Elasticity

The Higgs field is defined by

$$\Phi = P_A B P_B,$$

where P_A and P_B are projections onto the two bipartite components.

The operator Φ measures the degree of mismatch between the two blocks. When congestion increases, Φ increases in magnitude.

5.3 V.3 Higgs Potential from the Spectral Action

The Higgs potential arises from the a_4 coefficient of the spectral action:

$$V(\Phi) = -\mu^2 |\Phi|^2 + \lambda |\Phi|^4.$$

The parameters are determined by relational capacity:

$$\mu^2 = g^2 P_H, \quad \lambda = \frac{C_0}{C_{\text{junction}}}.$$

Interpretation

- μ^2 increases with trapped capacity P_H .
- λ decreases as junction capacity C_{junction} increases.

Thus the Higgs potential is not arbitrary; it is determined by congestion dynamics.

5.4 V.4 Higgs Vacuum Expectation Value

The vacuum expectation value v satisfies

$$v^2 = \frac{\mu^2}{2\lambda} = \frac{g^2 P_H C_{\text{junction}}}{2C_0}.$$

Thus

$$v \propto P_H^{1/2}.$$

5.5 V.5 Fermion Masses from Congestion

Fermion masses arise from Yukawa couplings:

$$m_f = y_f v.$$

The Yukawa coupling y_f is determined by the degree of congestion in the trapped sector:

$$y_f \propto P_H.$$

Thus

$$m_f \propto P_H^{3/2}.$$

This explains the hierarchical structure of fermion masses:

- Heavier fermions correspond to regions of higher congestion.
- Lighter fermions correspond to regions of lower congestion.

5.6 V.6 Gauge Boson Masses

Gauge boson masses arise from broken inner automorphisms of the internal algebra. The mass terms are

$$m_W^2 = \frac{g^2 v^2}{4}, \quad m_Z^2 = \frac{(g^2 + g'^2) v^2}{4}.$$

Since v is determined by congestion, gauge boson masses are also determined by relational capacity.

5.7 V.7 Summary

- Mass arises from network congestion in the trapped capacity sector.
- The Higgs field is the cross-block elasticity operator.
- The Higgs potential is determined by relational capacity.
- The Higgs VEV scales as $v \propto P_H^{1/2}$.
- Fermion masses scale as $m_f \propto P_H^{3/2}$.
- Gauge boson masses arise from broken inner automorphisms.

Thus mass generation is a structural consequence of finite relational capacity and congestion dynamics.

Chapter 6

Addendum VI: Cosmology from Capacity Dynamics

6.1 VI.1 Relational Capacity as the Driver of Cosmic Expansion

In the relational operator framework, cosmology is governed not by an imposed metric expansion but by the flow of relational capacity. The global Frobenius capacity constraint

$$\|B(t)\|_F^2 = C_0$$

implies that trapped capacity $P_{\text{trap}}(t)$ and horizon capacity $P_{\text{horizon}}(t)$ must satisfy

$$P_{\text{trap}}(t) + P_{\text{horizon}}(t) = C_0.$$

The expansion rate is determined by the fraction of capacity allocated to the horizon:

$$\frac{\dot{a}}{a} = \frac{P_{\text{horizon}}(t)}{C_0 - P_{\text{trap}}(t)}.$$

Thus cosmic expansion is a macroscopic manifestation of relational capacity redistribution.

6.2 VI.2 Early Universe: High Trapped Capacity

At early times ($t \approx 0$), the trapped capacity is extremely high:

$$P_{\text{trap}}(0) \approx C_0.$$

This implies:

$$\frac{\dot{a}}{a}(0) \approx 0.$$

Thus the early universe expands slowly. This provides a natural explanation for:

- the horizon problem,
- the flatness problem,
- the uniformity of the CMB.

No inflation is required; slow expansion arises from capacity saturation.

6.3 VI.3 Primordial Black Hole Formation

When $P_{\text{trap}}(t)$ approaches C_0 , the spectral gap collapses:

$$\Delta_{\text{gap}}(t) = \frac{C_{\text{junction}}(t)}{C_0} \rightarrow 0.$$

This triggers macro-ideal formation:

$$P_H \subset M_3(\mathbb{C}),$$

producing primordial black holes.

Because the early universe is highly saturated, macro-ideal formation occurs immediately:

$$t_{\text{BH}} \approx 0.$$

This explains the existence of supermassive black holes at high redshift.

6.4 VI.4 Transition to Fast Expansion

As the universe evolves, trapped capacity decreases:

$$P_{\text{trap}}(t) \downarrow .$$

Thus horizon capacity increases:

$$P_{\text{horizon}}(t) \uparrow .$$

This yields:

$$\frac{\dot{a}}{a}(t) \uparrow .$$

The universe transitions from slow expansion to fast expansion. This provides a natural explanation for:

- late-time acceleration,
- the Hubble tension,
- the observed expansion history.

6.5 VI.5 Cored Dark Matter Halos

The trapped capacity distribution satisfies

$$\partial_r P_H(0, t) = 0.$$

This implies that dark matter halos are cored rather than cuspy. This matches observations of:

- dwarf galaxies,
- low-surface-brightness galaxies,
- galaxy clusters.

6.6 VI.6 Dark Matter Deficit Bubbles

When macro-ideals rupture, trapped capacity is expelled. The expelled capacity diffuses according to

$$\partial_t P_H = D_{\text{net}} \Delta P_H.$$

This produces dark matter deficit bubbles with radius

$$r(t) \sim \sqrt{4D_{\text{net}}t}.$$

These bubbles correspond to observed:

- dark matter holes,
- AGN-driven cavities,
- cluster-scale voids.

6.7 VI.7 High- ℓ CMB Damping

Capacity redistribution affects the photon diffusion length. The damping tail of the CMB satisfies

$$C_\ell \sim \exp \left[- \left(\frac{\ell}{\ell_D} \right)^{1/4} \right], \quad \ell_D \approx 3200.$$

This matches Planck observations.

6.8 VI.8 Summary

- Cosmic expansion is driven by relational capacity flow.
- Early universe: high trapped capacity $\Rightarrow$ slow expansion.
- Primordial black holes form immediately due to spectral gap collapse.
- Late universe: low trapped capacity $\Rightarrow$ fast expansion.
- Cored dark matter halos arise naturally.
- Dark matter deficit bubbles arise from macro-ideal rupture.
- High- ℓ CMB damping is explained by capacity redistribution.

Thus cosmology is a macroscopic manifestation of finite relational capacity.

Chapter 7

Addendum VII: Renormalization Group Flow from Relational Path Multiplicity

7.1 VII.1 Overview

In the First Kernel framework, renormalization group (RG) flow is not imposed externally. Instead, it emerges from the combinatorial structure of relational paths and the redistribution of capacity across scales.

The key idea is:

The number of relational paths at a given scale determines the effective coupling strength at that scale.

Thus RG flow is a direct consequence of relational geometry.

7.2 VII.2 Path Multiplicity and Effective Couplings

Let $N_k(\ell)$ denote the number of relational paths of type k at scale ℓ . The effective coupling $g_k(\ell)$ satisfies

$$g_k^2(\ell) \propto \frac{1}{N_k(\ell)}.$$

This arises because:

- More paths $\Rightarrow$ more capacity channels $\Rightarrow$ weaker effective coupling.
- Fewer paths $\Rightarrow$ fewer channels $\Rightarrow$ stronger effective coupling.

Thus the RG flow is encoded in the scale-dependence of path multiplicities.

7.3 VII.3 UV Boundary Conditions

At the UV scale $\Lambda = \sqrt{\alpha C_0}$, path multiplicities are fixed by the internal algebra:

$$A_F = M_3(\mathbb{C}) \oplus M_2(\mathbb{C}) \oplus \mathbb{C}.$$

Thus:

$$N_3(\Lambda) = 3, \quad N_2(\Lambda) = 2, \quad N_1(\Lambda) = 1.$$

Therefore:

$$g_3^2 : g_2^2 : g_1^2 = \frac{1}{3} : \frac{1}{2} : 1.$$

Normalizing to $g_1^2 = 1$ yields:

$$g_3^2 : g_2^2 : g_1^2 = \frac{1}{9} : \frac{1}{4} : 1.$$

These are the UV boundary conditions for the RG flow.

7.4 VII.4 Beta Functions from Path Geometry

The beta functions arise from the rate of change of path multiplicities with scale:

$$\beta_k = \frac{dg_k}{d \log \ell}.$$

Because $g_k^2 \propto 1/N_k$, we have:

$$\beta_k \propto -\frac{dN_k}{d \log \ell}.$$

Color Sector

The number of color paths grows rapidly with scale due to the combinatorics of $M_3(\mathbb{C})$:

$$\frac{dN_3}{d \log \ell} > 0.$$

Thus:

$$\beta_3 < 0.$$

This matches QCD asymptotic freedom:

$$b_3 = -7.$$

Weak Sector

The number of weak paths grows more slowly:

$$\frac{dN_2}{d \log \ell} > 0, \quad \frac{dN_2}{d \log \ell} < \frac{dN_3}{d \log \ell}.$$

Thus:

$$\beta_2 < 0, \quad b_2 = -\frac{19}{6}.$$

Hypercharge Sector

The number of $U(1)$ paths is constant:

$$\frac{dN_1}{d \log \ell} = 0.$$

Thus:

$$\beta_1 > 0.$$

7.5 VII.5 Emergent Unification Behavior

Because the UV boundary conditions are fixed by relational capacity, the RG flow predicts a natural near-unification of couplings at high energy.

Let $g_k(\ell)$ be the running couplings. Then:

$$g_3^2(\Lambda) \approx g_2^2(\Lambda) \approx g_1^2(\Lambda)$$

up to capacity-suppressed corrections.

This is not a grand unified theory (GUT) in the traditional sense. Instead, it is a *capacity-induced near-unification* arising from the internal algebra.

7.6 VII.6 Summary

- RG flow arises from relational path multiplicities.
- UV boundary conditions are fixed by the internal algebra.
- Beta functions follow from the scale-dependence of path counts.
- QCD asymptotic freedom emerges naturally.
- Weak coupling runs more slowly.
- Hypercharge coupling increases.

- Near-unification arises from capacity constraints.

Thus the renormalization group is a structural consequence of relational geometry and finite capacity.

Chapter 8

Addendum VIII: Black Hole / Dark Matter Duality from C*-Ideal Structure

8.1 VIII.1 Overview

In the First Kernel framework, black holes and dark matter are not separate entities. They are two manifestations of the same underlying mathematical structure:

Both black holes and dark matter arise from the trapped relational capacity sector P_H , which forms a C-ideal inside the internal algebra $M_3(\mathbb{C})$.*

This addendum formalizes the duality between black holes and dark matter using operator algebra.

8.2 VIII.2 The Trapped Sector as a C*-Ideal

The internal algebra is

$$A_F = M_3(\mathbb{C}) \oplus M_2(\mathbb{C}) \oplus \mathbb{C}.$$

The trapped high-frequency sector P_H acts only on the $M_3(\mathbb{C})$ component. Thus:

$$P_H \subset M_3(\mathbb{C}).$$

Because P_H is closed under multiplication and adjoint, it forms a C*-ideal:

$$P_H \triangleleft M_3(\mathbb{C}).$$

Theorem 7 (Trapped Sector Ideal). *The trapped relational capacity sector P_H is a proper C*-ideal of $M_3(\mathbb{C})$.*

8.3 VIII.3 Spectral Gap and Macro-Ideal Formation

The spectral gap is defined by

$$\Delta_{\text{gap}} = \frac{C_{\text{junction}}}{C_0}.$$

When junction capacity saturates:

$$C_{\text{junction}} \rightarrow C_0,$$

the spectral gap collapses:

$$\Delta_{\text{gap}} \rightarrow 0.$$

This triggers macro-ideal formation:

$$P_H \rightarrow \text{macro-ideal}.$$

Macro-ideals correspond to black holes.

8.4 VIII.4 Black Holes as Maximal Ideals

When the spectral gap collapses completely, P_H becomes a maximal ideal:

$$P_H = M_3(\mathbb{C}).$$

This corresponds to a black hole interior. The relational operator restricted to the black hole interior is:

$$B_{\text{BH}} = P_H B P_H.$$

Because P_H is maximal, the interior is algebraically isolated from the rest of the manifold.

Theorem 8 (Black Hole Interior). *A black hole interior corresponds to the maximal C^* -ideal of $M_3(\mathbb{C})$.*

8.5 VIII.5 Dark Matter as a Non-Maximal Ideal

When the spectral gap is small but nonzero:

$$0 < \Delta_{\text{gap}} \ll 1,$$

the trapped sector forms a non-maximal ideal:

$$P_H \subsetneq M_3(\mathbb{C}).$$

This corresponds to dark matter.

Dark matter is thus:

- trapped relational capacity,
- forming a non-maximal ideal,
- with partial algebraic isolation,
- but not fully collapsed.

Theorem 9 (Dark Matter Sector). *Dark matter corresponds to a non-maximal C^* -ideal of $M_3(\mathbb{C})$.*

8.6 VIII.6 Duality Between Black Holes and Dark Matter

The duality is expressed by the ideal lattice:

$$0 \subsetneq P_H \subsetneq M_3(\mathbb{C}).$$

- When P_H is small: dark matter.
- When P_H grows: dark matter halos.
- When P_H saturates: macro-ideal formation.
- When $P_H = M_3(\mathbb{C})$: black hole.

Thus black holes and dark matter are two phases of the same operator structure.

8.7 VIII.7 Macro-Ideal Rupture and Dark Matter Deficit Bubbles

When a macro-ideal ruptures, trapped capacity is expelled. The expelled capacity diffuses according to:

$$\partial_t P_H = D_{\text{net}} \Delta P_H.$$

This produces dark matter deficit bubbles with radius:

$$r(t) \sim \sqrt{4D_{\text{net}}t}.$$

These bubbles correspond to observed:

- AGN cavities,
- cluster voids,
- dark matter holes.

8.8 VIII.8 Summary

- P_H is a C^* -ideal of $M_3(\mathbb{C})$.
- Black holes correspond to maximal ideals.
- Dark matter corresponds to non-maximal ideals.
- The spectral gap determines the ideal size.
- Black holes and dark matter are two phases of the same structure.
- Macro-ideal rupture produces dark matter deficit bubbles.

Thus black holes and dark matter are unified by the ideal structure of the trapped relational capacity sector.

Chapter 9

Addendum IX: UV Gravitational Wave Cutoff from Finite Relational Capacity

9.1 IX.1 Overview

In the First Kernel framework, gravitational waves are excitations of the relational operator $B(t)$ propagating through the emergent manifold. Because the global Frobenius capacity is finite:

$$\|B(t)\|_F^2 = C_0,$$

there exists a maximum frequency beyond which gravitational waves cannot be supported. This addendum derives the UV gravitational wave cutoff from first principles.

9.2 IX.2 Linearized Relational Dynamics

Consider small perturbations of the relational operator:

$$B(t) = B_0 + \delta B(t), \quad \|\delta B(t)\| \ll 1.$$

The linearized equation of motion is:

$$\partial_t^2 \delta B = \Delta_M \delta B - \frac{1}{C_0} \Delta_M^2 \delta B.$$

The second term arises from the finite capacity constraint, which suppresses high-frequency modes.

9.3 IX.3 Modified Dispersion Relation

Let $\delta B \sim e^{i(px - \omega t)}$. Substituting into the linearized equation yields:

$$-\omega^2 = -p^2 + \frac{p^4}{C_0}.$$

Thus the dispersion relation is:

$$\omega^2 = p^2 - \frac{p^4}{C_0}.$$

Theorem 10 (Modified Dispersion). *Gravitational waves satisfy the dispersion relation*

$$\omega(p) = p\sqrt{1 - \frac{p^2}{C_0}}.$$

9.4 IX.4 Existence of a UV Cutoff

The dispersion relation implies:

$$\omega^2 \geq 0 \implies p^2 \leq C_0.$$

Thus the maximum allowed momentum is:

$$p_{\max} = \sqrt{C_0}.$$

The maximum frequency is:

$$\omega_{\max} = \sqrt{C_0}.$$

Theorem 11 (UV Gravitational Wave Cutoff). *No gravitational wave mode can exceed the frequency*

$$\omega_{\max} = \sqrt{C_0}.$$

9.5 IX.5 Physical Interpretation

The finite relational capacity C_0 acts as a UV regulator. High-frequency gravitational waves require large curvature, which in turn requires large operator norm. Because the operator norm is bounded, high-frequency modes are suppressed.

Thus:

- low-frequency gravitational waves propagate normally,
- intermediate-frequency waves propagate with dispersion,
- high-frequency waves are forbidden.

9.6 IX.6 Observational Consequences

Cutoff in LIGO/Virgo/KAGRA Band

The cutoff frequency may lie above or within the sensitivity band of current detectors. If within the band, one expects:

- suppression of high-frequency components,
- distortion of merger waveforms,
- deviation from GR predictions at high frequencies.

Primordial Gravitational Waves

The cutoff implies:

- no arbitrarily high-frequency primordial modes,
- a natural suppression of UV contributions,
- a modified tensor spectrum in the early universe.

9.7 IX.7 Summary

- Finite relational capacity modifies gravitational wave propagation.
- The dispersion relation becomes

$$\omega^2 = p^2 - \frac{p^4}{C_0}.$$

- High-frequency modes are suppressed.
- The maximum gravitational wave frequency is

$$\omega_{\max} = \sqrt{C_0}.$$

- This yields testable predictions for gravitational wave astronomy.

Chapter 10

Addendum X: Experimental Falsifiability of the Relational Capacity Framework

10.1 X.1 Overview

A physical theory must be falsifiable. This addendum enumerates the concrete, testable predictions of the First Kernel framework. Each prediction arises directly from finite relational capacity and the operator-theoretic structure of the trapped sector P_H .

The predictions fall into four categories:

1. gravitational wave signatures,
2. cosmological expansion behavior,
3. dark matter structure,
4. particle physics relations.

Each category contains at least one decisive, falsifiable prediction.

10.2 X.2 Gravitational Wave Falsifiability

UV Cutoff

The dispersion relation derived in Addendum IX,

$$\omega^2 = p^2 - \frac{p^4}{C_0},$$

implies a maximum gravitational wave frequency:

$$\omega_{\max} = \sqrt{C_0}.$$

If gravitational waves are observed above $\sqrt{C_0}$, the theory is falsified.

High-Frequency Suppression

The amplitude of high-frequency modes must satisfy:

$$A(\omega) \sim \sqrt{1 - \frac{\omega^2}{C_0}}.$$

Thus:

- high-frequency components of merger waveforms must be suppressed,
- ringdown spectra must deviate from GR at high frequencies,
- primordial gravitational waves must show a UV cutoff.

Falsification Criterion

Detection of unsuppressed high-frequency gravitational waves falsifies the theory.

10.3 X.3 Cosmological Falsifiability

Early-Time Slow Expansion

The theory predicts:

$$\frac{\dot{a}}{a}(0) \approx 0.$$

Thus:

- no inflation,
- slow early expansion,
- horizon-scale uniformity from capacity saturation.

Late-Time Fast Expansion

As trapped capacity decreases:

$$\frac{\dot{a}}{a}(t) \uparrow.$$

Thus:

- late-time acceleration is inevitable,
- the Hubble tension is structural,
- the expansion rate must follow a specific curve determined by $P_{\text{trap}}(t)$.

Falsification Criterion

If the expansion history deviates from the predicted capacity-driven curve, the theory is falsified.

10.4 X.4 Dark Matter Falsifiability

Cored Halos

The trapped sector satisfies:

$$\partial_r P_H(0, t) = 0.$$

Thus dark matter halos must be cored, not cuspy.

Observation of universal cuspy halos falsifies the theory.

Dark Matter Deficit Bubbles

Macro-ideal rupture produces deficit bubbles with radius:

$$r(t) \sim \sqrt{4D_{\text{net}}t}.$$

Thus:

- AGN cavities,
- cluster voids,
- dark matter holes

must follow this diffusion law.

Falsification Criterion

If deficit bubbles do not follow the predicted diffusion scaling, the theory is falsified.

10.5 X.5 Particle Physics Falsifiability

Three Fermion Generations

Addendum IV showed:

$$n_{\text{gen}} = 3.$$

Thus:

Discovery of a fourth fermion generation falsifies the theory.

Mass Hierarchy Scaling

Masses scale as:

$$m_f \propto P_H^{3/2}.$$

Thus:

- fermion masses must follow a congestion law,
- deviations from the predicted scaling falsify the theory.

Gauge Coupling Near-Unification

Addendum VII predicts:

$$g_3^2(\Lambda) \approx g_2^2(\Lambda) \approx g_1^2(\Lambda).$$

Thus:

If couplings do not approach near-unification at high energy, the theory is falsified.

10.6 X.6 Summary of Falsifiable Predictions

- UV gravitational wave cutoff at $\omega_{\max} = \sqrt{C_0}$.
- High-frequency gravitational wave suppression.
- Slow early expansion, fast late expansion.
- Cored dark matter halos.
- Dark matter deficit bubbles with diffusion scaling.
- Exactly three fermion generations.
- Congestion-based mass hierarchy.
- Near-unification of gauge couplings.

Each prediction is decisive. Each prediction is falsifiable. Each prediction arises directly from finite relational capacity.

Chapter 11

Addendum XI: Unified Synthesis of the Relational Operator Framework

11.1 XI.1 Overview

This final addendum synthesizes the results of Addenda I–X into a unified mathematical and physical structure. The First Kernel framework is built on a single foundational principle:

All physical phenomena arise from finite relational capacity encoded in a bounded operator $B(t)$ acting on a composite Hilbert space.

Every emergent structure—geometry, matter, gauge fields, dark matter, black holes, cosmology, gravitational waves—follows from this single constraint.

11.2 XI.2 The Fundamental Constraint

The global Frobenius capacity constraint is:

$$\|B(t)\|_F^2 = C_0.$$

This constraint governs:

- the emergent manifold (Addendum I),
- gravitational dynamics (Addendum II),
- gauge sector structure (Addendum III),
- fermion generation count (Addendum IV),
- mass generation (Addendum V),
- cosmological expansion (Addendum VI),
- renormalization group flow (Addendum VII),

- black hole/dark matter duality (Addendum VIII),
- gravitational wave cutoff (Addendum IX),
- experimental falsifiability (Addendum X).

Thus C_0 is the single physical constant from which all others derive.

11.3 XI.3 Emergent Geometry

Addendum I showed that the graph-to-manifold limit yields a smooth Riemannian manifold (M, g) with metric:

$$d(x, y)^2 = \lim_{t \rightarrow 0^+} -4t \log K_M(x, y; t).$$

Thus geometry is emergent from relational diffusion.

11.4 XI.4 Emergent Gravity

Addendum II showed that the spectral action produces the Einstein–Hilbert term:

$$S_{\text{grav}} = \frac{f_2 \Lambda^2}{16\pi^2} \int_M R \, d\text{vol}_g,$$

with Newton’s constant:

$$G = \frac{\pi}{f_2 \alpha C_0}.$$

Gravity is therefore a macroscopic manifestation of finite relational capacity.

11.5 XI.5 Emergent Gauge Fields

Addendum III showed that the internal algebra:

$$A_F = M_3(\mathbb{C}) \oplus M_2(\mathbb{C}) \oplus \mathbb{C}$$

produces the Standard Model gauge group:

$$SU(3)_c \times SU(2)_L \times U(1)_Y.$$

Gauge fields arise from inner automorphisms of A_F .

11.6 XI.6 Fermion Generations

Addendum IV showed that the index of the Dirac operator is:

$$\text{Ind}(D) = 6,$$

yielding exactly three fermion generations.

This result is structural and unavoidable.

11.7 XI.7 Mass Generation

Addendum V showed that mass arises from congestion in the trapped sector:

$$m_f \propto P_H^{3/2}.$$

The Higgs field is the cross-block elasticity operator:

$$\Phi = P_A B P_B.$$

Thus mass is not fundamental; it is emergent.

11.8 XI.8 Cosmology

Addendum VI showed that cosmic expansion is driven by capacity flow:

$$\frac{\dot{a}}{a} = \frac{P_{\text{horizon}}(t)}{C_0 - P_{\text{trap}}(t)}.$$

This explains:

- slow early expansion,
- fast late expansion,
- primordial black holes,
- cored dark matter halos,
- dark matter deficit bubbles,
- high- ℓ CMB damping.

11.9 XI.9 Renormalization Group

Addendum VII showed that RG flow arises from path multiplicities:

$$g_k^2(\ell) \propto \frac{1}{N_k(\ell)}.$$

This yields:

- QCD asymptotic freedom,
- weak coupling behavior,
- hypercharge growth,
- near-unification at high energy.

11.10 XI.10 Black Hole / Dark Matter Duality

Addendum VIII showed that the trapped sector P_H is a C^* -ideal:

$$P_H \triangleleft M_3(\mathbb{C}).$$

- Non-maximal ideal $\Rightarrow$ dark matter.
- Maximal ideal $\Rightarrow$ black hole.

Thus black holes and dark matter are two phases of the same operator structure.

11.11 XI.11 Gravitational Wave Cutoff

Addendum IX showed that gravitational waves satisfy:

$$\omega^2 = p^2 - \frac{p^4}{C_0}.$$

Thus:

$$\omega_{\max} = \sqrt{C_0}.$$

This is a decisive, falsifiable prediction.

11.12 XI.12 Falsifiability

Addendum X enumerated the falsifiable predictions:

- UV gravitational wave cutoff,
- cored halos,
- deficit bubble diffusion,
- three fermion generations,
- congestion-based mass hierarchy,
- near-unification of couplings,
- capacity-driven expansion history.

Each prediction is decisive.

11.13 XI.13 Unified Picture

All physical phenomena emerge from:

Finite relational capacity acting on a composite Hilbert space.

This single constraint produces:

- geometry,
- gravity,
- gauge fields,
- fermions,
- masses,
- cosmology,
- dark matter,
- black holes,
- gravitational waves,
- falsifiable predictions.

The First Kernel is therefore a unified, mathematically rigorous, experimentally testable framework.

Closing Remarks

The First Kernel Addenda consolidate the mathematical foundations of the relational operator framework. They address critiques, strengthen derivations, and demonstrate that the theory is internally consistent, externally predictive, and experimentally falsifiable.

The addenda do not replace the main manuscript. They support it.

Together, the main paper and this supplementary volume form a complete presentation of the theory: its foundations, its consequences, and its testable predictions.

The First Kernel is not a reformulation of known physics. It is a reconstruction of physical law from relational capacity.

The Unified Relational Operator Framework

The expanded master equation for the entire relational operator framework, subject to the fundamental global Frobenius capacity constraint, can be expressed as follows.

The Total Spectral Action

The total action expands via the asymptotic Seeley-DeWitt spectral expansion as:

$$\begin{aligned} S_{\text{total}} &= \text{Tr} \left(f \left(\frac{D^2}{\Lambda^2} \right) \right) + \int_M \text{Tr}_{\mathcal{H}} (\Phi^\dagger (\Delta_M - \mu^2 + \lambda |\Phi|^2) \Phi) d\text{vol}_g \\ &= \frac{f_0 \Lambda^4}{16\pi^2} \int_M d\text{vol}_g + \frac{f_2 \alpha C_0}{16\pi^2} \int_M R d\text{vol}_g \\ &\quad + \frac{f_4}{16\pi^2} \int_M \left(\frac{1}{180} R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} - \frac{1}{180} R_{\mu\nu} R^{\mu\nu} + \frac{1}{72} R^2 \right) d\text{vol}_g + \mathcal{O}(C_0^{-1}) \end{aligned} \quad (11.1)$$

The Foundational Capacity Constraint

All emergent physics is bounded by the conserved global operator norm ceiling:

$$\|B(t)\|_F^2 \equiv \text{Tr}_{\mathcal{H}}(B^\dagger(t)B(t)) = C_0 \quad (11.2)$$

Key Framework Components

- **Fluctuated Dirac Operator:** $D = D_G + A + JAJ^{-1}$ acting on the composite Hilbert space $\mathcal{H} = L^2(M, S) \otimes (\mathbb{C}^3 \oplus \mathbb{C}^2 \oplus \mathbb{C})$.
- **Spectral UV Scale:** $\Lambda = \sqrt{\alpha C_0}$.
- **Internal Gauge Symmetry:** $\mathcal{A}_F = M_3(\mathbb{C}) \oplus M_2(\mathbb{C}) \oplus \mathbb{C} \implies SU(3)_c \times SU(2)_L \times U(1)_Y$.
- **Index Constraint (Generations):** $\text{Ind}(D) = \text{Ind}(D_G) \cdot \text{Ind}(D_F) = 1 \cdot 6 = 6 \implies n_{\text{gen}} = 3$.
- **Congestion Potential & Masses:** $\mu^2 = g^2 P_H$, $\lambda = \frac{C_0}{C_{\text{junction}}}$, and $m_f = y_f v \propto P_H^{3/2}$.
- **Cosmology & Dispersion:** Cosmic expansion is driven by $\frac{\dot{a}}{a} = \frac{P_{\text{horizon}}(t)}{C_0 - P_{\text{trap}}(t)}$, yielding the modified gravitational wave dispersion $\omega^2(p) = p^2 - \frac{p^4}{C_0}$ with a hard UV cutoff at $\omega_{\text{max}} = \sqrt{C_0}$.