A.

Patients newly treated with GLP-1 RA between 2010–2023 (N = 220,479)

- **Excluded**
- <12 months continuous enrollment (n= 47,499)
- Age <18 years (n=205)
- Prior GLP-1 RA or statin use (n=23,984)
- Baseline chronic liver disease etiologies (HBV, HCV, autoimmune hepatitis, alcohol-associated liver disease, cryptogenic liver disease, and HIV (n=3,345)
- SLD diagnosed after index date (n=39,843)
- Statin use prior to GLP-1 initiation (58,197)
- Statin add-on after 12 months (n=3,685)

Final GLP-1 RA based cohort (N = 47,406)

GLP-RA users with statin add-on within 12 months (N=5,991)

GLP-RA users without statin add-on during follow-up (N=37,730)

GLP-RA-statin combination users after time-conditional PS matching (N=4,846)

GLP-RA monotherapy users after time-conditional PS matching (N=19,327)

B.

Patients newly treated with statin between 2010–2023 (N = 893,776)

- **Excluded**
- <12 months continuous enrollment (n=378,137)
- Age <18 years (n=288)
- Prior GLP-1 RA or statin use (n=40,727)
- Baseline chronic liver disease etiologies (HBV, HCV, autoimmune hepatitis, alcohol-associated liver disease, cryptogenic liver disease, and HIV (n=6,616)
- SLD diagnosed after index date (n=215,162)
- GLP-1 RA use prior to statin initiation (13,921)
- GLP-1 RA add-on after 12 months (n=14,350)

Final statin based cohort (N = 238,925)

Statin users with GLP-1 RA add-on within 12 months (n=8,858)

Statin users without GLP-1 RA add-on during follow-up (n=215,717)

GLP-RA-statin combination users after time-conditional PS matching (n=7,653)

Statin monotherapy users after time-conditional PS matching (n=30,485)