

1 **Acoustic and perceptual impact of face masks on**
2 **speech: A scoping review**

3 Gursharan Badh¹, Thea Knowles^{1*}

4 *¹ Department of Communicative Disorders & Sciences, University at Buffalo, Buffalo, NY 14215, USA*

5

6 *Corresponding author

7 E-mail: theaknow@buffalo.edu

8 **Abstract**

9 During the COVID-19 pandemic, personal protective equipment such as facial masks
10 and coverings were mandated all over the globe to protect against the virus.
11 Although the primary aim of wearing face masks is to protect against viral
12 transmission, they pose a potential burden on communication. The purpose of this
13 scoping review was to identify the state of the evidence of the effect of facial
14 coverings on acoustic and perceptual speech outcomes. The scoping review followed
15 the framework created by Arksey & O'Malley (2005) and the Preferred Reporting
16 Items for Systematic Reviews and Meta-Analyses Extension for Scoping Reviews
17 guidelines (PRISMA-ScR; Tricco et al., 2018). The search was completed in May 2021
18 across the following databases: PubMed, EMBASE, PsycINFO, Web of Science, and
19 Google Scholar. A total of 3,846 records were retrieved from the database search.
20 Following the removal of duplicates, 3,479 remained for the title/abstract screen
21 and 149 were selected for the full-text review. Of these, 52 were included in the final
22 review and relevant data were extracted. The 52 articles included in the final review
23 consisted of; 11 studied perceptual outcomes only, 16 studied acoustic outcomes
24 only, and 14 studied both perceptual and acoustic outcomes. 13 of these
25 investigated acoustic features that could be used for mask classification. Although
26 the findings varied from article to article, many trends stood out. Many articles
27 revealed that face masks act as a low pass filter, dampening sounds at higher
28 frequencies; however, the frequency range and the degree of attenuation varied
29 based on face mask type. All but five articles that reported on perceptual outcomes

showed a common trend that wearing a face mask was associated with poorer speech intelligibility. The findings of the scoping review provided evidence that facial coverings negatively impacted speech intelligibility, which is likely due to a combination of auditory and visual cue degradation. Due to the continued prevalence of mask use, how facial coverings affect a wider variety of speaker populations, such as those with communication impairments, and strategies for overcoming communication challenges should be explored.

Introduction

The COVID-19 pandemic necessitated several restrictions in the interest of protecting public health, including the wearing of face masks by the general population. Throughout the pandemic, mask mandates and policies have varied from state to state and across the globe [1,2]. During major waves, the United States Centers for Disease Control and Prevention (CDC) recommendations were for all unvaccinated individuals over the the age of 2 to wear face masks indoors, and for vaccinated individuals to wear face masks in areas of high transmission. Mask recommendations from the CDC include masks that completely cover the nose and mouth and are made of two or more layers of breathable fabric [3]. Early anecdotal reports highlighted the unique challenges to communication brought about by wearing a covering over one's mouth and nose. While face masks had previously been used to protect against disease in, for example, health care settings, never before had their use been so highly encouraged across a vast population.

51 While the COVID-19 pandemic has resulted in a surge of new research focusing on
52 the effects of face coverings on communication, attempts at characterizing these
53 challenges date back many decades. [4] found that individuals working in toxic
54 environments indicated that they would often remove respirators since talking with
55 respirators was too difficult. With face mask use now ubiquitous, it is important to
56 investigate their impact on everyday communicative settings. A recent literature
57 review investigating the effects of respirators on speech, found that respirators
58 impact speech intelligibility and verbal communication ([5]). Round et al. 2021
59 literature review related findings back to the burden in medical settings and many
60 reports have been on settings such as professional healthcare or occupational
61 settings in which a face mask was necessary.

62 Respirators, which are devices used to protect the wearer from inhaling unwanted
63 toxic particles (CDC), likely impact communication due to a combination of acoustic
64 and visual disturbances. Respirators and anything else that blocks the mouth and
65 nose has the potential to impact the speech signal as well as block or distort visual
66 facial cues for listeners. These characteristics, and consequences, may be shared by
67 other types of face coverings, such as alternative coverings that are now more
68 frequently worn due to the COVID-19 pandemic. Given the widespread use of face
69 coverings in the general public, a better understanding of their impact on
70 communication is needed.

71 The purpose of this scoping review is to evaluate the state of the evidence at this
72 time regarding how face coverings, including those that were recommended for and

available to the public and not exclusively health care professionals, impact transmission of the speech signal by characterizing their acoustic and perceptual consequences. In the sections following we discuss the role of masks, standardization of masks and the potential burden of face masks on communication. To the authors' knowledge, this is the most comprehensive review on the topic to date. The following sections describe the role of face masks, how their use may be characterized and standardized, and a description of the potential burden they pose on communication.

Role of masks

Masks have long been recommended in certain healthcare settings for the purpose of protecting against the spread of disease, but facial coverings serve additional roles as well. This section briefly reviews other forms of facial coverings and the range of purposes. We are adopting a broad definition of facewear as used and described in more detail in [6].

The CDC define a respirator as "a personal protective device that is worn on the face or head and covers at least the nose and mouth" [7]. The use of a respirator is recommended not only to prevent the spread of airborne disease, but also to prevent inhalation of hazardous particles that may be transmitted via gases or vapors. The CDC has outlined seven types of respiratory protective devices: filtering facepiece respirators, elastomeric respirators (half and full facepiece), powered air-purifying respirators, supplied-air respirators, self-contained breathing apparatus, and combination respirators [8]. Of these, all but the filtering facepiece respirators

(which includes the N95) are reusable and either supply air or make use of a filtration cartridge or canister. These may be used not only in healthcare settings, but in military combat in which toxic fumes are a hazard (e.g., gas masks). Filtering facepiece respirators, conversely, are disposable half-facepiece masks that typically provide protection against particles but not vapors or gases.

Full head enclosures designed to protect against head trauma also often cover the face, such as motorcycle helmets. Activities requiring supplied air rely on full head enclosures as well, such as underwater or space travel. Face coverings have been recommended as a means to protect against poor air quality, such as during fires and highly polluted areas ([6]; [9]). Face coverings are commonplace in certain religious settings. For example, it is commonplace in certain Muslim communities for women to veil themselves in a niqab or a burqa, both of which cover part of or all of the face, including the mouth and nose. Certain garments designed to protect against the cold are designed to be worn over the face, such as balaclavas (which may or may not include a mouth hole), scarves, and neck gaiters. Disguise, either for entertainment or for criminal activity, may also use face coverings such as rubber masks, balaclavas, etc.

Standardization of face coverings

Medical grade masks and respirators

Some types of face coverings may be described in terms of a standard degree of filtration. Face coverings designed and produced for use in situations where a high

degree of filtration is necessary (e.g., healthcare settings) will typically be required to meet certain standards. In the US, for example, respirators must meet the standards of the National Institute for Occupational Safety and Health [10] and may fall into one of several classes that describe the degree of resistance and percentage of filtration of suspended particles [11]. For example, N95 and FFP2 masks are both designed to filter 95% of suspended particles and are “approximately equivalent” [11]. KN95 masks do not necessarily meet United States NIOSH standards but are designed to have the same filtration properties [12]. About 60% of the KN95 respirators in the United States are counterfeit and do not meet NIOSH standards [13].

Respirators may be distinguished from the broader class of “face masks” given that they are designed to filter out very small particles in the air. Face masks, generally, do not filter small particles from the air, they do however, provide a barrier from droplets (CCOHS). N95 respirators are the most widely available NIOSH approved respirator [13]. Respirators may be categorized as disposable and non-disposable. Disposable respirators consist of N95 and KN95 masks and pattern like face masks. Non-disposable respirators consist of powered air-purifying respirators and elastomeric half facepiece respirators. These respirators have air-purifying components to them and can filter out particles such as dust, and fumes [14].

Surgical masks are a commonly worn medical grade facial covering. Prior to the COVID-19 pandemic, surgical masks were mainly worn in a medical setting by healthcare personnel as a physical barrier to protect both the patient and the

healthcare personnel [15]. Surgical masks, which may also be referred to as disposable or medical procedural masks, are not NIOSH approved; however, they are cleared for use by the Food and Drug Administration [15].

Dust respirators are a disposable, non-medical class of respirators that are used to protect against dust during activities in which non-toxic particles may be present, such as mowing the grass or woodworking. Although some dust respirators may resemble N95 respirators, dust respirators are not NIOSH approved. Dust respirators offer a one way protection only and are not recommended to be used if being exposed to hazardous environments.

Non-medical grade face coverings

Due to the limited availability of medical grade face coverings at the onset of the COVID-19 pandemic, there was a rise in the recommendation and implementation of non-medical grade masks. An example of these are fabric masks. Fabric masks are now widely available and can be made at home or purchased. Cloth masks may have one layer, multiple layers or multiple layers separated by disposable filters. The level of protection provided from fabric masks are dependent on the layers of fabric and the type of fabric [16]. The CDC recommends wearing cloth masks with multiple layers and a nose wire to ensure proper protection [13]. Neck gaiters are another fabric facial covering that many wear; however, the CDC recommends wearing a gaiter with two layers. During the colder months many may wear scarfs, ski masks and balaclavas; however, the CDC does not consider these substitutes as face masks and recommends these be worn over face masks.

A rise in the use of transparent face masks occurred as a means of providing face coverings that provided visual access to the mouth during spoken communication. Many clear face masks and cloth masks with clear window inserts are available on the market today. These types of masks may be especially helpful when communicating with individuals who are hard of hearing or have a disability, young children who may be learning to read or learning a new language, and individuals who need to see the proper shape of the mouth. The FDA recently approved of a transparent medical face mask and indicated that these should be reserved for professionals and patients who require them [13]. Face shields are another type of transparent face covering. Face shields are typically constructed of materials such as polycarbonate or acetate and consist of a rigid, transparent visor that is often open at the bottom and is attached to a frame worn on the head [17]. They are considered adjunctive personal protective equipment and were not recommended to be used in place of, but rather in addition to face masks by the CDC as their effectiveness is not well established and they are not designed to protect against respiratory droplets [14,17,18].

Potential burden of masks

While in many cases the purpose of masking is to protect from hazardous conditions, a potential indirect burden of wearing a face covering is the detrimental impact it may have on communication. In a survey conducted of medical personnel in a hospital in Toronto, Ontario during the Severe Acute Respiratory Syndrome (SARS) outbreak, 47% of hospital workers indicated that wearing PPE was

182 associated with communication difficulty [19]. Due to an increase in the use of face
183 masks all over the globe, this same communication difficulty may be a burden to
184 many.

185 Face masks may pose a burden to communication for potentially three main
186 reasons. First, relevant visual information is lost due to the covering of the mouth,
187 obscuring lip movements and facial gestures and expression. Second, the mask itself
188 acts as a physical barrier between the listener and the speech source and may
189 absorb or attenuate acoustic information. Third, face coverings may introduce
190 physiological restrictions or behavioral adjustments that may have a bearing on
191 speech. For example, certain masks, such as fitted surgical masks like the N95, may
192 restrict jaw movement resulting in limited oral opening and changes to the filtered
193 speech signal. Wearing the mask may itself indirectly lead to changes in how speech
194 is produced by the wearer, either due to conscious or unconscious knowledge of the
195 previously mentioned barriers. These speech modifications will thus also result in a
196 modified acoustic speech signal. The Institute of Medicine has recommended that
197 there be an increased effort to improve speech intelligibility and reduce
198 communication interference while wearing masks. Although there is a lack of
199 research to support these claims, half-face elastomeric respirators are sometimes
200 marketed with claims of improved communication when manufactured with
201 speaking membranes and/or voice amplifiers [20].

202 As a physical barrier that covers the mouth, some degree of acoustic attenuation is
203 likely to occur with all types of face masks. Acoustic attenuation in this case refers to

the degree of sound energy dampened as a result of this barrier. The degree of attenuation and the frequency components affected are likely dependent on the material of the face mask [21]. When choosing a mask an individual may opt to choose a mask with greater protection; however these masks may be associated with greater reductions of sound transmission [22]. In environments where greater medical protection is necessary such as in a hospital setting, communicating with others may be a challenge.

The impact of facial coverings on communication may be exacerbated in certain populations, such as those susceptible to hearing loss. [23] investigated the impact of facial coverings on populations more susceptible to hearing loss such as older adults. These populations may often rely more on visual cues from the speaker to aid in their understanding of speech; therefore, communicating with someone wearing a face mask may hinder their communication [23]. [24] found that both acoustic and visual cues are fundamental in the listeners ability to recognize and perceive speech and the speaker's content. Individuals who have hearing loss, such as older adults, may have greater difficulty understanding speech in masks which may be attributed to the degraded signal and/or loss of visual cues. Given the widespread use of face masks it is important to consider how face masks pose a potential burden to different populations.

Rationale & Objectives of the current study

The rationale for the current review was to evaluate the state of the evidence regarding the impact of facial coverings on speech transmission. Our primary research questions included the following:

1. What is known from existing literature about the effect of face coverings on speech acoustic outcomes?
2. What is known from existing literature about the effect of face coverings on signal-based speech intelligibility?

While there are likely important visual effects of face coverings on a listeners' ability to understand speech, our review focuses on signal-based outcomes to determine the impact of the coverings on the audio transmission and its impact on listeners.

Methods

Protocol

This scoping review followed the framework posited by [25]. [25]'s framework includes five core stages: 1) identifying the research question, 2) identifying relevant studies, 3) study selection, 4) charting the data, and 5) summarizing the results. The authors followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) Extension for Scoping Reviews guidelines developed by the Enhancing the Quality and Transparency of Health Research Network [26].

Eligibility criteria

No filters were employed on the search in order to limit bias.

Information sources & search

In consultation with a librarian at the University at Buffalo, both authors iteratively developed a set of systematic search terminology. The initial search strategy was constructed with the help of the Yale Medical Subject Heading (MeSH) analyzer for use in PubMed using a combination of title/abstract and subject (MeSH) keywords. The following databases were included, with the search terminology translated for each one: PubMed, Embase, PsycINFO, Web of Science, Google Scholar. The final search was executed on May 10, 2021. Entries were exported into Zotero where duplicates were removed, then exported to the Rayyan software program which was used to manage the screening. The full search strategy for all databases can be found in the Appendix. The search strategy approach is summarized as:

- [Terms to identify acoustic and perceptual outcomes] AND
- [Terms to identify relevant face coverings] AND
- [Terms to identify studies related to speech and or voice]

Selection of sources of evidence

The Rayyan software program was used to manage the screening process ([27]). Screening followed two phases: a title/abstract screen and a full text screen. Both phases were carried out independently by both authors in the following way.

Following the formation of an initial set of inclusion and exclusion criteria, both authors screened a quasirandom subset of 30 articles in order to establish baseline reliability. Agreement at this stage was 96.7% (29/30). Authors then reviewed the articles independently, meeting after every 1000 articles in order to discuss conflicts. After the first 2000 articles, exclusion criteria were slightly modified to account for unanticipated themes, and previously screened articles were reviewed to ensure their fit in the updated criteria. For example, the exclusion of articles in which face masks were used exclusively for speech measurement (e.g., Phonatory Aerodynamic Systems) or treatment (e.g., Continuous Positive Airway Pressure) was added at this stage. The decision to exclude full helmets and full-face respirators (e.g., gas masks) was also made at this stage in order to maintain a focus on the presence of face masks like those recommended by the CDC. Conflicts were resolved through consensus discussion.

Screening and eligibility

Articles passing both levels of screening (title/abstract and full text) met the following criteria: 1) involved the study of at least one physical facial covering, 2) provided quantitative measures of speech production or perception, 3) presented new or original data and 4) were written in English. Exclusion criteria included 1) the study of full head enclosures (e.g., helmets, full or half-face respirators), 2) patents, opinion pieces, or media reports that did not present original data, and 3) occlusions exclusively used as a treatment or measurement tool (e.g., continuous positive airway pressure devices, ventilators, or phonatory aerodynamic systems).

Data charting process

Both authors completed the data charting process by filling out a customized fillable form and spreadsheet. Information extracted from the articles included 1) article information (title, authors, year, country of research), 2) face covering information (number and type of masks studied, whether a baseline no-mask condition was included), 3) speaker participant and speech methodological design information (speech stimuli versus non-speech auditory stimuli, live speakers versus pre-recorded speech, number and demographics of speakers if applicable), 4) listener participant and perceptual methodological design information where applicable (number and demographics of listeners, whether live listeners were included, perceptual conditions), 5) perceptual and/or acoustic outcome measures, and 6) main acoustic and perceptual results.

Data extraction fields

Charted data for all articles reporting on acoustic and/or perceptual outcomes appears in the Appendix. Explanations of the data extraction fields are reported below. No a-priori codes were established given the range of findings across the articles included in the final text review. Primary categories related to the types of masks, participants, audio stimuli, and experimental conditions were charted first in open-text fields. Major categorical themes were later identified and coded in additional columns which are detailed in the results fields below.

- 304 • *Outcome categories:* Articles were coded depending on whether they included
305 outcomes related to **speech acoustics, speech perception, both acoustics**
306 **and perception**, or **speaker classification**.
- 307 • *Number and type of face masks*
- 308 • *Speaker and speech source details:* The number of speakers, speaker gender, and
309 age were charted as they pertained to the study. Sound sources other than live
310 talkers, such as head and torso simulators, were charged. Additional relevant
311 information as it pertained to each study, such as speaker training or profession
312 if applicable, were also included in this section.
- 313 • *Listener details:* As with speaker details, the number of listeners, listener
314 gender, age, as well as other pertinent details for the study such as listener
315 training were charted as applicable. If non-human listeners were used, such as
316 with automatic speech recognition systems, this information was included.
- 317 • *Speech stimuli and conditions:* Details about speech stimuli, such as whether
318 word, sentences, or non-speech audio like pure tones or noise was charted
319 here. This section also included details of the number and type of experimental
320 conditions including mask conditions and, as applicable, other conditions.
- 321 • *Details on main acoustic outcomes*
 - 322 – *Acoustic attenuation and speech intensity outcomes:* This section included
323 any outcomes related to sound attenuation, transmission loss, overall
324 speech intensity, etc.

- *Other main acoustic outcomes:* Details reported here included, for example, outcomes related to changes in voice-quality acoustic outcomes, segmental acoustic outcomes, etc.
- *Acoustic category tallies:* All articles were coded as 1 or 0 with reference to whether they included outcomes related to attenuation, vocal intensity, voice quality, segmental level changes, or other acoustic results. These charting columns were added after the main data extraction had taken place and themes had been identified.
- *Details on main perceptual outcomes:* This section included main results reported on perceptual speech outcomes.
 - *Perceptual category tallies:* All articles were coded as 1 or 0 with reference to whether they included perceptual outcomes related to visual information, noise condition comparisons, listeners with hearing loss, the use of face shields, the use of transparent masks. These charting columns were added after the main data extraction had taken place and themes had been identified.
- *Details on other important results not reported elsewhere:* Additional results details were included here, as appropriate.

Synthesis of Results


All extracted data entered via the custom form were exported into a spreadsheet for analysis. Following the data extraction procedure, the authors completed a secondary charting process to record the unique types of face masks included across

all articles and the range of outcomes that were reported. Mask types were grouped into nine categories related to mask type and representation across the included articles: surgical, KN95, N95, shield, shield + mask combinations, transparent, inclusion of carbon filters, and other. For each mask category, this data synthesis, appearing in the Appendix, summarized the 1) minimum and maximum values of acoustic attenuation and/or 2) a summary of the whether the masks were reported to impact speech perception. These results along with descriptions of the other charted variables, are reported in the sections below.

Results and Discussion


A total of 3846 records were retrieved from the database search. Following the removal of duplicates, 3479 remained for the title/abstract screen and 149 were selected for the full-text review. An additional 5 were found during the data extraction and included in the final review. In total 52 were included in the final review and relevant data were extracted. The search procedures are included in Fig 1.

Figure 1: PRISMA flow diagram.



The majority of articles included in the final review ($n = 38$) were published in 2020 or 2021 and in response to the COVID-19 pandemic, as shown in Fig 2. Fourteen were published prior to 2020. Articles were classified into three main categories corresponding to the outcome data they reported: perceptual outcomes, acoustic outcomes, and acoustic classification.

Figure 2: Timeline of article publication.



370


371 **Types of face coverings**

372 Table 1: Range of main acoustic and perceptual results by mask type.

Mask category	Mask type	Number of articles reporting of acoustic/perceptual outcomes	Acoustic attenuation (RMS, dB SPL) (compared to baseline)	Perceptual results (compared to baseline)	Summary
Surgical	Surgical mask Polypropylene ASTM Level 2 (MediCom 2142) Polypropylene ASTM Level 3 (DemeTECH) Polypropylene (YY/T 0969) (Corey2020) procedure mask (Giuliani2020)	13	5 dB reduction (Atcherson2020inthesearch) 3.6 (3ft) - 4.2 dB (6ft) attenuation compared to no mask * 2 to 8 kHz (Atcherson2021) 3.6 dB (peak) (Corey2020) 3-4 dB reduction over 1.5 kHz (Giuliani2020) 2 dB reduction (Nguyen2021Acoustic) 6 dB reduction (Porschmann2020) ** estimate based on figure no attenuation (wolfe) 14 dB (Saedi2016)	lower % accuracy than no mask condition in individuals with moderate hearing loss (Atcherson2017effect) weak but significant effect of modality on consonant identification (subjects identified more consonants during condition when mouth could be seen) (Fecher2013) masks did NOT have detrimental impact on speech understanding. mendel2008 mask condition was sig. % words correct = 70% (Randazzo) no effect (Toscano2021)	Range of acoustic attenuation: ~0 dB (wolfe) to 14 dB (saedi2016). Change in perceptual accuracy: - reduction: n = 3 - no change: n = 3
Fabric	Fabric mask (various): Variations: with and without filter, single vs. multi-layer; weave/material Cotton jersey (generic) (Corey2020) Cotton plain (handmade) (generic) Cotton/spandex jersey (generic) Cotton/spandex jersey (LASC)(Corey2020) Cotton plain and denim (Jo-Ann)(Corey2020) Cotton percale bedsheet and polyester trim (handmade)(Corey2020)	9	5.1 dB (6ft handmade fabric) - 6.1 dB (3ft handmade fabric with HEPA filter) attenuation compared to no mask * 2 to 8 kHz (Atcherson2021) 4 dB - 12.6 dB (peak) (Corey2020) 14 dB reduction (Porschmann2020) ** estimate based on figure 2-3 dB of attenuation from 4,000 to 8,000 Hz (wolfe)	Cohn2021: no effect on word identification mask condition was sig. % words correct = 71% (Randazzo) sig effect of mask on speech perception (Rudge2020) sig effect (Toscano2021) no significant interaction (truong2021)	Range of acoustic attenuation: ~0 dB (wolfe) to 14 dB (saedi2016). Change in perceptual accuracy: - reduction: n = 3 - no change: n = 3
KN95	KN95	5	8.7 dB reduction (Atcherson2020inthesearch) 6.3 dB attenuation compared to no mask * 2 to 8 kHz * distance (3ft vs 6ft) made no difference * 2 to 8 kHz (Atcherson2021) 4 dB (peak) (Corey2020) 5.2 dB (Nguyen2021acoustic) 8 dB reduction (Porschmann2020) ** estimate based on figure	NA	Range of acoustic attenuation: 4 dB (Corey2020) to 8.7 dB (Atcherson2020inthesearch). Change in perceptual accuracy: NA - reduction: n = NA - no change: n = NA
N95	N95/FFP2/FFP3	8	10.9 dB reduction (Atcherson2020inthesearch) 6.2 dB (6ft) - 6.4 dB (3ft) attenuation compared to no mask * 2 to 8 kHz (Atcherson2021) 5.7 dB (peak) (Corey2020) Reduced intensity by 3- 10 dB over 1kHz (Giuliani2020) 10 dB (Randazzo) 4.3 db (vos) 2-3 dB of attenuation from 4,000 to 8,000 Hz (wolfe)	mask condition was sig. % words correct = 63% (Randazzo) significant at low SNR but insignificant at high SNR (Toscano2021) not significant (vos)	Range of acoustic attenuation: 2 dB (wolfe) to 10.9 dB (Atcherson2020inthesearch). Change in perceptual accuracy: - reduction: n = 2 - no change: n = 1
Shield	Face shield plastic shield (generic) (Atcherson2021) Humanity shield (Rapid Response PPE) (Atcherson 2021) Moog plastic shield with apron (handmade) (Atcherson2021)	5	13.7 (6ft) - 17.2 (3ft) attenuation compared to no mask * 2 to 8 kHz (Atcherson2021) face shields "split" sound propagation, mostly at higher frequencies cariato2021 13.7 dB (peak) (Corey2020) 3 to 6 dB of attenuation from 4,000 to 8,000 Hz (wolfe)	sig effect of mask on speech perception (Rudge2020)	Range of acoustic attenuation: 3 dB (wolfe) to 17.2 dB (Atcherson2021). Change in perceptual accuracy: - reduction: n = 1 - no change: n = 0

Mask category	Mask type	Number of articles reporting of acoustic/perceptual outcomes	Acoustic attenuation (RMS, dB SPL) (compared to baseline)	Perceptual results (compared to baseline)	Summary
Shield + mask combinations	Combination mask + face shield	6	20 dB (sm + face shield) - 29.2 dB (kn95 + face shield and transparent + face shield) dB reduction (Atcherson2020inthesearch) 18.0 (Polypropylene ASTM Level 3 (DemeTECH) + face shield) - 25.7 (Polypropylene ASTM Level 2 (MediCom 2142) + N95 + face shield) attenuation compared to no mask * 2 to 8 kHz (Atcherson2021) face shield + procedure mask: 6-10 reduction over 1 kHz (Giuliani2020) FFP3-vent+shield: 2.5 - 4kHz range: up to 18.3 dB attenuation (muzzi2020) FFP2-ventilated+shield: 12.5 - 2kHz range: up to 9.7 dB attenuation (muzzi2020) n95 + shield: 17.3 db (vos)	n95 + face shield: speech reception threshold ranged from 15 dB - 50 dB; speech discrimination score ranged from 90 - 95% (Bandaru) FFP3 + head visor: no significant different in two environments (office & emergency) but showed significant different in the intensive care unit and operating theatre. higher background noise in ICU and operating theatre. (Hampton2020) speech recog. was sig for n95 +face shield (vos)	Range of acoustic attenuation: 2.5 dB (wolfe) to 29.2 dB (Atcherson2020inthesearch). Change in perceptual accuracy: - reduction: n = 3 - no change: n = 0
Transparent	Transparent face masks: LIST TYPES transparent ("see-through") prototype surgical face mask (Atcherson2017effect) prototype FaceView transparent mask (Atcherson2020inthesearch) Safe 'N' Clear transparent surgical mask (Atcherson2020inthesearch) handmade transparent cloth mask(Atcherson2020inthesearch) The communicator (Safe 'N' Clear) (Atcherson2021) Cotton/polyester blend and vinyl window 1 (handmade) (Atcherson2021) Cotton/polyester blend and vinyl window 2 (handmade) (Atcherson2021) ClearMask (ClearMask LLC) (Atcherson2021) Cloth and vinyl window (handmade) (Corey2020) Cloth and PVC window (UTSDesignStore) (Corey2020)	7	transparent facemasks attenuated more than non-transparent face masks (Atcherson2021inthesearch; atcherson2017effect) 8.5 (6ft) - 16.1 (3ft) attenuation compared to no mask * 2 to 8 kHz (Atcherson2021) 10.8 - 12.5 dB (peak) (Corey2020) 11.32 dB attenuation for transparent, 13.64 dB for opaque (Thibodeau2021) * used only transparent mask and for opaque condition put fabric to cover window Communicator(TM) transparent mask: 8.4 dB ClearMask(TM): 10.9 dB (vos)	%correct: 25 to 100% (Atcherson2017effect) * reported on individuals with no HL, mod HL and SEV HL * also had a condition that included audiovisual component sig effect of mask on speech perception (Rudge2020) performance lower when compared to no mask (Thibodeau2021) * used only transparent mask and for opaque condition put fabric to cover window	Range of acoustic attenuation: 8.4 dB (vos) to 16.1 dB (Atcherson2021). Change in perceptual accuracy: - reduction: n = 3 - no change: n = 0
Carbon filter	carbon filter mask (PM2.5),	2	8.0 (3ft) - 8.4 dB (6ft) attenuation compared to no mask (Atcherson2021) 6 dB reduction (Porschmann2020) ** estimate based on figure	NA	Range of acoustic attenuation: 8 dB to 8.4 dB (Atcherson2021). Change in perceptual accuracy: NA
Other	Other: tape, hoodie, balaclava, rubber mask, niqab, helmet, mask (unspecified) a balaclava with a mouth hole, a balaclava without a mouth hole, a motorcycle helmet, a hooded sweatshirt (hoodie) and scarf combination, a niqāb (full- face Muslim veil), a rubber mask, and a piece of tape across the talker's mouth. (Fecher2013)	3	scarf: 2.5 dB @ 4.1 kHz (Saedi2016) rubber mask: 30 dB at 5.2 kHz. (Saedi2016)	weak but significant effect of modality on consonant identification (subjects identified more consonants during condition when mouth could be seen) (Fecher2013)	Range of acoustic attenuation: 2.5 dB (saedi2016) to 30 dB (saedi2016). Change in perceptual accuracy: - reduction: n = 1 - no change: n = 0

373 Figure 3: Types of face coverings.



374

375 Face coverings included a range of materials, though the majority of studies
 376 investigated one or more of the following: medical grade masks (e.g., surgical masks,
 377 N95 respirators), face shields and other types of transparent coverings, and various
 378 types of fabric coverings. A summary of the mask types used in the included articles
 379 is presented in Fig 3 and Table 1, including a summary of the impacts on acoustic
 380 attenuation and perceptual findings, as applicable.

381 **Perceptual outcomes of masks**

382 In total, 21 articles investigated the perceptual outcomes of face coverings. Of these,
 383 12 reported exclusively on perceptual outcomes, including word recognition and

sentence intelligibility, and 9 articles reported on both acoustic and perceptual outcomes.

While many types of face masks worn by a speaker were associated with poorer listener accuracy, this was mediated by the presence of visual information, type of material, environmental conditions (e.g., noise, reverberation), the presence of visual cues, and listener characteristics (e.g., hearing loss). Five studies investigated the presence of audio-visual cues from masks on speech perception. Three studies included listeners with hearing loss. Eight studies varied the noise level presented to listeners. Based on themes identified in the articles during the data charting process, the findings of these articles are summarized in the sections below.

Auditory-only versus auditory-visual impacts

Studies that explored the combined audio-visual effect of masks found that masks were associated with decreased accuracy when compared to audio-visual recordings of talkers without masks [24,28–32]. In a simulated video-conference call when talkers wore (surgical) face masks, listener participants were less accurate in word identification, less confident, and perceived greater listener effort [28]. Listeners were less accurate in consonant identification in audio-only conditions compared to audio-visual conditions in which talkers mouths and/or faces were covered [24,29,30]. [24] studied audio-only and audio-visual consonant recognition for talkers donning eight distinct types of face concealment. The authors found that, overall, consonant identification accuracy in audio-visual conditions exceeded the audio-only condition, especially in the presence of background noise,

and that this effect was stronger for some types of coverings over others. [29] found that while listener accuracy was lower in the audio-visual condition for three types of face coverings (niqab, surgical mask, balaclava), the overall number of consonant misperceptions was very small (2%). These confusions were driven by relatively few error types. The most prominent error types included confusion of stops with fricatives (/t/ ~ /θ/), stop voicing, and place of articulation for stops, fricatives, and nasals.

Two studies investigated the audio-visual effects of transparent masks on speech perception and included listeners with and without hearing loss [30,31]. Combined, findings suggest that when the talker is visible, transparent masks are associated with increased listener accuracy, but when the talker is not visible, transparent masks are associated with decreased accuracy [30,31]. Additionally, [31] found listeners reported higher confidence and lower concentration required to understand sentences spoken with a transparent mask compared to an opaque mask when the talker was visible. [30] found that visual cues from transparent masks aided individuals with moderate to severe hearing loss but not those with normal hearing.

Relatedly, six studies examined the effect of layering a face shield over a mask on speech perception in an auditory-only context [32–37]. In all cases, layering a face shield on top of a mask was associated with reduced intelligibility. These findings suggest the addition of a face shield disrupts acoustic transmission of the speech signal resulting in poorer listener accuracy. The increased perceptual difficulties in

auditory-only scenarios when a transparent mask or shield is used are likely related to increased acoustic attenuation and/or distortion from the rigid materials [38].

Varying levels of noise

Eight studies explored the effects of different noise levels on intelligibility of speech in masks [24,29,33,36,39–42]. Overall, results suggest intelligibility deficits of masks are worse in the presence of higher noise levels or in environments with greater reverberation. Strategies for improving speech intelligibility in masks included speaking more loudly [36] or more clearly [43] or using amplification such as an in-ear listening device [44] or a microphone [45].


Minimal impact

Although a common trend across all articles was that wearing a mask was typically associated with poorer speech intelligibility compared to not wearing a mask, this was not the case across the board. FiveXX articles reported minimal to no effects of masks on speech perception, most often for surgical masks [29,32,34,39,40,43]. [40] found no difference in intelligibility of speech produced with a surgical mask compared to without, regardless of the presence or absence of background noise. [32] found no differences in listener accuracy (evaluated in quiet) when talkers donned an N95 mask compared to without, but did find lower accuracy when the N95 mask was layered with a face shield. Similarly, [34] reported descriptive statistics on preliminary data suggesting that decreases in accuracy were observed when talkers wore face masks and further when layering with a face shield, they

noted that the relatively modest changes would likely not yield statistically meaningful differences. [29] found that speech with surgical masks, compared to niqabs and balaclavas, were the easiest to understand and were also associated with less acoustic transmission loss. While [39] found that listening effort did not differ significantly as a function of mask type, the authors reported that surgical masks had lowest probability of experiencing greater listening effort followed by fabric and the greatest listening effort was for N95s. [43] found no effect of a fabric face mask conditions when talkers spoke in a habitual speech style, but speech produced with a mask was *more* intelligible than without a mask when talkers were instructed to speak clearly. The authors attributed this to a targeted adaptation approach in which talkers overcompensated for the mask when adapting a clear speech style for listener comprehension.

Acoustic outcomes

Figure 4: Number of articles reporting on acoustic outcomes.



463

464 Of the 52 total included articles, 28 investigated the effect of face coverings on
 465 speech acoustic outcomes, which in most cases characterized the degree of
 466 attenuation of the speech signal. Other acoustic outcomes included overall speech
 467 intensity ($n = 4$), voice quality-related acoustic measures ($n = 5$), and segmental
 468 speech acoustics, such as properties of vowel and fricative productions ($n = 5$). Key
 469 differences in methodological approaches of note were whether the sound source
 470 included a live human talker, pre-recorded human talker, or non-speech sound, such
 471 as frequency sweeps. The following sections include mention of the range of
 472 acoustic characteristics reported in the included articles; the intention of this is to

provide an overview rather than a consensus, and the reader is cautioned to recall the extensive methodological variability in the acoustic results presented below.

The distribution of articles reporting on acoustic outcomes is shown in Fig 4.

Acoustic attenuation

21 articles reported on acoustic attenuation imposed by face coverings. Overall, most articles reporting on acoustic attenuation provided evidence that masks act as a low pass filter by dampening the amplitude of frequency components above 1 to 2 kHz. The range of attenuated frequencies and the degree to which masks dampened sound varied across mask types. A summary of the range of acoustic attenuation findings for each mask type is included in Table 1. Studies that compared multiple masks were consistent in reporting that surgical masks demonstrated the least amount of attenuation overall [21,38,46–49]. Attenuation imposed by surgical masks ranged from 0 - 2 dB in mean spectral amplitude between 1 - 8 kHz [48,50] to 14 dB at 4.5 kHz [51]. The greatest attenuation was observed for transparent masks and face shields, ranging from ~3 dB [50] to up to 29.2 dB [46]. Mask material and weave were reported to be the most important variables when considering the degree of attenuation [21]. [21] and [52] reported that the surgical mask and KN95 masks demonstrated peak attenuation at approximately 4 dB, and N95 respirators peaked at approximately 6 dB. [53] recorded white noise produced via a mannequin loudspeaker positioned 6 feet from a microphone in 4 mask and 4 mask plus face shield conditions. They reported preliminary data, identifying reductions in maximum sound pressure level across a spectral range of 0 to 8kHz. Reductions

ranged from 5 dB SPL (surgical mask only) to 29.2 dB SPL (KN95 mask plus face shield and transparent mask plus face shield). [38] expanded on the preliminary findings from [53], including 15 mask conditions (grouped by non-transparent and transparent types) recorded at a 3 foot and 6 foot distance. They found minimal attenuation below 1kHz, and a wide variation of reductions in the 2 - 8 kHz frequency range, and greater attenuation at the farther distance. Between 2 and 8 kHz, compared to no mask, surgical masks were associated with the least attenuation (~4 dB RMS at both 3 and 6-foot distances), and the Humanity Shield transparent mask was associated with the greatest attenuation (~17 dB RMS at both distances). Adding a face shield resulted in an additional 10 to 16 dB RMS attenuation, resulting in a total attenuation of 18 to 25 dB RMS. [40] reported on attenuation from one talker producing sentences with and without a standard surgical mask at a 1 foot distance under controlled speaking conditions (i.e., intentionally maintaining a steady duration and intensity). The authors found a small but significant difference in average RMS power across the speech spectra, suggesting that the talkers speech was actually less attenuated with the mask on, on the order of 0.21 dB SPL.

Two articles reported on spectral slope or tilt, which may be considered an indirect measures of attenuation as it captures the difference in low and high energy bands. [48] found no changes in spectral slopes (comparing 0-1 kHz to 1 - 8 kHz bands) in sustained vowels, but found higher relative low energy values in the surgical and KN95 masks, consistent with previous accounts of low-pass filtering. [54] reported an impact of surgical masks and N95 masks but not fabric masks on spectral tilt.

These differences were observed when measured via a headset and tabletop microphone.

Other acoustic consequences of face coverings

Overall vocal intensity-related outcomes

While attenuation of higher frequency components of the signal were found to be dampened by most masks, overall changes in vocal intensity were not consistently found. Of the studies reporting acoustic outcomes, 4 reported on overall speech intensity [48,54–56].

Of these, one controlled for behavioral adjustments in response to mask wearing by taking acoustic measurements from prerecorded samples. [55] presented prerecorded speech samples from a customized head and torso simulator loudspeaker with a mouth-to-microphone distance of 8 cm. The authors compared three mask types (surgical mask, FFP2, and transparent mask) and found that while the surgical mask was not associated with differences in speech intensity (extracted from a sustained vowel), the FFP2 and transparent masks resulted in speech signals that were approximately 1.5 dB SPL lower compared to no mask. Three studies reported no significant differences in vocal intensity for any of the face masks studied, including surgical masks, KN95 masks, N95 masks, and cloth masks [54,56]. Stimuli from these studies included sustained vowel production and/or connected speech tasks produced by live talkers. While [56] found no group differences in intensity, they reported that the majority (65% of 60 subjects) produced decreased

speech intensity from sustained vowels on the order of 1 to 2 dB with the surgical mask on, while 35% produced an increased intensity.

While not included in the counts above, two studies reported opposing findings on changes in speech intensity featuring a very small number of trained talkers (1 or 2) whose speech was used for perceptual testing. In both of these studies, the talkers were trained and instructed to carefully control speech intensity in the mask conditions. [36] found that a single talker (a trained researcher reading sentence test material) increased their speech-to-noise ratio compared to varying levels of background noise when wearing standard hospital PPE, which included a fit-tested FFP3 mask plus a visor. In low levels of background noise (i.e., 45 dB, simulating standard office noise), the talker increased their SNR by ~2 dB. This trend generally persisted as noise levels increased, with the exception of the highest noise levels (70 dB, simulating a surgical operating theatre), at which point SNR with and without the face coverings were approximately the same. The authors did not report on any statistical findings for this outcome. [43] reported on speech produced by two talkers producing sentences in three different styles (habitual, emotional, and clear) with and without a fabric mask, and found no differences in intensity across the mask conditions when talkers intentionally controlled their speech style in an attempt to be consistent with and without masks. It is worth reiterating that the talkers in both of these articles were trained and intentionally tried to produce consistent speech across the masks; the evidence suggests that masks alone may not impact speech intensity when this is the case.

Overall, these findings suggest that while vocal intensity differences may exist for heavier, thicker mask materials, these differences are small and inconsistent, and may sometimes be attributable to behavioral adjustments to masks.

Voice-quality related acoustic outcomes

5 articles reported on voice and voice-quality-related characteristics, suggesting inconsistent patterns regarding the impact of masks. Four of these reported on live talkers [48,54,56,57] and one reported on pre-recorded speech samples [55].

Overall, face masks (including surgical masks, FFP2 masks, and transparent masks) were not found to impact fundamental frequency [54–57].

The impact of masks on voice quality characteristics such as harmonics-to-noise ratio (HNR) and cepstral peak prominence smoothed (CPPS) varied substantially across the 5 studies. Three of the five articles found no differences in HNR or CPPS in unmasked versus surgical mask conditions in live talkers ([56]; [57]; [54]). [55], reporting on prerecorded sustained vowels, found that while surgical and FFP2 masks were not associated with significant differences in HNR compared to no mask, transparent masks were. All three mask types were associated with decreases in CPPS. [48] found that wearing either a surgical or KN95 mask was associated with an *increase* in HNR in live talkers producing a sustained /a/ vowel, but neither mask was associated with a change in CPPS (measured in vowel production, sentence reading, and passage reading). The authors suggested these results may be due to talkers subconsciously adopting their phonation style when wearing a mask.

Few differences were reported in jitter and shimmer. Specifically, surgical masks and FFP2 masks were not found to be associated with changes to jitter or shimmer ([56]; [57]; [55]; [54]), while transparent masks were associated with an increase in shimmer but no change in jitter ([55]). [55] also reported an increase in AVQI during pre-recorded read sentences across all mask conditions they included (surgical, FFP2, transparent). [54] reported no change in jitter and shimmer for N95, cloth, nor surgical masks.

Segmental acoustic outcomes

5 articles reported on segmental acoustic outcomes. Of these, XXfour reported on the speech of live talkers [58–61] and one reported on pre-recorded speech samples [55]. XXTwo articles reported on the effects of face coverings on vowel formant measures [55,60], and XXthree reported on fricative characteristics [58,59,61].

[55] reported on first and second formants and formant bandwidths of sustained /a/ vowels in prerecorded speech played via a mannequin loudspeaker wearing three different masks. An increase in first formant frequencies was found for a transparent mask (113 Hz), while no differences were found for surgical or FFP2 masks. There were no differences in first formant bandwidths for any of the masks. A decrease in second formant frequencies was found for both the FFP2 (56 Hz) and transparent mask (113 Hz), but not the surgical mask, and differences in second formant bandwidth was found for the transparent mask only. Findings suggest that transparent masks may amplify low frequency characteristics, while masks made of thicker or harder materials may dampen higher frequency characteristics. [60]

604 reported on ten talkers' production of central vowels in Pahari in three face cover
605 conditions: niqab, helmet, and mask (not specified). The authors reported on effects,
606 but not effect directions for F1, F2, and duration of two central vowels. The authors
607 reported an effect of all three face coverings on first and second formants of /ə/ but
608 not /a:/, and inconsistent effects on vowel durations. The authors did not report the
609 direction of effects on these outcomes, thus it is not possible to draw conclusions
610 from this study on the precise nature of the impact of face coverings on vowel
611 production.

612 The three articles reporting on fricative characteristics measured the impact of face
613 coverings that are often used in order to conceal the face (including, for example,
614 helmets, balaclavas, and party masks in addition to surgical masks). Overall, findings
615 suggest that face coverings of these sorts are associated with lower spectral center
616 of gravity, and inconsistent changes in fricative intensity and skewness and kurtosis
617 ([58]; [59]; [61]). Findings overall suggested greater sound absorption for rubber
618 masks, helmets, and tape covering the mouth compared to surgical masks and cloth
619 coverings (hoodies, balaclavas, and niqabs), leading to a shift in spectral
620 characteristics of fricatives. These spectral changes may have implications for
621 consonant identification. It is worth noting that [29], who is not included in the
622 counts here, also made qualitative statements about acoustic characteristics of
623 consonants produced with niqabs, balaclavas, and surgical masks in order to
624 speculate about reasons for listener consonant misidentification.

Other acoustic outcomes

Additional acoustic outcomes not reported above were investigated in 3 articles. These included temporal speech pause characteristics [54] and the acoustic Speech Transmission Index [22,62]. While mean pause length and variability of pause length did not differ in surgical, N95, and cloth masks compared to baseline, N95s and cloth masks were associated with a higher percentage of pauses.

Two articles reported on the STI [22,62]. The STI is a predictive measure of speech intelligibility that is calculated based on acoustic signal-to-noise ratios in the presence of pink ambient noise. Findings support previous conclusions of the relatively minimal effect of surgical masks, which were found to yield the smallest relative effect on the STI outcomes [22,62]. [22] measured the effects of 11 face coverings on sound propagation in high and low reverberant classroom settings using a logarithmic sine sweep emitted from a loudspeaker. Outcomes included including reverberation time, acoustic clarity, and the STI. The authors found that most face masks studied impacted sound propagation, and these differences were increased in rooms with greater reverberation and for male talkers. Surgical masks were least impactful overall. In contrast to previously reported findings of the increased attenuation imposed by transparent masks [CITE], [22] found that masks with transparent windows did not significantly impact speech transmission, though face shields had the effect of splitting sound propagation directivity in two (above and below the shield). [62] investigated the effect of multiple models of three types of face coverings on the STI (surgical masks, N95 masks, and air-purifying

respirators)¹. Surgical masks were associated with the least impact on the STI, followed by the N95 and then by the air-purifying respirators. Specifically, the surgical masks deviated from the no-mask condition by 3-4%, while the N95 masks deviated by 13 - 17%.

Acoustic classification of masks

13 articles were categorized as reporting outcomes related to acoustic classification of face masks, in which the aim of the authors was to identify whether a speaker was wearing a face mask from a set of acoustic features or algorithms. Of these, seven articles were published in the Proceedings of Interspeech 2020 as a part of the Computational Paralinguistics Mask Sub-Challenge [63–69]. The goal of this challenge was for authors to identify whether a speaker was wearing a surgical mask or not, by using machine learning techniques to identify acoustic features from audio recordings from the Mask Augsburg Speech Corpus (MASC) [70]. The MASC includes recordings of 32 speakers of German engaging in a variety of speech tasks with and without a surgical mask. Speech tasks included passage reading, question responses, reading and repeating words, and spontaneous picture descriptions. Of these, [66] was awarded as the challenge winner for achieving the highest accuracy (80.1% “Unweighted Average Recall”) using spectrogram image classification techniques. A summary of techniques used in this challenge are reported in [71].

¹ Air-purifying respirators were excluded in this review. Therefore, these results are not discussed in detail in this section.

Of the remaining six articles, three also reported on acoustic classification of the MASC data, but were published in conference proceedings other than Interspeech 2020 [72]. One article, published in Interspeech in 2015, reported on a corpus of 8 talkers engaging in sentence reading and spontaneous picture description tasks while wearing one of four types of face coverings: motorcycle helmet, rubber mask, hood and scarf, and a surgical mask [75]. Unlike the other classification articles in this category, the aim of this study was to identify whether an automatic speech recognition system could correctly identify individual talkers across the different face covering conditions.

An in depth summary of the outcomes across these investigations into machine learning algorithms aimed at the identification and prediction of the presence of face coverings is beyond the scope of the present review. Findings suggest, however, that the use and refinement of such algorithms may aid in future descriptions of the acoustic impacts of masks and the ability to monitor the speech of talkers while wearing masks. For a more in depth summary of issues of classification of masks, specifically in the context of the Interspeech 2020 ComParE Mask Sub-Challenge, readers are directed to [71].

Study limitations

This scoping review provides, to the authors' knowledge, the most comprehensive review of the effects of face masks on spoken communication. This review sought to characterize the effects of masks recommended for public use by the CDC at the start of the COVID-19 pandemic. As such, this review excluded certain types of face

coverings such as commercial, non-disposable respirators. For a review of the effects of respirators on speech intelligibility, the reader is directed to [5]. Given the rapid rise in interest in this topic, the authors chose not to restrict the review to exclusively peer-reviewed academic journals; white papers, magazine editorials, and proceedings papers have been included. Furthermore, as a part of the scoping review approach, no attempts were made to evaluate the quality of the articles included in the review. As such, some of the conclusions of the studies included may report on preliminary data or experimental designs in less rigorous settings. The full results of the data extraction, including citations and specific methodological details of interest are included in the Supplemental Materials to aid the reader in drawing their own conclusions. Another limitation is related to the timely nature of the topic. At the time of writing, a secondary search revealed several new potentially relevant articles. An update to the present review will be warranted, as new research related to the effects of face masks is a topic that is only continuing to grow. The purpose of the current review was to evaluate the state of the evidence at the current point in time.

Clinical and Research Implications

Overall, the evidence collected in this review suggest that face masks have been consistently shown to dampen higher frequency information of the speech signal. The effect of face masks on speech perception is less consistent, but studies overall point to a detrimental effect that is worsened by factors such as increased noise and the absence of visual cues. Surgical masks appear to have the least impact on

acoustic attenuation and perceptual outcomes. Most soft masks, such as disposable medical grade masks and fabric masks, dampen the acoustic signal to a lesser degree than transparent masks [22]. The thickness and weave of the face covering material appears to play a role in the acoustic impact of the masks [21], while rigid materials such as those used in transparent masks and face shields lead to even greater degrees of attenuation. Transparent masks and especially face shields with visors, though, may also amplify certain higher resonant frequencies and lead to changes in the direction of acoustic transmission [22,53]. From an auditory perspective, greater dampening appears to be associated with worse speech perception, but the presence of visual cues afforded by clear masks alleviates this effect in some circumstances. Areas of future research would benefit from identifying strategies that aid in overcoming the perceptual challenges of masks, including more behavioral modifications, such as clear speech [43] or loud speech [36] and the use of amplification [44,45].

A noticeable gap in the literature was related to representative sample populations that would carry clinical implications for the field of speech-language pathology and audiology. While a small number of studies ($n = 2$) examined the effects of masks on listeners with hearing loss, none reported on other clinical populations such as those with speech or voice disorders. Given the continued prevalence of mask use and the documented acoustic impact of masks, this is an area that should be addressed in the future.

Summary and Conclusion

This scoping review evaluated the state of the evidence regarding the impact of facial coverings on speech transmission at an acoustic and perceptual level. Fifty-two articles were included in the final review, encompassing a wide range of methodologies. Results suggest that face masks consistently are reported to attenuate higher frequency spectral information in the speech signal above 1 to 2 kHz, though this range was found to vary by mask type. Across articles included in this review, face masks are less consistently reported to impact other acoustic features of speech including vocal intensity, voice-quality related measures, and acoustic-phonetic aspects of speech production. While a common trend among the articles reporting on listener perceptual consequences of talkers wearing face masks suggested poorer overall speech intelligibility. However, the presence and magnitude of this effect varied widely and was subject to changes in background noise levels and listener characteristics. Areas of future work should include a wider range of talker characteristics.

References

[1] Badillo-Goicoechea E, Chang T-H, Kim E, LaRocca S, Morris K, Deng X, et al. Global trends and predictors of face mask usage during the COVID-19 pandemic. BMC Public Health 2021;21:2099. <https://doi.org/10.1186/s12889-021-12175-9>.

750 [2] Fischer CB, Adrien N, Silguero JJ, Hopper JJ, Chowdhury AI, Werler MM. Mask
751 adherence and rate of COVID-19 across the United States. PLOS ONE
752 2021;16:e0249891. <https://doi.org/10.1371/journal.pone.0249891>.

753 [3] CDC. Masks and Respirators 2022. [https://www.cdc.gov/coronavirus/2019-](https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/types-of-masks.html)
754 [ncov/prevent-getting-sick/types-of-masks.html](https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/types-of-masks.html) (accessed March 29, 2022).

755 [4] FAWCETT HH. Speech transmission through respiratory protective devices.
756 American Industrial Hygiene Association Journal 1961;22:170–4.
757 <https://doi.org/10.1080/00028896109343390>.

758 [5] Round M, Isherwood P. Speech intelligibility in respiratory protective equipment
759 - Implications for verbal communication in critical care. Trends in Anaesthesia &
760 Critical Care 2021;36:23–9. <https://doi.org/10.1016/j.tacc.2020.08.006>.

761 [6] Fecher N, Watt D. Speaker discrimination based on 'facewear speech'. IAFPA
762 2014 Conference 2014.

763 [7] CDC. Healthcare Workers: Personal Protective Equipment: Questions and
764 Answers 2020. [https://www.cdc.gov/coronavirus/2019-ncov/hcp/respirator-use-](https://www.cdc.gov/coronavirus/2019-ncov/hcp/respirator-use-faq.html)
765 [faq.html](https://www.cdc.gov/coronavirus/2019-ncov/hcp/respirator-use-faq.html) (accessed March 29, 2022).

766 [8] CDC. Healthcare Workers: Types of Respiratory Protection 2020.
767 [https://www.cdc.gov/coronavirus/2019-ncov/hcp/elastomeric-respirators-](https://www.cdc.gov/coronavirus/2019-ncov/hcp/elastomeric-respirators-strategy/respiratory-protection.html)
768 [strategy/respiratory-protection.html](https://www.cdc.gov/coronavirus/2019-ncov/hcp/elastomeric-respirators-strategy/respiratory-protection.html) (accessed March 29, 2022).

769 [9] Langrish JP, Mills NL, Chan JK, Leseman DL, Aitken RJ, Fokkens PH, et al.
770 Beneficial cardiovascular effects of reducing exposure to particulate air pollution

771 with a simple facemask. Part Fibre Toxicol 2009;6:8. <https://doi.org/10.1186/1743->
772 8977-6-8.

773 [10] CDC. National Institute for Occupational Safety & Health | NIOSH | CDC 2021.
774 <https://www.cdc.gov/niosh/index.htm> (accessed December 7, 2021).

775 [11] Li Y, Liang M, Gao L, Ayaz Ahmed M, Uy JP, Cheng C, et al. Face masks to prevent
776 transmission of COVID-19: A systematic review and meta-analysis. Am J Infect
777 Control 2021;49:900–6. <https://doi.org/10.1016/j.ajic.2020.12.007>.

778 [12] Duncan S, Bodurtha P, Naqvi S. The protective performance of reusable cloth
779 face masks, disposable procedure masks, KN95 masks and N95 respirators:
780 Filtration and total inward leakage. PLOS ONE 2021;16:e0258191.
781 <https://doi.org/10.1371/journal.pone.0258191>.

782 [13] CDC. Types of Masks and Respirators 2020.
783 [https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/types-of-](https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/types-of-masks.html)
784 masks.html (accessed December 7, 2021).

785 [14] CDC. Guidance for wearing masks 2020.
786 <https://www.cdc.gov/coronavirus/2019-nCoV/index.html> (accessed August 10,
787 2021).

788 [15] Rengasamy S, Shaffer R, Williams B, Smit S. A comparison of facemask and
789 respirator filtration test methods. Journal of Occupational and Environmental
790 Hygiene 2017;14:92–103.

791 [16] Konda A, Prakash A, Moss GA, Schmoldt M, Grant GD, Guha S. Aerosol Filtration
792 Efficiency of Common Fabrics Used in Respiratory Cloth Masks. *ACS Nano*
793 2020;14:6339–47. <https://doi.org/10.1021/acsnano.0c03252>.

794 [17] Roberge RJ. Face shields for infection control: A review. *Journal of Occupational*
795 & *Environmental Hygiene* 2016;13:235–42.
796 <https://doi.org/10.1080/15459624.2015.1095302>.

797 [18] CDC. COVID-19 and Your Health 2020.
798 [https://www.cdc.gov/coronavirus/2019-ncov/travelers/face-masks-public-](https://www.cdc.gov/coronavirus/2019-ncov/travelers/face-masks-public-transportation.html)
799 [transportation.html](https://www.cdc.gov/coronavirus/2019-ncov/travelers/face-masks-public-transportation.html) (accessed March 15, 2022).

800 [19] Nickell LA, Crighton EJ, Tracy CS, Al-Enazy H, Bolaji Y, Hanjrah S, et al.
801 Psychosocial effects of SARS on hospital staff: Survey of a large tertiary care
802 institution. *CMAJ* 2004;170:793–8. <https://doi.org/10.1503/cmaj.1031077>.

803 [20] Radonovich LJ, Yanke R, Cheng J, Bender B. Diminished speech intelligibility
804 associated with certain types of respirators worn by healthcare workers. *Journal of*
805 *Occupational and Environmental Hygiene* 2010;7:63–70.

806 [21] Corey RM, Jones U, Singer AC. Acoustic effects of medical, cloth, and transparent
807 face masks on speech signals. *The Journal of the Acoustical Society of America*
808 2020;148:2371. <https://doi.org/10.1121/10.0002279>.

809 [22] Caniato M, Marzi A, Gasparella A. How much COVID-19 face protections
810 influence speech intelligibility in classrooms? *Applied Acoustics* 2021;178:108051.
811 <https://doi.org/10.1016/j.apacoust.2021.108051>.

812 [23] Brotto D, Sorrentino F, Agostinelli A, Lovo E, Montino S, Trevisi P, et al. How
813 great is the negative impact of masking and social distancing and how can we
814 enhance communication skills in the elderly people? *Aging Clinical and*
815 *Experimental Research* 2021;33:1157–61. [https://doi.org/10.1007/s40520-021-](https://doi.org/10.1007/s40520-021-01830-1)
816 [01830-1](https://doi.org/10.1007/s40520-021-01830-1).

817 [24] Fecher N, Watt D. Effects of forensically-realistic facial concealment on
818 auditory-visual consonant recognition in quiet and noise conditions. *Auditory-Visual*
819 *Speech Processing* 2013.

820 [25] Arksey H, O'Malley L. Scoping studies: Towards a methodological framework.
821 *International Journal of Social Research Methodology* 2005;8:19–32.
822 <https://doi.org/10.1080/1364557032000119616>.

823 [26] Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA
824 extension for scoping reviews (PRISMA-ScR): Checklist and explanation. *Annals of*
825 *Internal Medicine* 2018;169:467–73.

826 [27] Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan—a web and
827 mobile app for systematic reviews. *Syst Rev* 2016;5:210.
828 <https://doi.org/10.1186/s13643-016-0384-4>.

829 [28] Giovanelli E, Valzolgher C, Gessa E,... Unmasking the Difficulty of Listening to
830 Talkers With Masks: Lessons from the COVID-19 pandemic. I ... 2021.

831 [29] Llamas C, Harrison P, Donnelly D, Watt D. Effects of different types of face
832 coverings on speech acoustics and intelligibility 2009.

833 [30] Atcherson SR, Mendel LL, Baltimore WJ, Patro C, Lee S, Pousson M, et al. The
834 Effect of Conventional and Transparent Surgical Masks on Speech Understanding in
835 Individuals with and without Hearing Loss. *Journal of the American Academy of*
836 *Audiology* 2017;28:58–67. <https://doi.org/10.3766/jaaa.15151>.

837 [31] Thibodeau LM, Thibodeau-Nielsen RB, Tran CMQ, de Souza Jacob RT.
838 Communicating During COVID-19: The Effect of Transparent Masks for Speech
839 Recognition in Noise. *Ear and Hearing* 2021.

840 [32] Vos TG, Dillon MT, Buss E, Rooth MA, Bucker AL, Dillon S, et al. Influence of
841 Protective Face Coverings on the Speech Recognition of Cochlear Implant Patients.
842 *The Laryngoscope* 2021.

843 [33] Muzzi E, Chermaz C, Castro V, Zaninoni M, Saksida A, a, et al. Short report on the
844 effects of SARS-CoV-2 face protective equipment on verbal communication.
845 *European Archives of Oto-Rhino-Laryngology : Official Journal of the European*
846 *Federation of Oto-Rhino-Laryngological Societies (EUFOS) : Affiliated with the*
847 *German Society for Oto-Rhino-Laryngology - Head and Neck Surgery* 2021:1–6.

848 [34] Wittum K, Feth L, Hoglund E. The effects of surgical masks on speech
849 perception in noise. *Proceedings of Meetings on ...* 2013.

850 [35] Bandaru SV, Augustine AM, Lepcha A, Sebastian S, Gowri M, Philip A, et al. The
851 effects of N95 mask and face shield on speech perception among healthcare workers
852 in the coronavirus disease 2019 pandemic scenario. *J Laryngol Otol* 2020;134:895–
853 8. <https://doi.org/10.1017/s0022215120002108>.

854 [36] Hampton T, Crunkhorn R, Lowe N, Bhat J, Hogg E, Afifi W, et al. The negative
855 impact of wearing personal protective equipment on communication during
856 coronavirus disease 2019. *The Journal of Laryngology and Otology* 2020;134:577–
857 81.

858 [37] Rudge A, Sonneveldt V, Brookes B. The effects of face coverings and remote
859 microphone technology on speech perception in the classroom. *The Moog Center for*
860 *Deaf Education White Paper* (Moog Center for Deaf Education, St Louis, MO) 2020.

861 [38] Atcherson SR, McDowell BR, Howard MP. Acoustic effects of non-transparent
862 and transparent face coverings. *The Journal of the Acoustical Society of America*
863 2021;149:2249–54. <https://doi.org/10.1121/10.0003962>.

864 [39] Bottalico P, Murgia S, Puglisi GE, Astolfi A, Kirk KI. Effect of masks on speech
865 intelligibility in auralized classrooms. *The Journal of the Acoustical Society of*
866 *America* 2020;148:2878.

867 [40] Mendel LL, Gardino JA, Atcherson SR. Speech understanding using surgical
868 masks: A problem in health care? *Journal of the American Academy of Audiology*
869 2008;19:686–95. <https://doi.org/10.3766/jaaa.19.9.4>.

870 [41] Toscano JC, Toscano CM. Effects of face masks on speech recognition in multi-
871 talker babble noise. *PloS One* 2021;16:e0246842.

872 [42] Thomas F, Allen C, Butts W, Rhoades C, Brandon C, Handrahan DL. Does
873 wearing a surgical facemask or N95-respirator impair radio communication? *Air*
874 *Medical Journal* 2011;30:97–102. <https://doi.org/10.1016/j.amj.2010.12.007>.

875 [43] Cohn M, Pycha A, Zellou G. Intelligibility of face-masked speech depends on
876 speaking style: Comparing casual, clear, and emotional speech. *Cognition*
877 2021;210:104570. <https://doi.org/10.1016/j.cognition.2020.104570>.

878 [44] Nguyen DL, Kay-Rivest E, Tewfik MA, Hier M, Lehmann A, re. Association of In-
879 Ear Device Use With Communication Quality Among Individuals Wearing Personal
880 Protective Equipment in a Simulated Operating Room. *JAMA Network Open*
881 2021;4:e216857.

882 [45] Rudge AM, Sonneveldt V, Brookes B. The effects of face coverings and remote
883 microphone technology on speech perception in the classroom. *The Moog Center for*
884 *Deaf Education White Paper* (Moog Center for Deaf Education, St Louis, MO) 2020.

885 [46] Atcherson SR, Finley ET, McDowell BR, Watson C. In the Search for Transparent
886 Face Coverings During the Covid-19 Pandemic. *Audiology Today* 2020;32:20–7.

887 [47] Giuliani N. For Speech Sounds, 6 Feet With a Mask Is Like 12 Feet Without.
888 *Leader Live* 2020.

889 [48] Nguyen DD, McCabe P, Thomas D, Purcell A, Doble M, Novakovic D, et al.
890 Acoustic voice characteristics with and without wearing a facemask. *Scientific*
891 *Reports* 2021;11:5651. <https://doi.org/10.1038/s41598-021-85130-8>.

892 [49] Pörschmann C, Lübeck T, Arend JM. Impact of face masks on voice radiation.
893 *The Journal of the Acoustical Society of America* 2020;148:3663.
894 <https://doi.org/10.1121/10.0002853>.

895 [50] Wolfe J, Smith J, Neumann S, Miller S,... Optimizing communication in schools
896 and other settings during COVID-19. The Hearing ... 2020.

897 [51] Saeidi R, Huhtakallio I, Alku P. Analysis of Face Mask Effect on Speaker
898 Recognition. Interspeech 2016.

899 [52] Goldin A, Weinstein B, Shiman N. How Do Medical Masks Degrade Speech
900 Reception? Hearing Review 2020;27:8–9.

901 [53] Atcherson S, Finley E, McDowell B, Watson C. More Speech Degradations and
902 Considerations in the Search for Transparent Face Coverings During the COVID-19
903 Pandemic 2020.

904 [54] Magee M, Lewis C, Noffs G, Reece H, Chan JC, Zaga CJ, et al. Effects of face masks
905 on acoustic analysis and speech perception: Implications for peri-pandemic
906 protocols. bioRxiv 2020.

907 [55] Maryn Y, Wuyts FL, Zarowski A. Are Acoustic Markers of Voice and Speech
908 Signals Affected by Nose-and-Mouth-Covering Respiratory Protective Masks?
909 Journal of Voice 2021:S0892199721000370.
910 <https://doi.org/10.1016/j.jvoice.2021.01.013>.

911 [56] Fiorella ML, Cavallaro G, Di Nicola V, Quaranta N. Voice Differences When
912 Wearing and Not Wearing a Surgical Mask. Journal of Voice 2021.
913 <https://doi.org/10.1016/j.jvoice.2021.01.026>.

914 [57] Cavallaro G, Di Nicola V, Quaranta N, Fiorella ML. Acoustic voice analysis in the
 915 COVID-19 era. *Acta Otorhinolaryngologica Italica : Organo Ufficiale Della Societa*
 916 *Italiana Di Otorinolaringologia E Chirurgia Cervico-Facciale* 2021;41:1–5.

917 [58] Fecher N, Watt D. Speaking under Cover: The Effect of Face-concealing
 918 Garments on Spectral Properties of Fricatives. *ICPhs* 2011.

919 [59] Fecher N. Spectral properties of fricatives: A forensic approach. *Fourth ISCA*
 920 *Workshop on Experimental Linguistics* 2011.

921 [60] Abbasi K, Khan AQ, Abbasi BA. Effects of Forensically Relevant Face Coverings
 922 on Acoustic Properties of Pahari Central Vowels. *International Review of Social*
 923 *Sciences* 2021;9:211–25.

924 [61] Saigusa J. The Effects of Forensically Relevant Face Coverings on the Acoustic
 925 Properties of Fricatives. *Lifespans and Styles* 2017.

926 [62] Palmiero AJ, Symons D, Morgan JW3, Shaffer RE. Speech intelligibility
 927 assessment of protective facemasks and air-purifying respirators. *Journal of*
 928 *Occupational and Environmental Hygiene* 2016;13:960–8.
 929 <https://doi.org/10.1080/15459624.2016.1200723>.

930 [63] Illium S, Müller R, Sedlmeier A,... Surgical Mask Detection with Convolutional
 931 Neural Networks and Data Augmentations on Spectrograms. *arXiv Preprint arXiv ...*
 932 2020.

933 [64] Klumpp P, Arias-Vergara T,... Surgical mask detection with deep recurrent
 934 phonetic models. *Proc Interspeech ...* 2020.

935 [65] Montacié C, Caraty M. Phonetic, Frame Clustering and Intelligibility Analyses for
 936 the INTERSPEECH 2020 ComParE Challenge. Lateral 2020.

937 [66] Szep J, Hariri S. Paralinguistic Classification of Mask Wearing by Image
 938 Classifiers and Fusion. ... INTERSPEECH Shanghai, China ... 2020.

939 [67] Markitantov M, Dresvyanskiy D, Mamontov D, Kaya H, Minker W, Karpov A.
 940 Ensembling End-to-End Deep Models for Computational Paralinguistics Tasks:
 941 ComParE 2020 Mask and Breathing Sub-Challenges. In: Interspeech 2020, ISCA;
 942 2020, pp. 2072–6. <https://doi.org/10.21437/Interspeech.2020-2666>.

943 [68] Yang Z, An Z, Fan Z, Jing C, Cao H. Exploration of Acoustic and Lexical Cues for
 944 the INTERSPEECH 2020 Computational Paralinguistic Challenge. In: Interspeech
 945 2020, ISCA; 2020, pp. 2092–6. <https://doi.org/10.21437/Interspeech.2020-2999>.

946 [69] Ristea N-C, Ionescu RT. Are you Wearing a Mask? Improving Mask Detection
 947 from Speech Using Augmentation by Cycle-Consistent GANs. In: Interspeech 2020,
 948 ISCA; 2020, pp. 2102–6. <https://doi.org/10.21437/Interspeech.2020-1329>.

949 [70] Schuller B, Batliner A,... The interspeech 2020 computational paralinguistics
 950 challenge: Elderly emotion, breathing & masks. Proceedings ... 2020.

951 [71] Mohamed MM, Nessiem MA, Batliner A, Bergler C, Hantke S, Schmitt M, et al.
 952 Face mask recognition from audio: The MASC database and an overview on the
 953 mask challenge. Pattern Recognition 2022;122:108361.
 954 <https://doi.org/10.1016/j.patcog.2021.108361>.

- 955 [72] Das RK, Li H. Classification of speech with and without face mask using acoustic
956 features. In: 2020 asia-pacific signal and information processing association annual
957 summit and conference (APSIPA ASC), IEEE; 2020, pp. 747–52.
- 958 [73] Wang D, Zou D, Cheng X, Xiao W. Classification of the Mask Augsburg Speech
959 Corpus (MASC) Using the Consistency Learning Method. 2020 5th International ...
960 2020.
- 961 [74] Xu X, Deng J, Zhang Z, Wu C, Schuller B. Identifying surgical-mask speech using
962 deep neural networks on low-level aggregation. Proceedings of the 36th ... 2021.
- 963 [75] Saeidi R, Niemi T, Karppelin H, Pohjalainen J,... Speaker recognition for speech
964 under face cover 2015.