

Doubly Robust Targeted Minimum Loss Based Estimation to Address Sampling Bias in Functional Connectivity Studies

Benjamin Risk¹, Daniel Lidstone², Liwei Wang¹, Stewart Mostofsky², David Benkeser¹, and Mary Beth Nebel²

¹Department of Biostatistics & Bioinformatics, Emory University

²Kennedy Krieger Institute and Johns Hopkins University School of Medicine



8 August 2021

Motion quality control exclusion

- Motion in the scanner produces artifacts (Power et al., 2012).
- Lenient criteria: < 5 min data after removing frames with > 3 mm or 3° from previous frame (Fassbender et al., 2017).
- Strict criteria: mean framewise displacement $> .2$ mm or < 5 min data after excluding FD $> .25$ mm (Ciric et al., 2017).




Figure: Quality control removes 30-83% of children with ASD and 12-60% of typically developing.

The problem: motion exclusion criteria in functional MRI causes sampling bias




Figure: During quality control, **more severe cases of autism are excluded.**

The solution: Deconfounded group difference via doubly robust targeted minimum loss based estimation

- $Y(1)$ is the counterfactual that a participant's scan is usable.
Define a novel parameter of interest:

$$\begin{aligned}\psi^* = & E^* \{ E^* (Y(1) | A = 1, W) | A = 1 \} \\ & - E^* \{ E^* (Y(1) | A = 0, W) | A = 0 \}\end{aligned}$$

- Define our target parameter:

$$\begin{aligned}\psi = & E \{ E(Y | \Delta = 1, A = 1, W) | A = 1 \} \\ & - E \{ E(Y | \Delta = 1, A = 0, W) | A = 0 \}.\end{aligned}$$

- $\psi^* = \psi$ under assumptions:
 - (A1.1) *Conditional Randomization*: for $a = 0, 1$,
 $E^* \{ Y(1) | A = a, W \} = E \{ Y | \Delta = 1, A = a, W \}.$
 - (A1.2) *Positivity*: for $a = 0, 1$ and all possible w ,
 $P(\Delta = 1 | A = a, W = w) > 0.$
 - (A1.3) *Consistency*: for all i such that $\Delta_i = 1, Y_i(1) = Y_i,$

- We call the target parameter ψ the **deconfounded group difference**.
- Estimate using doubly robust targeted minimum loss based estimation:
 - Propensity model predicting probability of inclusion to upweight usable data with small probabilities of inclusion.
 - Outcome model fit to usable data to predict functional connectivity in usable and unusable data.
 - Use ensemble of machine learning methods to fit the propensity and outcome models (Van Der Laan et al., 2007).
 - DRTMLE combines estimates in a manner such that mean and SEs robust to mis-specification of one of these models (Benkeser et al., 2017).

Data Analysis

- Resting-state fMRI scans from Kennedy Krieger Institute (either 5:20 or 6:45 seconds in length).
- 153 ASD children and 359 typically developing.
- Use the lenient criteria and residuals from a regression of motion and sex covariates.
- 108 ASD and 300 TD pass lenient criteria.
- SuperLearner with 10-fold CV for propensity and outcome models: `SL.earth`, `SL.glmnet`, `SL.gam`, `SL.glm`, `SL.ranger`, `SL.ridge`, `SL.step`, `SL.step.interaction`, `SL.svm`, `SL.xgboost`.
- Predictors: Primary diagnosis, Head coil, ADHD secondary, Age at scan, Sex, Handedness, Stimulants, Motor overflow, General ability index, Inattention, Hyperactivity, Social responsiveness, Autism diagnostic observation schedule.

Results and discussion




Figure: The deconfounded group difference via DRTMLE reveals more extensive differences between ASD and TD. Z-stats from partial correlations from the group ICA parcellation. Thresholded at $|Z| > 1.96$. Blue: ASD > TD. FDR 0.20: Naive: 2 edges, DTRMLE: 8 edges.

- Participant exclusion due to motion quality control creates large sampling biases.
- We use DRTMLE to estimate the deconfounded group difference in a large study of autism spectrum disorder.
- More extensive differences between ASD and TD when accounting for sampling biases via DRTMLE.
- Future directions: more work on inference. Develop permutation tests.
- Examine sensitivity to model assumptions: randomization (no unmeasured confounders) and e-values (Van Der Weele and Ding, 2017).
- Thank you for watching!
- Additional info: github.com/thebrisklab

References I

- Benkeser, D., Carone, M., Laan, M. J. V. D., and Gilbert, P. B. (2017). Doubly robust nonparametric inference on the average treatment effect. *Biometrika*, 104(4):863–880.
- Ciric, R., Wolf, D. H., Power, J. D., Roalf, D. R., Baum, G. L., Ruparel, K., Shinohara, R. T., Elliott, M. A., Eickhoff, S. B., Davatzikos, C., Gur, R. C., Gur, R. E., Bassett, D. S., and Satterthwaite, T. D. (2017). Benchmarking of participant-level confound regression strategies for the control of motion artifact in studies of functional connectivity. *NeuroImage*, 154:174–187.
- Fassbender, C., Mukherjee, P., and Schweitzer, J. B. (2017). Reprint of: Minimizing noise in pediatric task-based functional MRI; Adolescents with developmental disabilities and typical development. *NeuroImage*, 154:230–239.
- Power, J. D., Barnes, K. A., Snyder, A. Z., Schlaggar, B. L., and Petersen, S. E. (2012). Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion. *NeuroImage*, 59(3):2142–2154.
- Van Der Laan, M. J., Polley, E. C., and Hubbard, A. E. (2007). Super learner. *Statistical Applications in Genetics and Molecular Biology*, 6(1).
- Van Der Weele, T. J. and Ding, P. (2017). Sensitivity analysis in observational research: Introducing the E-Value. *Annals of Internal Medicine*, 167(4):268–274.