Representation theory of the symmetric groups

Based on a lecture series by Stuart Martin.

Michaelmas 2026

This is a fairly gentle introduction to the representation theory of the symmetric group. The flavour will be both algebraic and combinatorial, in particular a first course on representation theory is prerequisite.

Why study representations of symmetric groups? Firstly it is a nice theory. For instance, there is a *canonical* classification of the irreducibles — irreducible representations of $\mathfrak{S}_n$ over $\mathbb{C}$ are parametrised by *partitions* of n. The construction works uniformly for all symmetric groups, much unlike the case of general finite groups where multiple ad hoc constructions are needed for each case.

Another reason is that the theory of symmetric groups serves as fodder for the theory for other algebras that are diagrammatic in flavour, such as (pure) braid groups.

Schur–Weyl duality. Here is a third reason the representation theory of the symmetric group is nice. For V a $\mathbb{C}$-vector space, the tensor power $V^{\otimes 2}$ is a representation of $\mathsf{GL}(V) \times \mathfrak{S}_2$ where $\mathsf{GL}(V)$ acts on each factor, and $\mathfrak{S}_2$ acts by permuting factors (i.e. $v \otimes w \mapsto w \otimes v$). Then $V^{\otimes 2}$ splits into irreducible representations as $\operatorname{Sym}^2 V \oplus \bigwedge^2 V$; there exists a correspondence between representations of $\mathfrak{S}_2$ (namely the trivial and sign representation) and those of $\mathsf{GL}(V)$ (the two factors of $V^{\otimes 2}$ respectively). *Schur–Weyl duality* extends this theory from $\mathfrak{S}_2$ to $\mathfrak{S}_r$. That is, consider the representation $V^{\otimes r}$ of $\mathsf{GL}(V) \times \mathfrak{S}_r$ where $\mathsf{GL}(V)$ acts on each factor and $\mathfrak{S}_r$ permutes the factors. Then Schur–Weyl duality shows $V^{\otimes r} = \bigoplus_i A_i \otimes B_i$ where $\{A_i\}$ are irreducible representations of $\mathsf{GL}(V)$ and $\{B_i\}$ are irreducible representations of $\mathfrak{S}_r$. Again, there is a correspondence $A_i \leftrightarrow B_i$ between some representations of $\mathsf{GL}(V)$ and some representations of $\mathfrak{S}_r$.

Alas this correspondence is not usually bijective — one issue is that $\mathsf{GL}(V)$ has infinitely many irreducible representations while $\mathfrak{S}_r$ has a finite number has a finite number. Another problem is taking inverse powers of determinants.

But the correspondence is *almost* bijective if we stabilise — if we let $\dim(V)$ be large then we recover all representations of $\mathfrak{S}_r$, while if we let r be large then we recover all representations of $\mathsf{GL}(V)$ up to powers of the determinant. We won't look at this.

Topology. A fourth reason comes by taking any topological space X and considering its *configuration space* $C^n X$ (i.e. the moduli of n distinct unordered points in X). The group $\mathfrak{S}_n$ naturally acts on $C^n X$, giving a representation of $\mathfrak{S}_n$ on $H^i(C^n X, \mathbb{C})$.

In positive characteristic. Representations over fields of prime characteristic can be constructed in a 'similar' manner to the characteristic zero theory, which is again unlike the general case for finite groups (modular representation theory tends to be much, much harder in prime characteristic than over $\mathbb{C}$; even constructing characters is hard).

§ 0.1 About. This course on symmetric groups is a part of Part III of the Mathematics tripos at the University of Cambridge. The lectures were delivered in-person, and were transcribed live. The notes can be found online at `https://pas201.user.srcf.net/documents/2026-symmetric-group.pdf`.

Errors and corrections should be communicated to by email to `pas201@srcf.net`.

§ 1 Representations and modules

Lecture 1, 12/10/26

We go over some standard results in the representation theory of a finite group G over a field $\Bbbk$. By *module*, we will always mean finite dimensional left modules.

Definition 1.1. A *representation* of G on a finite dimensional $\Bbbk$-vector space V is any (and all) of the following.

(i) a homomorphism $\rho\colon G \to \mathsf{GL}(V)$;

(ii) a linear action of G on V, i.e. a map $G \times V \to V$ such that for all $g, h \in G$ and $v, w \in V$ we have $(gh)(v) = g(h(v))$, $1(v) = v$, $g(\alpha v + \beta w) = \alpha(gv) + \beta(gw)$;

(iii) a $\Bbbk G$-module structure on V, where $\Bbbk G$ is the *group algebra* (the free $\Bbbk$-module with basis G, with product given by extending the multiplication in G linearly).

Example 1.2. The one-dimensional trivial representation of G, denoted $\mathbf{1}_G$, is given by $V = \Bbbk$ with $g(v) = v$ for all $g \in G$, $v \in V$.

If W is a subrepresentation of V (i.e. a $\Bbbk G$-submodule) then we write $W \leqslant V$. A representation V is *simple* (or *irreducible*) if it is a simple $\Bbbk G$-module, i.e. it is non-zero and its only subrepresentations are $\mathbf{0}$ and V. We say V is *semisimple* (or *complete reducible*) if it is a direct sum of simple representations (this property is closed under taking sub-objects and quotients). We say V is *indecomposable* if it is not a direct sum of proper subrepresentations, we say it is *decomposable* otherwise.

Theorem 1.3 (Maschke). *If $\Bbbk$ has characteristic zero, then every subrepresentation of V is in fact a direct summand. In other words, irreducible and indecomposable representations of V coincide.*

A G-*homomorphism* between two $\Bbbk G$-modules V, W is a $\Bbbk G$-linear map $\alpha\colon V \to W$.

Suppose $\{L_i\}$ is a collection of pairwise non-isomorphic irreducible representations of G. Suppose V is semisimple. Then there is an isomorphism $f\colon \bigoplus L_i^{\oplus n_i} \to V$ for some non-negative integers n_i. The isomorphism is not canonical; but the embedding $f(L_i^{\oplus n_i}) \hookrightarrow V$ *is canonical* — it is the sum of the images of all maps $L_i \to V$, called the L_i-*isotypic piece*. Thus each $\Bbbk G$-module V is *canonically* isomorphic to the direct sum of its isotypic components, and each L_i-isotypic component admits a non-canonical decomposition into a direct sum of L_is.

Theorem 1.4 (Schur's lemma). *Suppose V, W are irreducible representations of G. Any G-homomorphism $V \to W$ is either zero or is an isomorphism. In particular, $\operatorname{End}_G V$ is a division algebra over $\Bbbk$ (if $\Bbbk$ is algebraically closed then $\operatorname{End}_G V \simeq \Bbbk$).*

Lecture 2, 14/10/26