

Derived Equivalences in Algebra and Geometry

Easter 2022

WHAT'S YOUR NEW YEAR'S RESOLUTION?

1920x1080. YOURS?

$$0 \rightarrow \mathcal{O}(-1) \otimes \Omega^1(1) \rightarrow \mathcal{O} \rightarrow \mathcal{O}_\Delta \rightarrow 0$$

*Both the computer scientist and the mathematician were confused
about why nobody invited them to the party that year.*

1	Categories of complexes	3
1.1	Bicomplexes	5
1.2	Three abelian categories	7
1.3	Homotopy and Derived categories	10
2	Coherent sheaves on \mathbb{P}^n	18
2.1	Fourier-Mukai transforms	19
2.2	Koszul resolutions	20
2.3	Beilinson's theorem	22
3	The Bernstein-Gel'fand-Gel'fand correspondence	29
3.1	Twisted functors	29
3.2	BGG resolutions	32
3.3	Descending to triangulated categories	37
3.4	The Tate resolution and Beilinson monads	39
3.5	Koszul duality	40
	References	42

Introduction

Fix a field k , and let X and Ξ be dual k -vector spaces of dimension $n + 1$ with dual bases (x_i) and (ξ_i) respectively. The goal of this exposition is to examine equivalences of various categories that arise naturally in this setting from algebro-geometric constructions. In particular, we look at chain complexes of

- (i) modules over the symmetric algebra $A := \text{Sym}^\bullet(X)$,
- (ii) modules over the exterior algebra $A^! := \bigwedge^\bullet(\Xi)$,
- (iii) coherent sheaves over $\mathbb{P}^n := \text{Proj}(\text{Sym}^\bullet(X))$, the projectivisation of Ξ .

To set the stage, we examine an algorithm to functorially obtain free resolutions of $k[x]$ -modules.

Example 0.1. In the simplest case when X, Ξ are one-dimensional, the data of a module over $A = k[x]$ involves a k -vector space M with a map $M \xrightarrow{x} M$ which can be seen as a complex $F(M)$ of k -vector spaces with differential d of degree 1. In other words, the underlying vector space $F(M) = M \oplus M$ is a module over the graded algebra $A^! = k[\xi]/(\xi^2)$, where the map $F(M) \xrightarrow{\xi} F(M)$ is given by

$$\begin{array}{ccc} M & \oplus & M \\ \searrow 1 & & \searrow \xi \\ M & \oplus & 0 \end{array}.$$

Consider the complex $G(F(M))$ of A -modules

$$\cdots \rightarrow 0 \rightarrow M \otimes_k A \xrightarrow{d \otimes 1 + \xi \otimes x} M \otimes_k A \rightarrow 0 \rightarrow \cdots,$$

concentrated in degrees -1 and 0 . This can be seen as the complex $F(M) \otimes_k A$, but the differential has been ‘twisted’ to remember the $A^!$ -action. This complex is exact everywhere except in degree 0 , where it has cohomology M . Since the modules appearing in it are free, we have recovered a free resolution of M . Moreover, the construction is functorial i.e. an A -module homomorphism $M \rightarrow M'$ naturally induces a chain map $G(F(M)) \rightarrow G(F(M'))$.

This is the first example of what may be called *Koszul duality*, a broad term encompassing various equivalences across algebra, geometry, and representation theory. The duality between symmetric and exterior algebras over finite dimensional vector spaces was first studied by Bernstein, Gel’fand & Gel’fand (1978), who exhibit an adjunction between the categories of complexes of graded modules over A and $A^!$.

Theorem. There are adjoint functors

$$\mathbf{C}(A\text{-grMod}) \xrightleftharpoons[F]{G} \mathbf{C}(A^!\text{-grMod})$$

such that any complex \mathbf{M} of graded A -modules has free resolution $GF(\mathbf{M})$, and any complex \mathbf{N} of graded $A^!$ -modules has injective resolution $FG(\mathbf{N})$.

In Section 3, we look at Eisenbud, Floystad & Schreyer’s (2003) treatment of the Bernstein-Gel’fand-Gel’fand (BGG) correspondence described above. In particular, we have a functorial method to obtain resolutions— this allows for a succinct proof of Hilbert’s theorem on syzygies which we discuss as an application.

To formulate a more precise result on Koszul duality, we need to employ the machinery of Verdier's *derived categories*. We begin by describing this and related constructions from homological algebra in Section 1. In particular, rings and schemes have associated derived categories whose objects are chain complexes of modules and sheaves respectively, considered up to *quasi-isomorphism* (i.e. chain maps that preserve homology). Thus, for instance, a sheaf and its injective resolution are the same object in the derived category. This provides a framework which lets us make clean statements about cohomology, and as we see, is in fact an enhancement of the classical notion of cohomology.

Certain schemes such as Grassmannians have a nice description of the associated derived category. In Section 2 we prove the celebrated theorem of Beilinson (1978) which asserts that the derived category of \mathbb{P}^n is built from $n + 1$ line bundles.

Theorem. The bounded derived category of \mathbb{P}^n is generated by the set $\{\mathcal{O}_{\mathbb{P}^n}, \mathcal{O}_{\mathbb{P}^n}(-1), \dots, \mathcal{O}_{\mathbb{P}^n}(-n)\}$.

For instance, every coherent sheaf \mathcal{A} on \mathbb{P}^n admits a resolution by finite direct sums of the line bundles $\mathcal{O}(-i)$ ($0 \leq i \leq n$). The use of Koszul resolutions and Fourier-Mukai transforms in Beilinson's proof gives an algorithm which lets us directly compute the explicit form of this resolution—in particular we show that this depends only on the cohomology of a few twists of \mathcal{A} . Eisenbud et al. (2003) uses the BGG functors to reconstruct this resolution, and comparing the two constructions provides an algorithm to compute sheaf cohomology.

Acknowledgement. The author is grateful to Dhruv Ranganathan for introducing them to derived categories; to Ismael Sierra del Rio for providing various perspectives on Koszul duality; and to Ian Grojnowski for guidance through the essay.

1 Categories of complexes

We set up the basic framework of homological algebra and derived categories necessary to formulate results in later sections. The material is largely taken from Weibel (2003) and the initial chapters of Huybrechts (2006), with heuristics taken from Thomas (2001).

A category \mathfrak{A} is *additive* if each hom-set $\text{Hom}_{\mathfrak{A}}(A, B)$ has the structure of an abelian group such that composition distributes over addition, \mathfrak{A} has finite products, and there is an object $0 \in \mathfrak{A}$ such that for any $A \in \mathfrak{A}$, $\text{Hom}_{\mathfrak{A}}(A, 0)$ and $\text{Hom}_{\mathfrak{A}}(0, A)$ are the trivial group. In this setting we can make sense of the kernel and cokernel morphisms, which are defined using their usual universal properties.

We say an additive category \mathfrak{A} is *abelian* if every morphism in \mathfrak{A} has a kernel and cokernel, and these behave as expected (i.e. every monomorphism is the kernel of its cokernel and every epimorphism is the cokernel of its kernel). Abelian categories provide the right framework to talk about exact sequences and cohomology, which is the essence of homological algebra. The prototypical example of an abelian category is the category $R\text{-Mod}$ of (left) modules on a ring R .

All the abelian categories occurring in this exposition come from categories of modules over a ring, or sheaves on a topological space. In both the cases, the notions of kernel, cokernel, image, exact sequence, and cohomology are the usual ones.

Example 1.1 (Endomorphism rings). For any object A in an additive category \mathfrak{A} , the group $\text{Hom}_{\mathfrak{A}}(A, A)$ naturally has the structure of a unital ring, where multiplication is given by composition. Then any unital ring R can be seen as an additive category with a single non-zero object $*$, such that $\text{Hom}_{\mathfrak{A}}(*, *) \cong R$. Thus rings are two-object abelian categories!

Example 1.2 (Chain complexes). A *chain complex* \mathbf{A} in an additive category \mathfrak{A} is a sequence of objects $(A^i)_{i \in \mathbb{Z}}$ and morphisms $d^i : A^i \rightarrow A^{i+1}$ (called the *differentials*) such that the compositions $d^i \circ d^{i-1}$ are all 0. We say the object A^i sits in differential degree i . If all but finitely many A^i 's are the zero object, then the chain complex is said to be *bounded*.

A *chain-map* $f : \mathbf{A} \rightarrow \mathbf{B}$ is a sequence $(f^i : A^i \rightarrow B^i)_{i \in \mathbb{Z}}$ of morphisms in \mathfrak{A} such that $f^{i+1} \circ d^i = d^i \circ f^i$ for all i . Then associated to \mathfrak{A} is a category $\mathbf{C}(\mathfrak{A})$ whose objects are chain complexes in \mathfrak{A} , and morphisms are chain-maps. This is naturally an abelian category, where the zero object is the complex which has 0 in every differential degree. Write $\mathbf{C}^b(\mathfrak{A})$, $\mathbf{C}^+(\mathfrak{A})$, and $\mathbf{C}^-(\mathfrak{A})$ for the full subcategories whose objects are bounded complexes, complexes bounded below, and complexes bounded above respectively. These are again examples of abelian categories.

The notions of additive and abelian categories come naturally with notions of functors which preserve the additional structures—these are called *additive* and *exact* functors.

Definition 1.3. A functor $F : \mathfrak{A} \rightarrow \mathfrak{B}$ between additive categories is called an *additive functor* if the maps $F : \text{Hom}_{\mathfrak{A}}(A, B) \rightarrow \text{Hom}_{\mathfrak{B}}(FA, FB)$ are group homomorphisms. We say F is *exact* if, in addition, it sends short exact sequences to short exact sequences.

Example 1.4 (Additive functors on chain complexes). Let \mathfrak{A} be any additive category.

1. There is a natural inclusion $\mathfrak{A} \rightarrow \mathbf{C}(\mathfrak{A})$ which sends an object A to the chain complex A^\bullet where $A^0 = A$, and $A^i = 0$ for $i \neq 0$. This is an exact functor, identifying \mathfrak{A} as a full abelian subcategory of $\mathbf{C}(\mathfrak{A})$.
2. Given $A^\bullet \in \mathbf{C}(\mathfrak{A})$, we define the *translate by 1* of A^\bullet to be the chain complex $A^\bullet[1]$, which has $A^{i+1}, -d^{i+1}$ in degree i . This defines an exact functor $[1] : \mathfrak{A} \rightarrow \mathfrak{A}$ which is an equivalence of categories. Write $[i]$ for the i -fold composition of $[1]$ with itself, and $[-i]$ for the functor inverse to $[i]$.
3. For any $i \in \mathbb{Z}$, the functor $H^i : \mathbf{C}(\mathfrak{A}) \rightarrow \mathfrak{A}$ which sends a complex \mathfrak{A} to its cohomology at A^i is an additive functor. We say \mathbf{A} is *exact at A^i* if $H^i(\mathbf{A}) = 0$. The complex is *acyclic* (or *exact*) if it is exact at every A^i .

1.1 Bicomplexes

If \mathfrak{A} is an abelian category, then so is $\mathbf{C}(\mathfrak{A})$ so we can construct a category $\mathbf{C}(\mathbf{C}(\mathfrak{A}))$ whose objects are *bicomplexes* of the form

$$\begin{array}{ccccccc}
 \vdots & & \vdots & & \vdots & & \\
 \uparrow & & \uparrow & & \uparrow & & \\
 \mathbf{A}^{p+1} : & \cdots \longrightarrow & \mathbf{A}^{p+1,q} \longrightarrow & \mathbf{A}^{p+1,q+1} \longrightarrow & \cdots & & \\
 \uparrow & & \uparrow & & \uparrow & & \\
 \mathbf{A}^p : & \cdots \longrightarrow & \mathbf{A}^{p,q} \longrightarrow & \mathbf{A}^{p,q+1} \longrightarrow & \cdots & & \\
 \uparrow & & \uparrow & & \uparrow & & \\
 \vdots & & \vdots & & \vdots & &
 \end{array} \quad (1)$$

where all the squares commute. Here the horizontal differentials (which we write $d_{>}$ for) come from the internal differentials of the chain complexes $\mathbf{A}^i \in \mathbf{C}(\mathfrak{A})$, while the vertical differentials (which we write d_{\wedge} for) come from the differentials of the complex in $\mathbf{C}(\mathbf{C}(\mathfrak{A}))$.

Such bicomplexes have an associated *total (direct sum) complex* in $\mathbf{C}(\mathfrak{A})$, given by

$$\cdots \rightarrow \bigoplus_{p+q=i-1} \mathbf{A}^{p,q} \longrightarrow \bigoplus_{p+q=i} \mathbf{A}^{p,q} \rightarrow \cdots$$

where the differential sends $a \in \mathbf{A}^{p,q}$ to $d_{\wedge}(a) + (-1)^i d_{>}(a)$. If direct products are taken instead of direct sums, we have the total direct product complex. In the case of bounded bicomplexes the two notions coincide.

Example 1.5. Given two chain complexes $\mathbf{A}, \mathbf{B} \in \mathbf{R}\text{-Mod}$ for some ring R , there is an associated bicomplex $\mathbf{A} \otimes_R \mathbf{B}$ given in degree (p, q) by $\mathbf{A}^p \otimes_R \mathbf{B}^q$. Then the *total tensor product complex* $\text{Tot}(\mathbf{A} \otimes_R \mathbf{B})$ is the total direct sum complex of this bicomplex.

Likewise, defining the bicomplex $\text{Hom}_R(\mathbf{A}, \mathbf{B})$ in degree (p, q) by $\text{Hom}_R(\mathbf{A}^p, \mathbf{B}^q)$. The *total Hom complex* $\text{Tot}(\text{Hom}_R(\mathbf{A}, \mathbf{B}))$ is the total direct product complex of this bicomplex. The usual $\otimes - \text{Hom}$ adjunction extends to these total complexes, the details can be found in Weibel (2003).

1.1.1 Spectral sequences.

Often, one is interested in the cohomology of the total complex associated to a bicomplex. *Spectral sequences* provide a bookkeeping tool for this purpose, and often allow us to extract the cohomology of the bicomplex from the cohomologies of the rows or columns. We will only deal with spectral sequences associated to bicomplexes of modules, so the category \mathfrak{A} is a module category for the purposes of this section.

The *spectral sequence (starting with horizontal cohomology)* \mathbf{E} of a bicomplex (1) is a sequence of *pages* \mathbf{E}_i , where each page has objects of \mathfrak{A} arranged in a grid $E_i^{p,q}$ with specified morphisms which

we now describe. The zeroth page is given by forgetting the vertical differentials in (1), as

$$\begin{array}{ccccccc}
 & & \vdots & & \vdots & & \\
 \dots & \longrightarrow & \mathbf{A}^{p+1,q} & \longrightarrow & \mathbf{A}^{p+1,q+1} & \longrightarrow & \dots \\
 & & \vdots & & \vdots & & \\
 \dots & \longrightarrow & \mathbf{A}^{p,q} & \longrightarrow & \mathbf{A}^{p,q+1} & \longrightarrow & \dots
 \end{array}$$

The objects in the first page will be the cohomologies of the sequences in \mathbf{E}_0 , with maps between them induced by the vertical differentials d_\wedge .

$$\begin{array}{ccccc}
 & \vdots & & \vdots & \\
 & \uparrow & & \uparrow & \\
 \dots & H^q(\mathbf{A}^{p+1}) & & H^{q+1}(\mathbf{A}^{p+1}) & \dots \\
 & \uparrow & & \uparrow & \\
 \dots & H^q(\mathbf{A}^p) & & H^{q+1}(\mathbf{A}^p) & \dots \\
 & \uparrow & & \uparrow & \\
 & \vdots & & \vdots &
 \end{array}$$

The subsequent pages are defined likewise— in particular, the morphisms on the i th page go from $E_i^{p,q}$ to $E_i^{p+i,q-i-1}$ and the compositions of the morphisms, whenever defined, are zero (so that the i th page contains a sequence of complexes.) The objects on E_i are then the cohomologies of the complexes on E_{i-1} , and the morphisms described above are induced from the previous pages.

All the spectral sequences we consider in this exposition are *regular*, i.e. there is an $r \geq 2$ such that the morphisms on every page after E_r are zero. In this case, we have $E_r^{p,q} = E_{r+1}^{p,q} = \dots = E_\infty^{p,q}$ for some object $E_\infty^{p,q} \in \mathfrak{A}$.

Definition 1.6. Given a sequence $H^\bullet = (\dots, H^{-1}, H^0, H^1, \dots)$ of objects in \mathfrak{A} , we say a (regular) spectral sequence E *converges weakly* to H^\bullet if for every i there exists a filtration

$$\dots \subseteq F^2 H^i \subseteq F^1 H^i \subseteq F^0 H^i = H^i$$

such that for every $p \geq 0$ and for every q , we have

$$\frac{F^p H^{p+q}}{F^{p+1} H^{p+q}} \cong E_\infty^{p,q}.$$

If in addition for all i we have $\bigcap_p F^p H^i = 0$ and $H^i = \lim_p H^i / F^p H^i$, then we say the spectral sequence *converges* to H^\bullet .

For a general bicomplex as in (1), not much can be said about convergence of the spectral sequence. However, if the bicomplex is *bounded* i.e. for every i there are only finitely many non-zero $A^{p,q}$ with $p + q = i$ then we have the following result.

Proposition 1.7. If the bicomplex (1) is bounded, then the spectral sequence E converges to the sequence H^\bullet where H^i is the i th homology object of the total (direct sum) complex of the bicomplex.

Proof. See Lemma 12.25.3 of The Stacks project authors (2022). \square

1.2 Three abelian categories

Before proceeding with more constructions from homological algebra, we describe the three abelian categories central to this exposition—these are the module categories of the symmetric and exterior algebra, and the category of coherent sheaves on \mathbb{P}^n . These are defined in the usual way, but we record the constructions involved for completeness of exposition, and to set conventions.

Let k , X , and Ξ be as in the introduction.

1.2.1 Symmetric and exterior algebras.

Definition 1.8. Given an $n + 1$ -dimensional k -vector space V , the *tensor algebra* is the k -vector space

$$T(V) = k \oplus \bigoplus_{i \geq 1} \underbrace{(V \otimes_k V \otimes_k \dots \otimes_k V)}_{i \text{ times}}$$

with a product $\nabla : T(V) \otimes T(V) \rightarrow T(V)$ induced by the natural identifications $V^{\otimes i} \otimes V^{\otimes j} \xrightarrow{\sim} V^{\otimes(i+j)}$. This is an associative algebra with a natural $\mathbb{Z}_{\geq 0}$ -grading.

The *symmetric algebra* $\text{Sym}^\bullet(V)$ and the *exterior algebra* $\bigwedge^\bullet(V)$ are then the graded algebras defined as quotients of $T(V)$ by certain two-sided ideals, namely

$$\text{Sym}^\bullet(V) = \frac{T(V)}{(x \otimes y - y \otimes x \mid x, y \in V)}, \quad \bigwedge^\bullet(V) = \frac{T(V)}{(x \otimes x \mid x \in V)}.$$

Since the ideals are generated by homogeneous elements, these algebras inherit gradings from $T(V)$.

We continue to use ∇ for the product morphism on either algebra, though the corresponding bilinear map on $\bigwedge^\bullet V$ is often written \wedge .

Remark 1.9. We can repeat the above constructions in the category of R -modules for any ring R . In this case, we write $T_R(M)$, $\text{Sym}_R^\bullet(M)$, $\bigwedge_R^\bullet(M)$ respectively for the tensor, symmetric, and exterior algebras over $M \in R\text{-Mod}$. In particular,

$$T(M) = R \oplus \bigoplus_{i \geq 1} \underbrace{(M \otimes_R M \otimes_R \dots \otimes_R M)}_{i \text{ times}}.$$

Since we are primarily concerned with the algebras $A = \text{Sym}^\bullet(X)$ and $A^! = \bigwedge^\bullet(\Xi)$, we redefine the grading on $A^!$ as $A^!_{-i} = \bigwedge^i \Xi$. This amounts to a change of sign from the usual grading, but the convention ensures that the dual vector spaces X and Ξ lie in degrees 1 and -1 in their respective algebras.

Graded modules. A graded A -module is a \mathbb{Z} -graded k -vector space $\mathbf{M} = \bigoplus_i M_i$ with an A -module structure such that $A_i M_j \subseteq M_{i+j}$. For any i , we say the elements of M_i are homogeneous of degree i . A morphism of graded modules then is an A -module homomorphism that preserves the degree of homogeneous elements. Define the category $A\text{-grMod}$ to have finitely generated graded A -modules as its objects, and graded module homomorphisms as the morphisms. This is an abelian category.

The abelian category $A^!\text{-grMod}$ is defined likewise, with objects being finitely generated graded $A^!$ -modules.

1.2.2 The exterior coalgebra

The *exterior coalgebra* on Ξ is defined as the linear dual of $A^!$, written $A^! := \text{Hom}_k(A^!, k)$. $A^!$ has the \mathbb{Z} -grading $A^!_i = \text{Hom}_k(A^!_{-i}, k)$ and is naturally an $A^!$ -module via

$$a \cdot f(a') = (-1)^{\deg a} f(a \wedge a')$$

whenever $a \in A^!$ is homogeneous, and $f \in \text{Hom}(A^!, k)$.

For any vector space N , there is the natural isomorphism of $A^!$ -modules $\text{Hom}_k(A^!, N) \cong A^! \otimes_k N$.

Choosing a basis x_i for X fixes an isomorphism $X \cong \text{Hom}_k(\Xi, k) = A^!_1$, which can be extended to get the isomorphism of graded k -vector spaces

$$A^! = \bigoplus_i \text{Hom}_k(\wedge^i \Xi, k) \cong \bigoplus_i \wedge^i X = \wedge^\bullet(X).$$

In particular, X is a subspace of both $A^!$ and A . This observation is essential in defining the Koszul duality functors, so we write $\tau : A^! \rightarrow A$ for the k -linear map which identifies the subspaces of $A^!$ and A corresponding to X , and is 0 elsewhere.

The coproduct on $A^!$. Being the linear dual of a finite dimensional algebra, $A^!$ has a natural (coassociative counital) coalgebra structure which comes from dualising the (associative unital) product $\nabla : A^! \otimes_k A^! \rightarrow A^!$. This is called the *shuffle coproduct*, and it is helpful to have an explicit description of it which we now describe.

Given a collection of indices $\underline{\alpha} = \{\alpha_1 < \dots < \alpha_i\} \subseteq \{0, \dots, n\}$, write $x_{\underline{\alpha}}$ for the standard basis element of $A^!$ given by $x_{\alpha_1} \wedge x_{\alpha_2} \wedge \dots \wedge x_{\alpha_i}$ (in particular, $x_{\emptyset} = 1$). The vector $\xi_{\underline{\alpha}}$ is defined similarly. We say a tuple $(\underline{\beta}, \underline{\beta}')$ of subsets is a *break* of $\underline{\alpha}$ if $(\beta_1 < \dots < \beta_p, \beta'_1 < \dots < \beta'_q)$ is a permutation of $(\alpha_1 < \dots < \alpha_i)$ (in other words, $\underline{\alpha} = \underline{\beta} \sqcup \underline{\beta}'$). The *sign* of this break, written $\langle \underline{\beta}, \underline{\beta}' \rangle$, is defined to be the sign of the corresponding permutation. Thus we have

$$\nabla(x_{\underline{\beta}} \otimes x_{\underline{\beta}'}) = x_{\underline{\beta}} \wedge x_{\underline{\beta}'} = \langle \underline{\beta}, \underline{\beta}' \rangle x_{\underline{\alpha}}.$$

This allows us to write the coproduct on $A^!$ as

$$\Delta(x_{\underline{\alpha}}) = \sum_{(\underline{\beta}, \underline{\beta}') \in \text{br}(\underline{\alpha})} \langle \underline{\beta}, \underline{\beta}' \rangle x_{\underline{\beta}} \otimes x_{\underline{\beta}'}$$

where $\text{br}(\underline{\alpha})$ is the set of all breaks of $\underline{\alpha}$. Recalling that $A^! \otimes_k A^!$ is \mathbb{Z} -graded with $\bigoplus_{p+q=i} A^!_p \otimes A^!_q$ in degree i , we observe that the map Δ respects grading hence $A^!$ is a *graded coalgebra*.

1.2.3 Graded chain complexes

Objects of $\mathbf{C}(A\text{-grMod})$ are chain complexes of graded A -modules in which the differentials are morphisms in $A\text{-grMod}$ (i.e. A -module homomorphisms which preserve degree). Such an object can be viewed as a \mathbb{Z}^2 -graded k -vector space $\mathbf{M} = \bigoplus_{i,j} M_j^i$ with an endomorphism d (the differential) such that

1. $d \circ d = 0$,
2. d has degree $(1, 0)$ i.e. $d(M_j^i) \subseteq M_j^{i+1}$, and
3. for each $i \in \mathbb{Z}$, $M_\bullet^i = \bigoplus_j M_j^i$ is a graded A -module.

Likewise, an object $\mathbf{N} \in \mathbf{C}(A^!\text{-grMod})$ can be seen as a \mathbb{Z}^2 -graded k -vector space $\bigoplus_{i,j} N_j^i$ with a differential ∂ of degree $(1, 0)$. We shall use the two viewpoints on interchangeably, switching between them whenever convenient to provide a clearer picture. In particular, the ability to view a complex as a single module with additional structure allows for cleaner definitions and proofs, see for instance Theorem 3.2.

For a chain complex $\mathbf{M} = \bigoplus_{i,j} M_j^i$, we say the lower indices denote the *internal* (or *Adam's*) grading, while the upper indices denote the *differential* (or *cohomological*) degree. We use $\langle \cdot \rangle$ to denote shifts in Adam's gradings, continuing to use $[\cdot]$ to denote shifts in differential gradings. Thus for example we have $\mathbf{M}\langle q \rangle_j^i = M_{q+j}^i$.

1.2.4 Coherent sheaves

The details of all the constructions described in this section can be found in Sections II.5 and II.8 of Hartshorne (2008).

For any scheme \mathbf{X} , the category $\mathcal{S}\mathcal{h}\mathbf{X}$ of sheaves on \mathbf{X} is abelian with the usual notions of kernel and cokernel. If the scheme \mathbf{X} is noetherian, then the (co)kernel of a morphism of coherent sheaves is coherent. In this case, the category $\mathcal{C}\mathcal{O}\mathcal{h}\mathbf{X}$ whose objects are coherent sheaves of $\mathcal{O}_{\mathbf{X}}$ -modules is abelian.

The category $\mathcal{C}\mathcal{O}\mathcal{h}\mathbf{X}$ supports an additional operation, the tensor product \otimes . The tensor product of coherent sheaves will always be over the structure sheaf, unless otherwise specified. Given a coherent sheaf \mathcal{E} , we can then define the sheaves of algebras $T\mathcal{E}$, $\text{Sym } \mathcal{E}$, and $\bigwedge^\bullet \mathcal{E}$ as coming from the presheaves which assign to an open set $U \subset \mathbf{X}$ the corresponding tensor operation applied to $\mathcal{E}(U)$ as an $\mathcal{O}_{\mathbf{X}}(U)$ -module.

If $\mathbf{X} \xrightarrow{f} \mathbf{Y}$ is a morphism of noetherian schemes, then the *pullback* of a coherent sheaf on \mathbf{Y} is again coherent. This gives an additive functor

$$f^* : \mathcal{C}\mathcal{O}\mathcal{h}\mathbf{Y} \longrightarrow \mathcal{C}\mathcal{O}\mathcal{h}\mathbf{X}.$$

The pullback functor commutes with the various tensor operations described above.

Sheaves on \mathbb{P}^n . The *projectivisation* of Ξ is the k -scheme defined as $\mathbb{P}^n = \text{Proj}(\text{Sym}(\Xi^\vee))$. Since $\text{Sym}(\Xi^\vee) = A$ is the polynomial algebra on $n + 1$ variables, we see that \mathbb{P}^n is the usual projective n -space over $\text{Spec}(k)$. This is a noetherian scheme, and so $\mathcal{C}\mathcal{O}\mathcal{h}\mathbb{P}^n$ is an abelian category.

Serre gives a correspondence between $A\text{-grMod}$ and $\mathcal{Coh} \mathbb{P}^n$ using a construction similar to Proj, called the ‘sheafification’ of graded modules. This correspondence is functorial, but *not* an equivalence of categories. Indeed, much of Section 2 is dedicated to obtaining a precise formulation of the equivalence between coherent sheaves on \mathbb{P}^n and finitely generated graded A -modules.

In particular, the module $A \in A\text{-grMod}$ corresponds to the structure sheaf $\mathcal{O}_{\mathbb{P}^n}$. Since \mathbb{P}^n is central to the exposition, we omit the subscript \mathbb{P}^n unless there is a possibility of confusion— thus writing \mathcal{O} for the structure sheaf.

The invertible sheaf $\mathcal{O}(i)$ is defined as the sheafification of the graded module $A\langle i \rangle$. In particular, the global sections of $\mathcal{O}(i)$ correspond to degree i polynomials. The various sheaves $\mathcal{O}(i)$ form an abelian group under \otimes , there being natural isomorphisms $\mathcal{O}(i) \otimes \mathcal{O}(j) \cong \mathcal{O}(i+j)$. Note that tensor products of sheaves are always taken over the structure sheaf.

For any sheaf $\mathcal{E} \in \mathcal{Coh} \mathbb{P}^n$, its i th *twist* is defined by $\mathcal{E}(i) = \mathcal{E} \otimes \mathcal{O}(i)$. Since the sheaves $\mathcal{O}(i)$ are flat, taking the i th twist is an exact functor.

The *cotangent sheaf* Ω and the *tangent sheaf* $\mathcal{T} = \mathcal{H}om(\Omega, \mathcal{O})$ are locally free sheaves of rank n which fit in dual exact sequences given by

$$0 \rightarrow \Omega \rightarrow \mathcal{O}(-1)^{\oplus(n+1)} \rightarrow \mathcal{O} \rightarrow 0, \quad 0 \rightarrow \mathcal{O} \rightarrow \mathcal{O}(1)^{\oplus(n+1)} \rightarrow \mathcal{T} \rightarrow 0. \quad (2)$$

These are called the *Euler exact sequences*, and form our primary means of getting hold of the sheaves. In particular, Ω corresponds to the graded module given by the kernel of

$$A^{n+1}\langle -1 \rangle \longrightarrow A; \quad (a_0, \dots, a_n) \longmapsto a_0x_0 + \dots + a_nx_n.$$

1.3 Homotopy and Derived categories

We return to our discussion of homological algebra, to build the framework of *triangulated categories* and the derived category. Before describing these, we provide some motivation as to why we construct them by first reviewing how homological algebra is typically used to find invariants in mathematics.

Given a category \mathcal{C} , we choose for each $X \in \mathcal{C}$ a complex $\Phi(X) \in \mathbf{C}(\mathfrak{A})$ for some abelian category \mathfrak{A} . Then the compositions $H^i \circ \Phi : \mathcal{C} \rightarrow \mathfrak{A}$ allow us to assign a sequence of \mathfrak{A} -objects associated to X , and this assignment (in the good cases) is functorial. The following examples illustrate this.

1. Consider the category \mathcal{Simp} of simplicial complexes (in the sense of algebraic topology), and the functor $F : \mathcal{Simp} \rightarrow \mathbb{Z}\text{-Mod}$ which associates to a simplicial complex $\mathbf{X} \in \mathcal{Simp}$ the chain complex $F(\mathbf{X})$ of abelian groups where the group in degree $-i$ is generated by i -simplices, and the differentials are boundary maps. If $\mathcal{T}\mathcal{r}$ is the category of triangulable topological spaces, then we can choose a map $\Phi : \mathcal{T}\mathcal{r} \rightarrow \mathcal{Simp}$ which associates to each $\mathbf{X} \in \mathcal{T}\mathcal{r}$ a simplicial complex which is its triangulation. The i th *simplicial homology* of \mathbf{X} is defined as $H^{-i}(F(\Phi(\mathbf{X})))$.
2. For a ring R , let $\mathbf{C}_{\text{free}}^-(R\text{-Mod})$ be the full subcategory of $\mathbf{C}^-(R\text{-Mod})$ whose objects are complexes of free R -modules. Choose for each $M \in R\text{-Mod}$, choose a complex $\Phi(M) \in \mathbf{C}_{\text{free}}^-(R\text{-Mod})$ such that $\Phi(M)$ is exact in non-zero degrees and has $H^0(\Phi(M)) = M$. Then for a fixed R -module M , we define the i th Ext functor by $\text{Ext}_R^i(M, -) = H^{-i} \circ \text{Hom}_R(M, -) \circ \Phi$.

3. For a scheme \mathbf{X} , let $\mathbf{C}_{\text{inj}}^+(\mathcal{Q}\mathcal{C}\mathcal{H}\mathbf{X})$ be the full subcategory of $\mathbf{C}^+(\mathcal{Q}\mathcal{C}\mathcal{H}\mathbf{X})$ whose objects are complexes given in each degree by an injective sheaf. If \mathbf{X} is noetherian, then the category $\mathcal{Q}\mathcal{C}\mathcal{H}\mathbf{X}$ has enough injectives (see Section III.3 of Hartshorne (2008)) and for any coherent sheaf \mathcal{E} it is possible to choose a complex $\Phi(\mathcal{E}) \in \mathbf{C}_{\text{inj}}^+(\mathcal{Q}\mathcal{C}\mathcal{H}\mathbf{X})$ such that $\Phi(\mathcal{E})$ is exact in non-zero degrees and has $H^0(\Phi(\mathcal{E})) = \mathcal{E}$. Then the i th *sheaf cohomology* of \mathcal{E} is defined as $H^i(\Gamma(\Phi(\mathcal{E})))$, where Γ is the global sections functor.

All of the constructions above have two ‘problems’– the first is that the choice of Φ is often neither unique nor functorial. However, it turns out in all of the examples that the actual homology computed is functorial and independent of the choice of Φ . For triangulable spaces, proving the independence of homology from choice of triangulation amounts to showing that whenever $\Phi_1(\mathbf{X})$ and $\Phi_2(\mathbf{X})$ are two triangulations of \mathbf{X} , there is a third simplicial complex $\Phi_3(\mathbf{X})$ (a ‘common refinement’ of the two triangulations) such that there is a diagram

$$\begin{array}{ccc} & \Phi^3(\mathbf{X}) & \\ \swarrow & & \searrow \\ \Phi^1(\mathbf{X}) & & \Phi^2(\mathbf{X}) \end{array}$$

in which the two morphisms induce isomorphisms on cohomology.

Definition 1.10. A morphism $f : \mathbf{A} \rightarrow \mathbf{B}$ in $\mathbf{C}(\mathfrak{A})$ is a *quasi-isomorphism* if for all $i \in \mathbb{Z}$, the induced maps $H^i(f) : H^i(\mathbf{A}) \rightarrow H^i(\mathbf{B})$ are isomorphisms.

Cohomology is hard to compute directly in the original category \mathfrak{C} , so we choose to pass through an intermediate category $\mathbf{C}(\mathfrak{A})$. In an ideal situation, we would have, in place of $\mathbf{C}(\mathfrak{A})$, another category $\mathbf{D}(\mathfrak{A})$ such that the homology functors $H^i : \mathbf{D}(\mathfrak{A}) \rightarrow \mathfrak{A}$ are just as easy to compute, but the assignment $\Phi : \mathfrak{C} \rightarrow \mathbf{D}(\mathfrak{A})$ would actually be functorial. The derived category will play this role.

The second problem with cohomology is that it is too crude an invariant– the triangulable spaces $S^1 \times S^1$ and $S^1 \vee S^1 \vee S^2$ have the same simplicial homology, but are clearly not homotopy equivalent. In fact, by Whitehead’s theorem, two simply connected triangulable spaces \mathbf{X}, \mathbf{Y} are homotopy equivalent if and only if there are maps

$$\begin{array}{ccc} & \mathbf{Z} & \\ \swarrow & & \searrow \\ \mathbf{X} & & \mathbf{Y} \end{array}$$

which induce isomorphisms on homology. Thus instead of simply stating if two objects in \mathfrak{C} have isomorphic cohomology, we also wish to specify whether these isomorphisms are induced by morphisms in \mathfrak{C} . Thus a better invariant than cohomology is the associated complex itself, identified with other complexes it maps to via quasi-isomorphisms. In the words of Thomas (2001),

Complexes good, (co)homology bad.

Accordingly, the *derived category* of an abelian category \mathfrak{A} is defined via a universal property.

Theorem 1.11. There is a category $\mathbf{D}(\mathfrak{A})$ with an additive functor $\Theta : \mathbf{C}(\mathfrak{A}) \rightarrow \mathbf{D}(\mathfrak{A})$ universal among additive functors that send quasi-isomorphisms to isomorphisms, i.e. whenever $F : \mathbf{C}(\mathfrak{A}) \rightarrow \mathfrak{D}$ is an additive functor such that every quasi-isomorphism f in $\mathbf{C}(\mathfrak{A})$ is sent to an isomorphism $F(f)$ in \mathfrak{D} , then F factors uniquely through Θ .

$$\begin{array}{ccc} \mathbf{C}(\mathfrak{A}) & \xrightarrow{F} & \mathfrak{D} \\ \downarrow \Theta & \nearrow \exists! & \\ \mathbf{D}(\mathfrak{A}) & & \end{array}$$

Proof. See Chapter 10 of Weibel (2003). □

We describe the construction and state some useful properties. The derived category is constructed in two steps— we first pass to the homotopy category where homotopic morphisms are identified, and then invert quasi-isomorphisms by the operation of localisation.

1.3.1 The homotopy category

To any abelian category \mathfrak{A} , we associate a category $\mathbf{K}(\mathfrak{A})$ (the *homotopy category*) such that there is an additive functor $\mathbf{C}(\mathfrak{A}) \rightarrow \mathbf{K}(\mathfrak{A})$ which H^i factors through.

Definition 1.12. A chain map $A \xrightarrow{f} B$ in $\mathbf{C}(\mathfrak{A})$ is said to be *nullhomotopic* if there are morphisms $s^i : A^i \rightarrow B^{i-1}$ such that $f^i = d^i \circ s^{i+1} + s^i \circ d^i$ for all i . Two chain maps $f, g : A \rightrightarrows B$ are *homotopic* if the difference $f - g$ is nullhomotopic. We call s the *chain homotopy*.

We define the *homotopy category* $\mathbf{K}(\mathfrak{A})$ to have the same objects as $\mathbf{C}(\mathfrak{A})$ and morphisms given by equivalence classes of chain homotopic maps. For any $A, B \in \mathbf{C}(\mathfrak{A})$, the nullhomotopic chain maps form a subgroup $N(\mathfrak{A}, \mathfrak{B})$ of $\text{Hom}_{\mathbf{C}(\mathfrak{A})}(A, B)$. Then we have

$$\text{Hom}_{\mathbf{K}(\mathfrak{A})}(A, B) = \frac{\text{Hom}_{\mathbf{C}(\mathfrak{A})}(A, B)}{N(A, B)}.$$

The categories $\mathbf{K}^+(\mathfrak{A})$, $\mathbf{K}^-(\mathfrak{A})$, and $\mathbf{K}^b(\mathfrak{A})$ are analogously defined from $\mathbf{C}^+(\mathfrak{A})$, $\mathbf{C}^-(\mathfrak{A})$, and $\mathbf{C}^b(\mathfrak{A})$ respectively.

Isomorphisms in $\mathbf{K}(\mathfrak{A})$ are called *homotopy equivalences*.

Proposition 1.13. If f and g are homotopic chain maps, then the induced morphisms $H^i(f)$ and $H^i(g)$ on homology are the same.

Thus techniques of homological algebra fail to distinguish between homotopic morphisms. We often use the result above to show a complex is exact, by showing first that the identity map on the complex is nullhomotopic. Such complexes are called *contractible*.

1.3.2 Localisation and the derived category

Let \mathbf{K} be a triangulated subcategory of $\mathbf{K}(\mathfrak{A})$, and Q be the class of all quasi-isomorphisms in \mathbf{K} . We define the associated derived category $\mathbf{D} = Q^{-1}\mathbf{K}$ by ‘adding inverses’ to quasi-isomorphisms, in the sense of localisations defined below.

Definition 1.14. Let S be a collection of morphisms in a category \mathcal{C} . A localisation of \mathcal{C} with respect to S is a category $S^{-1}\mathcal{C}$ with a functor $q : \mathcal{C} \rightarrow S^{-1}\mathcal{C}$ such that the following hold.

1. For every $s \in S$, $q(s)$ is an isomorphism in $S^{-1}\mathcal{C}$.
2. Any functor $F : \mathcal{C} \rightarrow \mathcal{D}$ such that F takes elements of S to isomorphisms factors uniquely through q .

Example 1.15 (Localisation of rings). Considering a commutative ring R to be an additive category \mathfrak{A} with a single non-zero object $*$ as in Example 1.1, we can choose S to be a multiplicative subset of $R = \text{Hom}_{\mathfrak{A}}(*, *)$. Then $S^{-1}\mathfrak{A}$ is again additive with a single non-zero object $*$. The corresponding ring is precisely the ring of fractions $S^{-1}R$.

We describe the objects and morphisms of $\mathbf{D}(\mathfrak{A})$ explicitly, directing the reader to Weibel (2003) for proofs and further motivations.

- Proposition 1.16.**
1. Under the localisation map $q : \mathbf{K}(\mathfrak{A}) \rightarrow \mathbf{D}(\mathfrak{A})$, the objects of the two categories are identified.
 2. The cohomology functors $H^i : \mathbf{K}(\mathfrak{A}) \rightarrow \mathfrak{A}$ factor through $q : \mathbf{K}(\mathfrak{A}) \rightarrow \mathbf{D}(\mathfrak{A})$, so the cohomology objects $H^i(\mathbf{A})$ of any $\mathbf{A} \in \mathbf{D}(\mathfrak{A})$ are well-defined.
 3. Viewing an object $A \in \mathfrak{A}$ as a complex concentrated in degree 0 yields an equivalence between \mathfrak{A} and the full subcategory of $\mathbf{D}(\mathfrak{A})$ whose objects are all complexes \mathbf{A} with $H^i(\mathbf{A}) = 0$ for all $i \neq 0$.

To describe the morphisms, we go back to Example 1.15. Recall that elements of the ring of fractions $S^{-1}R$ can be given by a tuple (s, r) (conventionally written r/s) for $s \in S$, $r \in R$. Likewise, morphisms $\mathbf{A} \rightarrow \mathbf{B}$ in the localised category $\mathbf{D}(\mathfrak{A})$ are given by diagrams in $\mathbf{K}(\mathfrak{A})$ of the form

$$\begin{array}{ccc} & \mathbf{C} & \\ \swarrow \scriptstyle \text{qis} & & \searrow \\ \mathbf{A} & & \mathbf{B} \end{array}$$

where the first map is a quasi-isomorphism.

The derived categories $\mathbf{D}^+(\mathfrak{A})$, $\mathbf{D}^-(\mathfrak{A})$, and $\mathbf{D}^b(\mathfrak{A})$ are defined analogously as localisations of $\mathbf{K}^+(\mathfrak{A})$, $\mathbf{K}^-(\mathfrak{A})$, and $\mathbf{K}^b(\mathfrak{A})$ respectively.

1.3.3 Triangulated categories

Both the homotopy category and the derived category of \mathfrak{A} are additive, but neither is usually abelian since (co)kernels are no longer guaranteed to be well-defined. Instead, we describe a structure on the homotopy and derived categories that ‘remembers’ short exact sequences.

Define the *mapping cone* of $f : \mathbf{A} \rightarrow \mathbf{B}$ to be the complex $\text{Cone}f$ given in degree i by $A^{i+1} \oplus B^i$, with differential

$$\begin{aligned} \dots \rightarrow A^{i+1} \oplus B^i &\longrightarrow A^{i+2} \oplus B^{i+1} \rightarrow \dots \\ (a, b) &\longmapsto (-da, db + fa). \end{aligned}$$

The complex $\mathcal{C}one f$ has natural injections from and projections to \mathbf{B} and $\mathbf{A}[1]$ respectively. This gives us a triple of morphisms

$$(\mathbf{A} \rightarrow \mathbf{B}, \mathbf{B} \rightarrow \mathcal{C}one f, \mathcal{C}one f \rightarrow \mathbf{A}[1])$$

in $\mathbf{K}(\mathfrak{A})$, which we call a *strict triangle*.

Definition 1.17. We say a triple of morphisms (u, v, w) in $\mathbf{K}(\mathfrak{A})$ is a *distinguished* (or *exact*) triangle if it is isomorphic to a strict triangle, i.e. there is a strict triangle (f, g, h) and a commuting diagram in $\mathbf{K}(\mathfrak{A})$

$$\begin{array}{ccccccc} \mathbf{A} & \xrightarrow{f} & \mathbf{B} & \xrightarrow{g} & \mathcal{C}one f & \xrightarrow{h} & \mathbf{A}[1] \\ \downarrow \alpha & & \downarrow & & \downarrow & & \downarrow \alpha[1] \\ \mathbf{E} & \xrightarrow{u} & \mathbf{F} & \xrightarrow{v} & \mathbf{G} & \xrightarrow{w} & \mathbf{E}[1] \end{array}$$

where the vertical maps are isomorphisms.

Strict triangles and distinguished triangles in $\mathbf{D}(\mathfrak{A})$ are analogously defined.

Remark 1.18. Here is how exact triangles in the derived category correspond to short exact sequences. Whenever

$$0 \rightarrow \mathbf{A} \xrightarrow{f} \mathbf{B} \rightarrow \mathbf{C} \rightarrow 0$$

is a short exact sequence in $\mathbf{C}(\mathfrak{A})$, the category $\mathbf{K}(\mathfrak{A})$ (and hence also $\mathbf{D}(\mathfrak{A})$) has a distinguished triangle

$$(\mathbf{A} \rightarrow \mathcal{C}yl f, \mathcal{C}yl f \rightarrow \mathcal{C}one f, \mathcal{C}one f \rightarrow \mathbf{A}[1])$$

where the complex $\mathcal{C}yl f$ (called the *mapping cylinder* of f), is defined as the mapping cone of $\mathcal{C}one f[-1] \rightarrow \mathbf{A}$. It can be shown that there are quasi-isomorphisms $\mathcal{C}yl f \rightarrow \mathbf{B}$ and $\mathcal{C}one f \rightarrow \mathbf{C}$.

The structure given to $\mathbf{K}(\mathfrak{A})$ by exact triangles is abstracted by the notion of a *triangulated category*, which is an additive category \mathfrak{T} with an automorphism $T : \mathfrak{T} \rightarrow \mathfrak{T}$ (called the *translation*) and a collection of triples of morphisms $(A \rightarrow B, B \rightarrow C, C \rightarrow TA)$ (called *distinguished triangles*) subject to four axioms (see Definition 10.2.1 of Weibel (2003)). The derived category of an abelian category satisfies these axioms, with the role of T played by $[1]$, and the distinguished triangles being the exact triangles.

Remark 1.19. The first axiom satisfied by a triangulated category suggests that every morphism fits into an exact triangle. In the derived category, this exact sequence is the one associated to the cone of the morphism. The other axioms put more conditions on the class of exact triangles— in particular, it is closed under rotations (i.e. if (u, v, w) is an exact triangle then so is $(v, w, u[1])$), and $(A \xrightarrow{\text{id}} A, A \rightarrow 0, 0 \rightarrow A[1])$ is an exact triangle for every A .

A *morphism of triangulated categories* (or *exact functor*) is an additive functor that commutes with translation and sends distinguished triangles to distinguished triangles.

We say \mathfrak{T}' is a *triangulated subcategory* of \mathfrak{T} if it is a full subcategory such that the inclusion functor is a morphism of triangulated categories, and every exact triangle in \mathfrak{T} is also exact in \mathfrak{T}' . Thus to show a full subcategory $\mathfrak{T}' \subset \mathfrak{T}$ is a triangulated subcategory, it suffices to check that \mathfrak{T}' is closed under translation, and whenever $(A \rightarrow B, B \rightarrow C, C \rightarrow TA)$ is an exact triangle such that $A, B \in \mathfrak{T}'$, we have an object in \mathfrak{T}' isomorphic to C .

Remark 1.20. The localisation functor q defined in Definition 1.14 is not necessarily additive, however Weibel (2003) describes conditions on the class S which ensure $q : \mathcal{C} \rightarrow S^{-1}\mathcal{C}$ is well-behaved. For the purposes of this exposition, the following fact suffices: when \mathcal{A} is the abelian category of sheaves or modules, the class Q of quasi-isomorphisms in a (triangulated subcategory of) $\mathbf{K}(\mathcal{A})$ is sufficiently nice that the localisation functor q and the functors induced by the universal property are morphisms of triangulated categories.

1.3.4 Generators of a triangulated category

Triangulated categories have two ‘fundamental operations’ built in– translations and taking mapping cones (i.e. completing morphisms to exact triangles). Then we say a collection of objects S in a triangulated category \mathcal{C} *generates it* if, up to isomorphism, every object of \mathcal{C} can be reached by taking these objects, shifting them, taking arbitrary morphisms between them, taking mapping cones of these morphisms, and repeating these operations finitely many times.

Definition 1.21. If \mathcal{T} is a triangulated category, we say a collection S of objects generates \mathcal{T} if the smallest triangulated subcategory of \mathcal{T} containing S is equivalent to \mathcal{T} .

Example 1.22. Let $\{E_1, \dots, E_i\}$ be a collection of objects in the abelian category \mathcal{A} . We define $\mathbf{K}^b\{E_1, \dots, E_i\}$ to be the full subcategory of $\mathbf{K}^b(\mathcal{A})$ whose objects are bounded complexes given in each degree by finite direct sums of the E_j . This category is closed under translation and taking mapping cones, so defines a triangulated subcategory of $\mathbf{K}^b(\mathcal{A})$. In this case, $\{E_1, \dots, E_i\}$ is a set of generators for $\mathbf{K}^b\{E_1, \dots, E_i\}$.

Generators give us an explicit handle on the triangulated category, as is illustrated by the following lemma from Beilinson (1978).

Lemma 1.23. Let $F : \mathcal{T} \rightarrow \mathcal{U}$ be a morphism of triangulated categories, and $\{X_i\}$ a collection of generators of \mathcal{T} such that the collection $\{FX_i\}$ generates \mathcal{U} . If for any pair X_i, X_j and any integer m the map

$$F : \text{Hom}_{\mathcal{T}}(X_i[m], X_j) \rightarrow \text{Hom}_{\mathcal{U}}(FX_i[m], FX_j)$$

is an isomorphism, then F is an equivalence of categories.

Proof. Note the image of F is a triangulated subcategory containing all the FX_i , i.e. the whole of \mathcal{U} . It suffices to show that the functor F is fully faithful, i.e. that $\text{Hom}(X, Y) \cong \text{Hom}(FX, FY)$ for all $X, Y \in \mathcal{C}$.

Let \mathcal{T}' be the full subcategory of those $X \in \mathcal{T}$ satisfying $\text{Hom}_{\mathcal{T}}(X[m], X_i) \cong \text{Hom}_{\mathcal{U}}(FX[m], FX_i)$ for all X_i and all m . We show that this is a triangulated subcategory of \mathcal{T} . Since it contains all the X_i by assumption, we have $\mathcal{T}' = \mathcal{T}$. Note \mathcal{T}' is clearly closed under taking translations. If $(X \rightarrow Y, Y \rightarrow Z, Z \rightarrow X)$ is an exact triangle in \mathcal{T} such that X and Y are in \mathcal{T}' , then for any generator X_i we have a commuting diagram

$$\begin{array}{ccccccccc} \text{Hom}(X, X_i) & \longrightarrow & \text{Hom}(Y, X_i) & \longrightarrow & \text{Hom}(Z, X_i) & \longrightarrow & \text{Hom}(X[1], X_i) & \longrightarrow & \text{Hom}(Y[1], X_i) \\ \downarrow \cong & & \downarrow \cong & & \downarrow & & \downarrow \cong & & \downarrow \cong \\ \text{Hom}(FX, FX_i) & \rightarrow & \text{Hom}(FY, FX_i) & \rightarrow & \text{Hom}(FZ, FX_i) & \rightarrow & \text{Hom}(FX[1], FX_i) & \rightarrow & \text{Hom}(FY[1], FX_i) \end{array}$$

in $\mathbf{D}(\mathbb{Z}\text{-Mod})$. By the derived version of the five lemma (see Weibel (2003) Exercise 10.2.2), the morphism in the middle is also an isomorphism. This shows \mathfrak{C}' is closed under taking mapping cones hence is a triangulated subcategory, as required.

A similar argument shows that if \mathfrak{C}'' is the full subcategory containing all objects $Y \in \mathfrak{C}$ satisfying $\text{Hom}_{\mathfrak{C}}(X, Y) \cong \text{Hom}_{\mathfrak{C}}(X, Y)$ for all $X \in \mathfrak{C}$, then \mathfrak{C}'' is a triangulated subcategory and contains all the X_i . Thus $\mathfrak{C}'' = \mathfrak{C}$, and $\text{Hom}(X, Y) \cong \text{Hom}(FX, FY)$ for all $X, Y \in \mathfrak{C}$ as required. \square

1.3.5 Functors between derived categories

Morphisms between homotopy and derived categories will typically come from additive functors $F : \mathbf{C}(\mathfrak{A}) \rightarrow \mathbf{C}(\mathfrak{B})$ on the corresponding chain complex categories.

Lemma 1.24. An additive functor $F : \mathbf{C}(\mathfrak{A}) \rightarrow \mathbf{C}(\mathfrak{B})$ descends to a morphism of triangulated categories $F : \mathbf{K}(\mathfrak{A}) \rightarrow \mathbf{K}(\mathfrak{B})$ if it takes cones to cones, i.e. for any morphism f in $\mathbf{C}(\mathfrak{A})$ we have $F(\mathcal{C}one f) = \mathcal{C}one F(f)$.

Proof. It is immediate from the splitting lemma that a chain map $f : \mathbf{A} \rightarrow \mathbf{B}$ is nullhomotopic if and only if the short exact sequence

$$0 \rightarrow \mathbf{B} \rightarrow \mathcal{C}one f \rightarrow \mathbf{A}[1] \rightarrow 0$$

is split. Since F maps cones to cones, the sequence above is mapped by F to

$$0 \rightarrow F(\mathbf{B}) \rightarrow \mathcal{C}one F(f) \rightarrow F(\mathbf{A})[1] \rightarrow 0.$$

But additive functors send split exact sequences to split exact sequences, so $F(f)$ is nullhomotopic and F descends to an additive functor $\mathbf{K}(\mathfrak{A}) \rightarrow \mathbf{K}(\mathfrak{B})$. Moreover, the image of a strict triangle is a strict triangle so we have a morphism of triangulated categories. \square

In order to induce a functor $\mathbf{D}(\mathfrak{A}) \rightarrow \mathbf{D}(\mathfrak{B})$, the functor F must send quasi-isomorphisms to quasi-isomorphisms. Often, however, this is not the case and we resort to one of two options– restrict to a triangulated subcategory (for example the bounded derived category) so that F does map quasi-isomorphisms to isomorphisms, or define functors on the derived category that preserve *some* of the properties of F . We describe both the approaches.

Descent to smaller triangulated subcategories. Given a morphism $F : \mathbf{K}(\mathfrak{A}) \rightarrow \mathbf{K}(\mathfrak{B})$ of triangulated categories, we say a complex $\mathbf{A} \in \mathbf{K}(\mathfrak{A})$ is *F-acyclic* if the complex $F(\mathbf{A})$ is acyclic.

Proposition 1.25. Suppose \mathbf{K} is a triangulated subcategory of $\mathbf{K}(\mathfrak{A})$ such that every acyclic complex in \mathbf{K} is also *F-acyclic*. Then the restricted functor $F : \mathbf{K} \rightarrow \mathbf{K}(\mathfrak{B})$ descends to a morphism of triangulated categories $F : \mathbf{D} \rightarrow \mathbf{D}(\mathfrak{B})$ where \mathbf{D} is the derived category of \mathbf{K} .

Proof. If $f : \mathbf{A} \rightarrow \mathbf{B}$ is a quasi-isomorphism in \mathbf{K} , then examining the long exact sequence in cohomology associated to $0 \rightarrow \mathbf{B} \rightarrow \mathcal{C}one f \rightarrow \mathbf{A}[1] \rightarrow 0$ shows that $\mathcal{C}one f$ is acyclic. Hence by assumption $F(\mathcal{C}one f)$ is acyclic. Now F preserves exact triangles, so the triangle

$$(F(\mathbf{A}) \rightarrow F(\mathbf{B}), F(\mathbf{B}) \rightarrow F(\mathcal{C}one f), F(\mathcal{C}one f) \rightarrow F(\mathbf{A})[1])$$

is exact and gives a short exact sequence in \mathfrak{A} . Examining the associated long exact sequence on homology shows that $F(f) : F(\mathbf{A}) \rightarrow F(\mathbf{B})$ is a quasi-isomorphism. Thus the composite $\mathbf{K} \rightarrow \mathbf{K}(\mathfrak{B}) \rightarrow \mathbf{D}(\mathfrak{B})$

sends quasi-isomorphisms to isomorphisms. By the universal property of localisation we have an induced functor $\mathbf{D} \rightarrow \mathbf{D}(\mathfrak{A})$. By Remark 1.20, we are done. \square

This approach is used, for instance, to establish the Bernstein-Gel'fand-Gel'fand correspondence of Section 3 which gives an equivalence of bounded derived categories but fails to extend to the unbounded situation directly.

Derived functors. Let \mathbf{K} be a triangulated subcategory of $\mathbf{K}(\mathfrak{A})$, and $q : \mathbf{K} \rightarrow \mathbf{D}$ the localisation map with respect to the class of quasi-isomorphisms. For $F : \mathbf{K} \rightarrow \mathbf{K}(\mathfrak{A})$ a morphism of triangulated categories, we define associated *derived functors* by their universal properties.

Definition 1.26. A (total) right derived functor of F on \mathbf{K} is a morphism of triangulated categories $\mathbf{R}F : \mathbf{D} \rightarrow \mathbf{D}(\mathfrak{B})$ with a natural transformation

$$\begin{array}{ccccc} \mathbf{K} & \xrightarrow{F} & \mathbf{K}(\mathfrak{B}) & \xrightarrow{q} & \mathbf{D}(\mathfrak{B}) \\ & & \Downarrow \zeta & & \\ \mathbf{K} & \xrightarrow{q} & \mathbf{D} & \xrightarrow{\mathbf{R}F} & \mathbf{D}(\mathfrak{B}) \end{array}$$

such that if $G : \mathbf{D} \rightarrow \mathbf{D}(\mathfrak{B})$ is a morphism equipped with a natural transformation $\zeta' : qF \Rightarrow Gq$, then there is a unique natural transformation $\eta : \mathbf{R}F \Rightarrow G$ such that $\zeta'_A = \eta_{qA} \circ \zeta_A$ for every $A \in \mathbf{D}$.

The (total) left derived functor of F on \mathbf{K} is a morphism $\mathbf{L}F : \mathbf{D} \rightarrow \mathbf{D}(\mathfrak{B})$ with a natural transformation $\zeta : (\mathbf{L}F)q \Rightarrow qF$ satisfying a universal property similar to the right derived functor.

We provide a brief account of various right and left derived functors that come up in this exposition.

Example 1.27. If $F : \mathfrak{A} \rightarrow \mathfrak{B}$ is an exact functor, then the induced functor $F : \mathbf{C}(\mathfrak{A}) \rightarrow \mathbf{C}(\mathfrak{B})$ preserves quasi-isomorphisms (since f is a quasi-isomorphism if and only if $\mathcal{C}one f$ is an acyclic complex) and we have a functor $F : \mathbf{D}(\mathfrak{A}) \rightarrow \mathbf{D}(\mathfrak{B})$. In effect, F is its own right and left derived functor on $\mathbf{K}(\mathfrak{A})$.

Proposition 1.25 can be seen as a generalisation of above, saying that if a triangulated subcategory \mathbf{K} is such that every F -acyclic complex is F -acyclic then F is its own right and left derived functor on \mathbf{K} . Often, we can choose the subcategory \mathbf{K} (with localisation \mathbf{D}) such that $\mathbf{D} \cong \mathbf{K}(\mathfrak{A})$. This allows us to show existence of derived functors in certain special cases.

Example 1.28. For a noetherian scheme \mathbf{X} , let $\mathbf{K}_{\text{inj}}^+(\mathcal{Q}coh \mathbf{X})$ be the full subcategory of $\mathbf{K}(\mathcal{Q}coh \mathbf{X})$ containing complexes of injective sheaves that are bounded below. Since injective sheaves are flasque, every acyclic complex in $\mathbf{K}_{\text{inj}}^+(\mathcal{Q}coh \mathbf{X})$ is Γ -acyclic where Γ is the global sections functor.

Now every quasi-isomorphism in $\mathbf{K}_{\text{inj}}^+$ is an isomorphism and moreover, every complex in $\mathbf{K}^+(\mathcal{Q}coh \mathbf{X})$ is quasi-isomorphic to a bounded below complex of injectives. This can be used to show that the derived category $\mathbf{D}^+(\mathcal{Q}coh \mathbf{X})$ is equivalent to $\mathbf{K}_{\text{inj}}^+(\mathcal{Q}coh \mathbf{X})$. Thus Γ has left and right derived functors $\mathbf{D}^+(\mathcal{Q}coh \mathbf{X}) \rightarrow \mathbf{D}(\mathbb{Z}\text{-Mod})$. In practice, computing these on a complex \mathcal{E} involves first finding a quasi-isomorphic complex of injectives and then applying Γ . In particular, if \mathcal{E} is a complex concentrated in degree 0, then the i th sheaf cohomology is $H^i(\mathbf{R}\Gamma(\mathcal{E}))$.

In fact, Weibel (2003) shows that the right derived functor extends to the whole derived category, giving a morphism of triangulated categories $\mathbf{R}\Gamma : \mathbf{D}(\mathcal{Q}coh \mathbf{X}) \rightarrow \mathbf{D}(\mathbb{Z}\text{-Mod})$.

Example 1.29. If \mathfrak{A} has enough injectives (i.e. any object of \mathfrak{A} can be resolved by injectives), then for each complex $\mathbf{A} \in \mathbf{K}(\mathfrak{A})$ the functor $\text{Tot}(\text{Hom}(\mathbf{A}, -))$ has left and right derived functors $\mathbf{D}^+(\mathfrak{A}) \rightarrow \mathbf{D}(\mathbb{Z}\text{-Mod})$. Then Weibel (2003) shows that the right derived functor is a bifunctor

$$\mathbf{R}\text{Hom} : \mathbf{D}(\mathfrak{A})^{\text{op}} \times \mathbf{D}^+(\mathfrak{A}) \rightarrow \mathbf{D}(\mathbb{Z}\text{-Mod}).$$

Moreover, if \mathbf{A} and \mathbf{B} are both complexes bounded below, then we have the *hyperext* groups

$$\text{Ext}^i(\mathbf{A}, \mathbf{B}) \cong H^i(\mathbf{R}\text{Hom}(\mathbf{A}, \mathbf{B})) = \text{Hom}_{\mathbf{D}(\mathbf{A})}(\mathbf{A}[-i], \mathbf{B}).$$

Likewise, if $\mathfrak{A} = \mathbf{R}\text{-Mod}$ has enough projectives, then the total tensor product functor has a left derived functor

$$\otimes_{\mathbf{R}}^{\mathbf{L}} : \mathbf{D}^-(\mathfrak{A}) \times \mathbf{D}^-(\mathfrak{A}) \rightarrow \mathbf{D}(\mathbb{Z}\text{-Mod}).$$

The cohomologies of this are the *hypertor* functors

$$\text{Tor}_i^{\mathbf{R}}(\mathbf{A}, \mathbf{B}) = H^{-i}(\mathbf{A} \otimes_{\mathbf{R}}^{\mathbf{L}} \mathbf{B}),$$

which are computed as usual using projective resolutions.

If \mathbf{X} is a noetherian scheme, the derived tensor product can be constructed if one uses locally free sheaves instead of projectives. Thus we have a bifunctor

$$(- \otimes^{\mathbf{L}} -) : \mathbf{D}^b(\mathcal{Qcoh} \mathbf{X}) \times \mathbf{D}^b(\mathcal{Qcoh} \mathbf{X}) \rightarrow \mathbf{D}^b(\mathcal{Qcoh} \mathbf{X}).$$

Weibel (2003) also shows that if $f : \mathbf{X} \rightarrow \mathbf{Y}$ is a proper morphism of projective schemes, then the derived functors $\mathbf{R}f^*$ and $\mathbf{L}f_*$ exist on bounded derived categories of quasi-coherent sheaves.

2 Coherent sheaves on \mathbb{P}^n

For a noetherian scheme \mathbf{X} , we write $\mathbf{D}^b(\mathbf{X})$ to mean $\mathbf{D}^b(\mathcal{Coh} \mathbf{X})$.

The goal of this section is to prove Beilinson's theorem, which asserts that $\{\mathcal{O}, \mathcal{O}(-1), \dots, \mathcal{O}(-n)\}$ and $\{\mathcal{O}, \Omega(1), \dots, \Omega^n(n)\}$ are both generating sets for the derived category $\mathbf{D}^b(\mathbb{P}^n)$. We provide Beilinson's (1978) original proof, following the treatment in Caldararu (2005) and Carbone (2016). There are two key ideas involved– the first is that the identity functor on $\mathbf{D}^b(\mathbb{P}^n)$ admits a factorisation

$$\begin{array}{ccc} \mathbf{D}^b(\mathbb{P}^n \times \mathbb{P}^n) & \xrightarrow{-\otimes^{\mathbf{L}} \mathcal{O}_{\Delta}} & \mathbf{D}^b(\mathbb{P}^n \times \mathbb{P}^n) \\ \pi_2^* \uparrow & & \downarrow \mathbf{R}\pi_{1*} \\ \mathbf{D}^b(\mathbb{P}^n) & \xrightarrow{\text{id}} & \mathbf{D}^b(\mathbb{P}^n) \end{array}$$

where $\pi_1, \pi_2 : \mathbb{P}^n \times \mathbb{P}^n \rightarrow \mathbb{P}^n$ are the projection maps, and $\mathcal{O}_{\Delta} \in \mathcal{Coh}(\mathbb{P}^n \times \mathbb{P}^n)$ is the structure sheaf of the diagonal subscheme. This follows from the geometric theory of a *Fourier-Mukai transform* associated to a pair of schemes \mathcal{X}, \mathcal{Y} , and we briefly sketch the construction in Section 2.1.

The second observation, called *Beilinson's resolution of the diagonal*, follows from the algebraic theory of *Koszul resolutions* and shows that \mathcal{O}_{Δ} admits a resolution by locally free sheaves of the form $\pi_1^*(\Omega^i(i)) \otimes \pi_2^*(\mathcal{O}(-i))$, where Ω is the sheaf of differentials on \mathbb{P}^n . Combined with the factorisation of identity, this provides an algorithm to resolve any coherent sheaf on \mathbb{P}^n in terms of the $\mathcal{O}(i)$ thus proving Beilinson's result.

2.1 Fourier-Mukai transforms

The material in this section is from Huybrechts (2006), where the topic occupies a central position. Given two smooth projective k -schemes \mathbf{X}_1 and \mathbf{X}_2 , we associate to each object $\mathcal{E} \in \mathbf{D}^b(\mathbf{X}_1 \times \mathbf{X}_2)$ a morphism of triangulated categories $\Phi_{\mathcal{E}} : \mathbf{D}^b(\mathbf{X}_1) \rightarrow \mathbf{D}^b(\mathbf{X}_2)$.

Definition 2.1. The *Fourier-Mukai transform with kernel \mathcal{E}* of a complex $\mathcal{A} \in \mathbf{D}^b(\mathbf{X}_1)$ is defined as

$$\Phi_{\mathcal{E}}(\mathcal{A}) = \mathbf{R}\pi_{1*}(\pi_2^* \mathcal{A} \otimes^{\mathbf{L}} \mathcal{E}) \in \mathbf{D}^b(\mathbf{X}_2).$$

Here $\pi_i : \mathbf{X}_1 \times \mathbf{X}_2 \rightarrow \mathbf{X}_i$ ($i = 1, 2$) be the projection maps, these are flat so the pullback functors π_i^* are exact and need no derivation by Example 1.27. Being the composition of three exact functors, the Fourier-Mukai transform $\Phi_{\mathcal{E}}$ is an exact functor. Moreover, the dependence on the kernel is functorial— for a fixed $\mathcal{A} \in \mathbf{D}^b(\mathbf{X}_1)$, the map

$$\begin{aligned} \Phi_{-}(\mathcal{A}) : \mathbf{D}^b(\mathbf{X}_1 \times \mathbf{X}_2) &\longrightarrow \mathbf{D}^b(\mathbf{X}_2) \\ \mathcal{E} &\longmapsto \Phi_{\mathcal{E}}(\mathcal{A}) \end{aligned}$$

is the composite $\pi_{1*}(\pi_2^* \mathcal{A} \otimes^{\mathbf{L}} -)$, hence is an exact functor.

Remark 2.2. The name comes from the following analogy with functional analysis— given a finite-dimensional vector space X and its dual Ξ , to any smooth function $E(x, \xi) : X \times \Xi \rightarrow \mathbb{C}$ we can associate a linear map $\phi_E : L^2(X) \rightarrow L^2(Y)$ between the spaces of square-integrable functions, given by $f \mapsto \int_X f(x) E(x, \xi) dx$. If $E(x, \xi) = e^{2\pi i \langle x, \xi \rangle}$, then ϕ_E is an isomorphism called the *Fourier transform*.

The Fourier-Mukai transform yields interesting functors based on choice of \mathcal{E} . The following result is useful in order to study these.

Lemma 2.3 (Projection formula). If $f : \mathbf{X}_1 \rightarrow \mathbf{X}_2$ is a proper morphism of projective schemes over k , then for any complexes $\mathcal{E} \in \mathbf{D}^b(\mathbf{X}_2)$ and $\mathcal{F} \in \mathbf{D}^b(\mathbf{X}_1)$, we have a natural isomorphism

$$\mathbf{R}f_*(\mathcal{F} \otimes^{\mathbf{L}} \mathbf{L}f^*(\mathcal{E})) \cong \mathbf{R}f_*(\mathcal{F}) \otimes^{\mathbf{L}} \mathcal{E}.$$

Proof. See Section 3.3 of Huybrechts (2006). □

Example 2.4. If $\mathbf{X}_1 = \mathbf{X}_2 = \mathbf{X}$ and $\mathbf{X} \xrightarrow{\iota} \mathbf{X} \times \mathbf{X}$ is the diagonal inclusion, then we can consider the Fourier-Mukai transform with kernel $\mathcal{O}_{\Delta} = \iota_* \mathcal{O}_{\mathbf{X}}$, the structure sheaf of the diagonal subscheme. Since ι is a closed immersion, the pushforward ι_* is exact and $\mathbf{R}\iota_* = \iota_*$ as derived functors. Hence $\mathcal{O}_{\Delta} = \mathbf{R}\iota_* \mathcal{O}_{\mathbf{X}}$ in $\mathbf{D}^b(\mathbf{X})$, and we can use the projection formula to get

$$\begin{aligned} \Phi_{\mathcal{O}_{\Delta}}(\mathcal{A}) &= \mathbf{R}\pi_{1*}(\pi_2^* \mathcal{A} \otimes^{\mathbf{L}} \mathbf{R}\iota_* \mathcal{O}_{\mathbf{X}}) \\ &= \mathbf{R}\pi_{1*} \circ \mathbf{R}\iota_*(\mathbf{L}\iota^* \pi_2^* \mathcal{A} \otimes^{\mathbf{L}} \mathcal{O}_{\mathbf{X}}) \\ &= \mathbf{R}(\pi_1 \circ \iota)_*(\mathbf{L}(\pi_2 \circ \iota)^* \mathcal{A} \otimes^{\mathbf{L}} \mathcal{O}_{\mathbf{X}}) \\ &= \mathcal{A} \otimes^{\mathbf{L}} \mathcal{O}_{\mathbf{X}}. \end{aligned}$$

But $\mathcal{O}_{\mathbf{X}}$ is a locally free sheaf so the functor $(- \otimes^{\mathbf{L}} \mathcal{O}_{\mathbf{X}})$ is the same as $(- \otimes \mathcal{O}_{\mathbf{X}})$, which is identity. In other words, the Fourier-Mukai transform with kernel \mathcal{O}_{Δ} is the identity functor.

Replacing the trivial bundle $\mathcal{O}_{\mathbf{X}}$ in the above computation with some other line bundle \mathcal{L} on \mathbf{X} , we see that the derived functor $(- \otimes \mathcal{L})$ is the Fourier-Mukai transform with kernel $\iota_* \mathcal{L}$. Similarly,

one can show that the Fourier-Mukai kernel $\mathcal{O}_\Delta[1]$ yields the shift functor $\mathcal{A} \mapsto \mathcal{A}[1]$. Thus Fourier-Mukai transforms generalise many familiar constructions. It is in fact a theorem of Orlov that any fully faithful exact functor $\mathbf{D}^b(\mathbf{X}_1) \rightarrow \mathbf{D}^b(\mathbf{X}_2)$ that admits adjoints must arise as the Fourier-Mukai transform for some kernel determined uniquely up to isomorphism.

2.2 Koszul resolutions

Given a ring R and a sequence (r_0, \dots, r_n) of elements in R , the associated *Koszul complex* is a very useful construction which detects various homological properties of the ring, and often yields free resolutions of the R -module $R/(r_0, \dots, r_n)$. The construction and theory of Koszul complexes is treated in its full generality in Eisenbud (1995); here we only study the behaviour in two special cases which we use to resolve the diagonal– the first is when (r_0, \dots, r_n) generate the unit ideal, and the second is when they form a regular sequence.

Definition 2.5. Given a ring R and a sequence (r_0, \dots, r_n) of elements in R , the associated *Koszul complex* is the complex of R -modules given by

$$K_R(r_0, \dots, r_n) : \quad 0 \rightarrow \bigwedge_R^{n+1}(R^{n+1}) \rightarrow \bigwedge_R^n(R^{n+1}) \rightarrow \dots \rightarrow \bigwedge_R^2(R^{n+1}) \rightarrow R^{n+1} \rightarrow R \rightarrow 0$$

$$d(e_{\alpha_1} \wedge \dots \wedge e_{\alpha_i}) = \sum_j (-1)^{i+j+1} r_{\alpha_j} \cdot (e_{\alpha_1} \wedge \dots \hat{e}_{\alpha_j} \dots \wedge e_{\alpha_i})$$

where e_0, \dots, e_n are the standard generators of R^{n+1} , and $\hat{}$ denotes omission of a term. We put the term $\bigwedge_R^i(R^{n+1})$ in differential degree $-i$.

Observe that the modules appearing in $K_R(r_0, \dots, r_n)$ are free, so acyclic Koszul complexes yield free R -resolutions. In the simplest case when the sequence contains a single element r_0 , the Koszul complex is given by

$$K(r_0) : \quad 0 \rightarrow R \xrightarrow{r_0} R \rightarrow 0,$$

so it is exact if and only if r_0 is a unit in R . This result generalises to sequences (r_0, \dots, r_n) that generate the unit ideal.

Proposition 2.6. If R is a ring and $(r_0, \dots, r_n) = A$, then the Koszul complex $K_R(r_0, \dots, r_n)$ is exact.

Proof. We show that the identity on $K_R(r_0, \dots, r_n)$ is chain homotopic to the zero morphism. By assumption, there are elements $\lambda_0, \dots, \lambda_n \in R$ such that $\sum_i \lambda_i r_i = -1$. Then consider the map given by

$$h : \bigwedge_R^i(R^{n+1}) \rightarrow \bigwedge_R^{i+1}(R^{n+1})$$

$$h(e) = \sum_j \lambda_j e \wedge e_j$$

A straightforward basis-wise check shows $d \circ h + h \circ d = \text{id}$, showing h is the required chain homotopy. Thus the complex $K_R(r_0, \dots, r_n)$ is contractible, hence exact. \square

Looking again at the Koszul complex for a single element $r_0 \in R$, we have that $H^1(K(r_0)) = 0$ if and only if r_0 is not a zero-divisor in R – in this case the complex is a free resolution of $R/(r_0)$. Recall that (r_0, \dots, r_n) is an *R -regular sequence* if r_0 is not a zero-divisor in R , and for every $0 \leq i < n$, r_{i+1} is not a zero-divisor for the module $R/(r_0, \dots, r_i)$. Then Eisenbud (1995) proves that whenever (r_0, \dots, r_n) is a regular sequence, the associated Koszul complex is exact everywhere except in

degree 0 where it has cohomology $R/(r_0, \dots, r_n)$. We prove a special case of the result, r_0, \dots, r_n are the indeterminates in a polynomial algebra.

Proposition 2.7 (Loday (2012)). Suppose R contains a field of characteristic 0. Then for the polynomial ring $S = R[x_0, \dots, x_n]$, we have

$$H^i(K_S(x_0, \dots, x_n)) = \begin{cases} R, & i = 0 \\ 0, & \text{otherwise} \end{cases}.$$

Proof. Write M for the rank $n+1$ free R -module with generators m_0, \dots, m_n . Then we can identify $\bigwedge_S^\bullet(S^{n+1})$ with the algebra $\bigwedge_R^\bullet(M) \otimes_k S$, these are graded so that M lies in degree 1. Considering S as an R -algebra graded by degree, we see that $K = K_S(x_0, \dots, x_n)$ is a complex of graded R -algebras given by

$$\begin{aligned} \mathbf{K}: \quad 0 \rightarrow \bigwedge_R^{n+1}(M) \otimes_R S\langle -n-1 \rangle \rightarrow \bigwedge_R^n(M) \otimes_R S\langle -n \rangle \rightarrow \dots \rightarrow S \rightarrow 0 \\ d((m_{\alpha_1} \wedge \dots \wedge m_{\alpha_i}) \otimes s) = \sum_j (-1)^{i+j+1} (m_{\alpha_1} \wedge \dots \wedge \hat{m}_{\alpha_j} \wedge \dots \wedge m_{\alpha_i}) \otimes s x_{\alpha_j}. \end{aligned}$$

Note the differential d preserves internal grading, so we can write the complex above as a direct sum $\mathbf{K} = \bigoplus_r \mathbf{K}_r$ where \mathbf{K}_r is the complex of R -modules formed at Adam's degree r (called the r th strand of \mathbf{K}). Since cohomology is an additive functor, we have $H^i(\mathbf{K}) = \bigoplus_r H^i(\mathbf{K}_r)$.

Now the strands in negative degrees vanish everywhere, and the \mathbf{K}_0 has the module R concentrated in differential degree 0. Thus it suffices to prove every other strand is exact. We do this by showing that the identity map on \mathbf{K}_r is nullhomotopic whenever $r > 0$. In this case, we know by assumption that $r \in R$ is a unit so consider the map

$$\begin{aligned} h: \bigwedge^i(M) \otimes_k S_{r-i} \rightarrow \bigwedge^{i+1}(M) \otimes_k S_{r-i-1} \\ h(m \otimes (x_{\beta_1} \dots x_{\beta_{r-i}})) = -\frac{1}{r} \sum_j (m \wedge x_{\beta_j}) \otimes (x_{\beta_1} \dots \hat{x}_{\beta_j} \dots x_{\beta_{r-i}}). \end{aligned}$$

A straightforward basis-wise check shows $h \circ d + d \circ h = \text{id}$ as required. \square

Koszul complexes in geometry. We use Koszul complexes in the following geometric setting— on the affine scheme $\mathbf{X} = \text{Spec } R$, an $(n+1)$ -tuple $s = (r_0, \dots, r_n)$ in A can be seen as a global section of the free sheaf $\mathcal{E} = \mathcal{O}_{\mathbf{X}}^{\oplus(n+1)}$. Then the Koszul complex associated to (r_0, \dots, r_n) yields a complex of coherent sheaves, given by

$$\mathcal{K}_{\mathbf{X}}(s): \quad 0 \rightarrow \bigwedge^{n+1} \mathcal{E}^\vee \rightarrow \bigwedge^n \mathcal{E}^\vee \rightarrow \dots \rightarrow \mathcal{E}^\vee \rightarrow \mathcal{O}_{\mathbf{X}} \rightarrow \mathcal{O}_{\mathbb{V}(s)} \rightarrow 0$$

where $\mathbb{V}(s)$ is the zero locus of s , i.e. the closed subscheme corresponding to the ideal (r_0, \dots, r_n) . The differential then can be interpreted as ‘contracting’ an i -form in $\bigwedge^i \mathcal{E}^\vee$ with the section s . If (r_0, \dots, r_n) is a regular sequence then we have shown that the complex $\mathcal{K}_{\mathbf{X}}(s)$ is exact, giving a locally free resolution of $\mathcal{O}_{\mathbb{V}(s)}$.

The construction automatically extends to arbitrary schemes \mathbf{X} — given global section of a locally free sheaf $\mathcal{E} \in \text{Coh } \mathbf{X}$, we can cover \mathbf{X} by affine open subschemes $\mathbf{X} = \bigcup_\alpha U_\alpha$; then the associated Koszul complexes $\mathcal{K}_{U_\alpha}(s|_{U_\alpha})$ glue to give a Koszul complex $\mathcal{K}_{\mathbf{X}}(s)$ of coherent sheaves on \mathbf{X} .

Remark 2.8. The exactness of this complex can be checked stalk-locally– given $p \in \mathbf{X}$, the section s gives a tuple (r_0, \dots, r_n) in the local ring $\mathcal{O}_{\mathbf{X},p}$. Localising the complex $\mathcal{K}_{\mathbf{X}}(s)$ at p then yields the Koszul complex $K_{\mathcal{O}_{\mathbf{X},p}}(r_0, \dots, r_n)$. In particular, if $p \notin \mathbb{V}(s)$ then (r_0, \dots, r_n) generate the unit ideal so the localised complex is exact by Proposition 2.6. Thus it suffices to check that the complex $\mathcal{K}_{\mathbf{X}}(s)$ is exact at points of $\mathbb{V}(s)$.

Example 2.9 (Generalised Euler exact sequences). Consider the locally free sheaf $\mathcal{E} = \mathcal{O}(1)^{\oplus n+1}$ on \mathbb{P}^n , and the global section $s = (x_0, \dots, x_n)$. Since $\mathbb{V}(s) = \emptyset$, we have the exact Koszul complex

$$0 \rightarrow \bigwedge^{n+1} \mathcal{E}^\vee \rightarrow \bigwedge^n \mathcal{E}^\vee \rightarrow \dots \rightarrow \mathcal{E}^\vee \rightarrow \mathcal{O} \rightarrow 0.$$

On the standard affine neighbourhood \mathbf{U}_0 , we have $\mathcal{E}^\vee(\mathbf{U}_0) = M \oplus \Omega(\mathbf{U}_0)$ where $\Omega(\mathbf{U}_0)$ is the submodule of those elements that vanish when contracted with s . Thus M has rank 1 (with generator $\frac{1}{x_0}$) and we have

$$\bigwedge^i \mathcal{E}^\vee(\mathbf{U}_0) = \Omega^i(\mathbf{U}_0) \oplus M \otimes_{\mathcal{O}(\mathbf{U}_0)} \Omega^{i-1}(\mathbf{U}_0).$$

The differential $\bigwedge^i \mathcal{E}^\vee(\mathbf{U}_0) \rightarrow \bigwedge^{i-1} \mathcal{E}^\vee(\mathbf{U}_0)$ in the Koszul complex is seen to be the composition

$$\bigwedge^i \mathcal{E}^\vee(\mathbf{U}_0) \twoheadrightarrow M \otimes_{\mathcal{O}(\mathbf{U}_0)} \Omega^{i-1}(\mathbf{U}_0) \cong \Omega^i(\mathbf{U}_0) \hookrightarrow \bigwedge^{i-1}(\mathbf{U}_0).$$

Hence $\ker(\bigwedge^i \mathcal{E}^\vee \rightarrow \bigwedge^{i-1} \mathcal{E}^\vee) = \Omega^i$ and the Koszul complex breaks into short exact sequences

$$0 \rightarrow \Omega^i \rightarrow \bigwedge^i \mathcal{E}^\vee \rightarrow \bigwedge^{i-1} \mathcal{E}^\vee \rightarrow 0. \quad (3)$$

In case $i = 0$, this can be seen to be the Euler exact sequence (2).

2.3 Beilinson's theorem

Let $\iota : \mathbb{P}^n \rightarrow \mathbb{P}^n \times \mathbb{P}^n$ be the inclusion of the diagonal subscheme Δ , and write $\pi_1, \pi_2 : \mathbb{P}^n \times \mathbb{P}^n \rightarrow \mathbb{P}^n$ for the two coordinate projections. To prove Beilinson's theorem, we will show that the Koszul complex associated to a particular locally free sheaf $\mathcal{E} \in \mathcal{Coh}(\mathbb{P}^n \times \mathbb{P}^n)$ gives a resolution of $\mathcal{O}_\Delta = \iota_* \mathcal{O}_{\mathbb{P}^n}$. The sheaf \mathcal{E} is chosen so that the sheaves arising in the resolution are of the form $\pi_1^* \mathcal{O}(-i) \otimes \pi_2^* \Omega^i(i)$. Given such a resolution, the functoriality (in kernel) of the Fourier-Mukai transform can be used to obtain a resolution of any $\mathcal{A} \in \mathbf{D}^b(\mathbb{P}^n)$.

In particular, the (-1) th twist of the second Euler exact sequence (2) yields an associated long exact sequence of cohomology

$$0 \rightarrow \Gamma(\mathcal{O}(-1)) \rightarrow \Gamma(\mathcal{O}^{\oplus(n+1)}) \rightarrow \Gamma(\mathcal{T}(-1)) \rightarrow H^1(\mathcal{O}(-1)) \rightarrow \dots$$

But all cohomologies of $\mathcal{O}(-1)$ vanish, so we have an isomorphism $\Gamma(\mathcal{T}(-1)) \cong \Gamma(\mathcal{O}^{\oplus(n+1)})$. Thus the global sections of $\mathcal{T}(-1)$ form an $n + 1$ -dimensional k -vector space. Write $\frac{\partial}{\partial x_0}, \dots, \frac{\partial}{\partial x_n}$ for the standard basis.

2.3.1 Resolution of the diagonal

Given two coherent sheaves \mathcal{A}, \mathcal{B} on \mathbb{P}^n , write $\mathcal{A} \boxtimes \mathcal{B}$ for the coherent sheaf $\pi_1^* \mathcal{A} \otimes \pi_2^* \mathcal{B}$ on $\mathbb{P}^n \times \mathbb{P}^n$. Using this notation, the locally free sheaf on $\mathbb{P}^n \times \mathbb{P}^n$ (and associated global section) which we use to obtain the Koszul resolution is given by

$$\mathcal{E} = \mathcal{O}(1) \boxtimes \mathcal{T}(-1), \quad s = \sum_{\alpha=0}^n x_\alpha \boxtimes \frac{\partial}{\partial y_\alpha} \in \Gamma(\mathcal{E}),$$

where $(x_0 : \dots : x_n)$ and $(y_0 : \dots : y_n)$ are homogeneous coordinates on the first and second copy of \mathbb{P}^n respectively. To understand how the section s is defined, note that global sections of a sheaf \mathcal{A} are same as morphisms from the structure sheaf to \mathcal{A} . Then by functoriality of pullbacks (and since pullback of the structure sheaf is again the structure sheaf), we lift $x_\alpha \in \Gamma(\mathcal{O}(1))$ to a global section $\pi_1^*(x_\alpha) \in \Gamma(\pi_1^*\mathcal{O}(1))$. The sections $\pi_2^*(\frac{\partial}{\partial y_\alpha}) \in \Gamma(\pi_2^*\mathcal{T}(-1))$ are likewise defined. Thus we have a global section $\pi_1^*(x_\alpha) \otimes \pi_2^*(\frac{\partial}{\partial y_\alpha})$ of the tensor product *presheaf*, which gets mapped to a global section $x_\alpha \boxtimes \frac{\partial}{\partial y_\alpha}$ of the tensor product sheaf by the canonical map

$$\Gamma(\pi_1^*\mathcal{O}(1)) \otimes_k \Gamma(\pi_2^*\mathcal{T}(-1)) \rightarrow \Gamma(\mathcal{O}(1) \boxtimes \mathcal{T}(-1)).$$

The following results justify our choice of \mathcal{E} and s .

Lemma 2.10. For \mathcal{E} and s as given, the section s vanishes precisely on the diagonal subscheme.

Proof. We check this on affine patches— writing $U_\alpha = \mathbb{P}^n \setminus \mathbb{V}(x_\alpha)$ and $V_\beta = \mathbb{P}^n \setminus \mathbb{V}(y_\beta)$, we see that the various open sets of the form $U_\alpha \times V_\beta$ are isomorphic to \mathbb{A}^2 and form an open cover of $\mathbb{P}^n \times \mathbb{P}^n$. We focus on the patch $U_0 \times V_0$, though the calculation on other patches is similar.

Write $x_{\alpha/0} = x_\alpha/x_0$ ($1 \leq \alpha \leq n$) for the standard coordinates on U_0 , the coordinates on V_0 are written similarly. Then writing $S = k[y_{1/0}, \dots, y_{n/0}]$ for the coordinate ring of V_0 , we can read off the module of sections of $\mathcal{T}(-1)$ on V_0 as the quotient

$$T = \frac{S^{n+1}}{S \cdot (1, y_{1/0}, \dots, y_{n/0})}.$$

Since the quotient map commutes with restriction of sections, the restriction of $\frac{\partial}{\partial y_\alpha} \in \Gamma(\mathcal{T}(-1))$ is the image in T of the α th standard generator of S^{n+1} . Then we must have

$$\frac{\partial}{\partial y_0} \Big|_{V_0} + y_{1/0} \frac{\partial}{\partial y_1} \Big|_{V_0} + \dots + y_{n/0} \frac{\partial}{\partial y_n} \Big|_{V_0} = 0.$$

Writing $S' = k[x_{1/0}, \dots, x_{n/0}]$ for the coordinate ring on U_0 , it is clear that $\mathcal{O}(1)$ restricts to the module S' . The restriction of the section $x_\alpha \in \Gamma(\mathcal{O}(1))$ is given by $\frac{x_\alpha}{x_0}$.

Write $R = k[x_{1/0}, \dots, x_{n/0}] \cong S' \otimes_k S$ for the structure sheaf of $U_0 \times V_0$. Since pulling back a sheaf on an affine schemes corresponds to taking tensor product with the coordinate ring, the restriction of $\mathcal{O}(1) \boxtimes \mathcal{T}(-1)$ to $U_0 \times V_0$ corresponds to the R -module

$$(S' \otimes_{S'} R) \otimes_R (R \otimes_S T) \cong R \otimes_S T.$$

Then we have

$$\begin{aligned} s|_{U_0 \times V_0} &= \sum_{\alpha=0}^n x_\alpha \Big|_{U_0} \otimes \frac{\partial}{\partial y_\alpha} \Big|_{V_0} \\ &= \sum_{\alpha=1}^n \left(-1 \otimes y_{\alpha/0} \frac{\partial}{\partial y_\alpha} + x_{\alpha/0} \otimes \frac{\partial}{\partial y_\alpha} \Big|_{V_0} \right) \\ &= \sum_{\alpha=1}^n (x_{\alpha/0} - y_{\alpha/0}) \otimes \frac{\partial}{\partial y_\alpha} \Big|_{V_0}. \end{aligned}$$

Thus the restriction of $\mathbb{V}(s)$ to $U_0 \times V_0$ is defined by the ideal $(x_{1/0} - y_{1/0}, \dots, x_{n/0} - y_{n/0})$, which gives the closed subscheme $\Delta \cap (U_0 \times V_0)$. \square

Lemma 2.11. For \mathcal{E} as given and $i \geq 1$, there is a natural isomorphism $\bigwedge^i(\mathcal{E}^\vee) \cong \mathcal{O}(-i) \boxtimes \Omega^i(i)$.

Proof. Observe we have natural isomorphisms

$$\begin{aligned} \mathcal{E}^\vee &= \mathcal{H}om(\pi_1^* \mathcal{O}(1) \otimes \pi_2^* \mathcal{T}(-1), \mathcal{O}_{\mathbb{P}^n \times \mathbb{P}^n}) \\ &\cong \mathcal{H}om(\pi_1^* \mathcal{O}(1), \mathcal{H}om(\pi_2^* \mathcal{T}(-1), \mathcal{O}_{\mathbb{P}^n \times \mathbb{P}^n})) \\ &\cong \mathcal{H}om(\pi_1^* \mathcal{O}(1), \mathcal{O}_{\mathbb{P}^n \times \mathbb{P}^n}) \otimes \mathcal{H}om(\pi_2^* \mathcal{T}(-1), \mathcal{O}_{\mathbb{P}^n \times \mathbb{P}^n}) \\ &= (\pi_1^* \mathcal{O}(1))^\vee \otimes (\pi_2^* \mathcal{T}(-1))^\vee \end{aligned}$$

where the first isomorphism comes from the $\otimes - \mathcal{H}om$ adjunction, and the second uses the fact that $\mathcal{O}(1)$ (and hence its pullback) is locally free; see Section II.5 of Hartshorne (2008). Moreover, since pullback commutes with dualising for locally free sheaves of finite rank, we have the isomorphisms

$$\begin{aligned} (\pi_1^* \mathcal{O}(1))^\vee &\cong \pi_1^* \mathcal{O}(-1), & (\pi_2^* \mathcal{T}(-1))^\vee &\cong \pi_2^* (\mathcal{T}(-1))^\vee \\ & & &\cong \pi_2^* (\mathcal{T}^\vee \otimes \mathcal{O}(1)) \\ & & &\cong \pi_2^* \Omega(1) \end{aligned}$$

hence $\mathcal{E}^\vee = \mathcal{O}(-1) \boxtimes \Omega(1)$, proving the result for $i = 1$. To prove the result for $i > 1$, we record the following observation.

Claim For locally free coherent sheaves \mathcal{L}, \mathcal{M} such that \mathcal{L} has rank 1, there is a natural isomorphism

$$\bigwedge^i(\mathcal{L} \otimes \mathcal{M}) \cong \mathcal{L}^{\otimes i} \otimes \bigwedge^i \mathcal{M}.$$

Given this, it is immediate that we have the isomorphisms

$$\begin{aligned} \bigwedge^i(\mathcal{E}^\vee) &= \bigwedge^i(\pi_1^* \mathcal{O}(-1) \otimes \pi_2^* \Omega(1)) \\ &\cong (\pi_1^* \mathcal{O}(-1))^{\otimes i} \otimes \bigwedge^i(\pi_2^* \Omega(1)) \\ &\cong \pi_1^* (\mathcal{O}(-1))^{\otimes i} \otimes \pi_2^* (\bigwedge^i \Omega(1)) \\ &\cong \pi_1^* \mathcal{O}(-i) \otimes \pi_2^* (\bigwedge^i \Omega \otimes (\mathcal{O}(1))^{\otimes i}) \\ &\cong \mathcal{O}(-i) \boxtimes \Omega^i(i). \end{aligned}$$

To prove the claim, observe that there is a natural map

$$(\mathcal{M} \otimes \mathcal{L})^{\otimes i} \rightarrow \mathcal{L}^{\otimes i} \otimes \bigwedge^i \mathcal{M}.$$

We show that this induces the required isomorphism $\bigwedge^i(\mathcal{L} \otimes \mathcal{M}) \cong \mathcal{L}^{\otimes i} \otimes \bigwedge^i \mathcal{M}$. This can be checked locally— on an affine open $\text{Spec } R$, say \mathcal{L} and \mathcal{M} are given by the free R -modules L and M respectively. Then the natural map

$$T_R(L \otimes_R M) \rightarrow \bigoplus_j (L^{\otimes j} \otimes_R \bigwedge_R^j M)$$

on the tensor algebra sends $(\ell \otimes m) \otimes (\ell \otimes m) \mapsto 0$, thus descending to a map on $\bigwedge^\bullet_R(L \otimes_R M)$. This induced map preserves grading. Thus we have a map $\bigwedge^i(\mathcal{L} \otimes \mathcal{M}) \rightarrow (\mathcal{L})^{\otimes i} \otimes \bigwedge^i \mathcal{M}$. Comparing ranks of the sheaves, this must be an isomorphism. \square

We now show that the Koszul complex of s gives the required resolution of the diagonal.

Theorem 2.12 (Beilinson (1978)). There is an exact sequence in $\mathcal{Coh}(\mathbb{P}^n \times \mathbb{P}^n)$ of the form

$$0 \rightarrow \mathcal{O}(-n) \boxtimes \Omega^n(n) \rightarrow \cdots \rightarrow \mathcal{O}(-1) \boxtimes \Omega(1) \rightarrow \mathcal{O}_{\mathbb{P}^n \times \mathbb{P}^n} \rightarrow \mathcal{O}_\Delta \rightarrow 0.$$

Proof. From Lemma 2.10 and Lemma 2.11, it is clear that the Koszul complex $\mathcal{K}_{\mathbb{P}^n \times \mathbb{P}^n}(s)$ has the required form. It remains to show the complex is exact. From the discussion in Remark 2.8, we can check this on the affine open sets $U_\alpha \times V_\alpha$ which cover $V(s)$.

Using the notation of Lemma 2.10, the coordinate ring of $U_0 \times V_0$ can be written as $R = S[z_1, \dots, z_n]$ where $z_\alpha = x_{\alpha/0} - y_{\alpha/0}$. Then the section $s|_{U_0 \times V_0}$ corresponds to the n -tuple (z_1, \dots, z_n) . Hence the restriction of $\mathcal{K}_{\mathbb{P}^n \times \mathbb{P}^n}(s)$ is given by $K_R(z_1, \dots, z_n)$, the Koszul complex associated to the regular sequence (z_1, \dots, z_n) in R . By Proposition 2.7, this is exact. \square

2.3.2 Generating the category

We now show that either of the two sets $\{\mathcal{O}, \mathcal{O}(-1), \dots, \mathcal{O}(-n)\}$ and $\{\mathcal{O}, \Omega(1), \dots, \Omega^n(n)\}$ generates $\mathbf{D}^b(\mathbb{P}^n)$. To do this, first observe the following consequence of the resolution of the diagonal.

Lemma 2.13. If \mathcal{A} is a bounded complex of coherent sheaves on \mathbb{P}^n , then \mathcal{A} is isomorphic to an object in the triangulated subcategory generated by

$$S = \{\Phi_{\mathcal{O}(-n) \boxtimes \Omega^n(n)}(\mathcal{A}), \Phi_{\mathcal{O}(-n+1) \boxtimes \Omega^{n-1}(n-1)}(\mathcal{A}), \dots, \Phi_{\mathcal{O}_{\mathbb{P}^n \times \mathbb{P}^n}}(\mathcal{A})\}.$$

Proof. We first split the exact sequence of Theorem 2.12 into short exact sequences

$$\begin{aligned} 0 \rightarrow \mathcal{O}(-n) \boxtimes \Omega^n(n) &\rightarrow \mathcal{O}(-n+1) \boxtimes \Omega^{n-1}(n-1) \rightarrow \mathcal{E}_1 \rightarrow 0 \\ 0 \rightarrow \mathcal{E}_1 &\rightarrow \mathcal{O}(-n+2) \boxtimes \Omega^{n-2}(n-2) \rightarrow \mathcal{E}_2 \rightarrow 0 \\ &\vdots \\ 0 \rightarrow \mathcal{E}_{n-2} &\rightarrow \mathcal{O}(-1) \boxtimes \Omega(1) \rightarrow \mathcal{E}_{n-1} \rightarrow 0 \\ 0 \rightarrow \mathcal{E}_{n-1} &\rightarrow \mathcal{O}_{\mathbb{P}^n \times \mathbb{P}^n} \rightarrow \mathcal{O}_\Delta \rightarrow 0 \end{aligned}$$

for coherent sheaves \mathcal{E}_i . These give distinguished triangles in $\mathbf{D}^b(\mathbb{P}^n)$. Since $\Phi_{-}(\mathcal{A}) : \mathbf{D}^b(\mathbb{P}^n) \rightarrow \mathbf{D}^b(\mathbb{P}^n)$ is an exact functor, we have distinguished triangles

$$\begin{aligned} \Phi_{\mathcal{O}(-n) \boxtimes \Omega^n(n)}(\mathcal{A}) &\rightarrow \Phi_{\mathcal{O}(-n+1) \boxtimes \Omega^{n-1}(n-1)}(\mathcal{A}) \rightarrow \Phi_{\mathcal{E}_1}(\mathcal{A}) \rightarrow \Phi_{\mathcal{O}(-n) \boxtimes \Omega^n(n)}(\mathcal{A})[1] \\ \Phi_{\mathcal{E}_1}(\mathcal{A}) &\rightarrow \Phi_{\mathcal{O}(-n+2) \boxtimes \Omega^{n-2}(n-2)}(\mathcal{A}) \rightarrow \Phi_{\mathcal{E}_2}(\mathcal{A}) \rightarrow \Phi_{\mathcal{E}_1}(\mathcal{A})[1] \\ &\vdots \\ \Phi_{\mathcal{E}_{n-2}}(\mathcal{A}) &\rightarrow \Phi_{\mathcal{O}(-1) \boxtimes \Omega(1)}(\mathcal{A}) \rightarrow \Phi_{\mathcal{E}_{n-1}}(\mathcal{A}) \rightarrow \Phi_{\mathcal{E}_{n-2}}(\mathcal{A})[1] \\ \Phi_{\mathcal{E}_{n-1}}(\mathcal{A}) &\rightarrow \Phi_{\mathcal{O}_{\mathbb{P}^n \times \mathbb{P}^n}}(\mathcal{A}) \rightarrow \Phi_{\mathcal{O}_\Delta}(\mathcal{A}) \rightarrow \Phi_{\mathcal{E}_{n-1}}(\mathcal{A})[1]. \end{aligned}$$

Hence the complexes $\Phi_{\mathcal{E}_1}(\mathcal{A}), \dots, \Phi_{\mathcal{E}_{n-1}}(\mathcal{A})$, and $\Phi_{\mathcal{O}_\Delta}(\mathcal{A}) \cong \mathcal{A}$ all lie in the triangulated subcategory generated by S . \square

Lemma 2.14. For any complex $\mathcal{A} \in \mathbf{D}^b(\mathbb{P}^n)$ and $i > 0$, the complex $\Phi_{\mathcal{O}(-i) \boxtimes \Omega^i(i)}(\mathcal{A})$ lies in the triangulated subcategory generated by $\mathcal{O}(-i)$, and also the triangulated subcategory generated by $\Omega^i(i)$.

Proof. Note the functors $(-\otimes \Omega^i(i))$, π_1^* , π_2^* are all exact so we have $\mathcal{O}(-i) \boxtimes \Omega^i(i) \cong \pi_1^* \mathcal{O}(-i) \otimes^L \pi_2^* \Omega^i(i)$ in $\mathbf{D}^b(\mathbb{P}^n \times \mathbb{P}^n)$. Now we can use the projection formula to show that

$$\begin{aligned} \Phi_{\mathcal{O}(-i) \boxtimes \Omega^i(i)}(\mathcal{A}) &= \mathbf{R}\pi_{1*}(\pi_2^* \mathcal{A} \otimes^L (\mathcal{O}(-1) \boxtimes \Omega^i(i))) \\ &= \mathbf{R}\pi_{1*}(\pi_2^* \mathcal{A} \otimes^L \pi_1^* \mathcal{O}(-i) \otimes^L \pi_2^* \Omega^i(i)) \\ &= \mathbf{R}\pi_{1*} \pi_2^*(\mathcal{A} \otimes \Omega^i(i)) \otimes^L \mathcal{O}(-i). \end{aligned}$$

From the commuting diagram

$$\begin{array}{ccc} \mathbb{P}^n \times \mathbb{P}^n & \xrightarrow{\pi_1} & \mathbb{P}^n \\ \downarrow \pi_2 & & \downarrow \tau \\ \mathbb{P}^n & \xrightarrow{\tau} & \text{Spec } k \end{array},$$

we can invoke flat base change (see Chapter III, Proposition 9.3 of (Hartshorne 2008)) to see that

$$\begin{aligned} \mathbf{R}\pi_{1*} \pi_2^*(\mathcal{A} \otimes \Omega^i(i)) &= \tau^*(\mathbf{R}\tau_*(\mathcal{A} \otimes \Omega^i(i))) \\ &= \tau^* \mathbf{R}\Gamma(\mathcal{A} \otimes \Omega^i(i)) \end{aligned}$$

is a complex of constant sheaves on \mathbb{P}^n . Now $\mathbf{R}\Gamma(\mathcal{A} \otimes \Omega^i(i))$ is the complex

$$\cdots \rightarrow 0 \rightarrow \Gamma(\mathcal{I}^0) \rightarrow \Gamma(\mathcal{I}^1) \rightarrow \Gamma(\mathcal{I}^2) \rightarrow \cdots$$

where $0 \rightarrow \mathcal{A} \otimes \Omega^i(i) \rightarrow \mathcal{I}^\bullet$ is an injective resolution. Being a complex of k -vector spaces, it splits and is quasi-isomorphic to the complex of its homologies

$$\cdots \rightarrow H^j(\mathcal{A} \otimes \Omega^i(i)) \rightarrow H^{j+1}(\mathcal{A} \otimes \Omega^i(i)) \rightarrow \cdots$$

where the morphisms are all 0. It follows that $\Phi_{\mathcal{O}(-i) \boxtimes \Omega^i(i)}(\mathcal{A})$ is quasi-isomorphic to the complex

$$\cdots \rightarrow H^j(\mathcal{A} \otimes \Omega^i(i)) \otimes_k \mathcal{O}(-i) \rightarrow H^{j+1}(\mathcal{A} \otimes \Omega^i(i)) \otimes_k \mathcal{O}(-i) \rightarrow \cdots$$

with zero differentials. In particular, it lies in the triangulated subcategory of $\mathbf{D}^b(\mathbb{P}^n)$ generated by $\{\mathcal{O}(-i)\}$, as required.

The result for $\Omega^i(i)$ follows similarly, switching the roles of π_1 and π_2 in the projection formula above. \square

The above result extends to show that $\Phi_{\mathcal{O}_{\mathbb{P}^n \times \mathbb{P}^n}}(\mathcal{A})$ lies in the triangulated subcategory generated by \mathcal{O} , since $\mathcal{O}_{\mathbb{P}^n \times \mathbb{P}^n} \cong \mathcal{O} \boxtimes \mathcal{O}$. Thus we have the desired conclusion.

Theorem 2.15. The category $\mathbf{D}^b(\mathbb{P}^n)$ is generated by either of the two sets $\{\mathcal{O}, \mathcal{O}(-1), \dots, \mathcal{O}(-n)\}$ and $\{\mathcal{O}, \Omega(1), \dots, \Omega^n(n)\}$.

Proof. Immediate from Lemma 2.13 and Lemma 2.14. \square

Example 2.16. On \mathbb{P}^1 , the Euler exact sequence (2) shows that $\Omega \cong \mathcal{O}(-2)$ is quasi-isomorphic to a complex of the form

$$\cdots \rightarrow 0 \rightarrow (\mathcal{O}(-1))^{\oplus 2} \rightarrow \mathcal{O} \rightarrow 0 \rightarrow \cdots. \quad (4)$$

Here is how we can deduce the same fact from Beilinson's theorem— applying $\Phi_-(\mathcal{O}(i))$ to the resolution of the diagonal yields an exact triangle in $\mathbf{D}^b(\mathbb{P}^1)$ given by

$$\mathbf{R}\Gamma(\mathcal{O}(i-1)) \otimes_k \mathcal{O}(-1) \xrightarrow{f} \mathbf{R}\Gamma(\mathcal{O}(i)) \otimes_k \mathcal{O} \rightarrow \mathcal{O}(i) \rightarrow \mathbf{R}\Gamma(\mathcal{O}(i-1)) \otimes_k \mathcal{O}(-1)[1].$$

Using the fact that $\mathbf{R}\Gamma(\mathcal{O}(i-1))$ and $\mathbf{R}\Gamma(\mathcal{O}(i))$ split, the map f is given by

$$\begin{array}{ccccccc} \cdots & \longrightarrow & 0 & \longrightarrow & A_{i-1} \otimes_k \mathcal{O}(-1) & \xrightarrow{0} & A_{-i-1} \otimes_k \mathcal{O}(-1) \longrightarrow 0 \longrightarrow \cdots \\ & & \downarrow & & \downarrow f^0 & & \downarrow f^1 \\ \cdots & \longrightarrow & 0 & \longrightarrow & A_i \otimes_k \mathcal{O} & \xrightarrow{0} & A_{-i-2} \otimes_k \mathcal{O} \longrightarrow 0 \longrightarrow \cdots \end{array},$$

where A is the ring $k[x_0, x_1]$. By definition of exact triangles, we must have that $\mathcal{O}(i)$ is quasi-isomorphic to the complex $\mathcal{Cone}(f)$ which has form

$$\cdots \rightarrow 0 \rightarrow A_{i-1} \otimes_k \mathcal{O}(-1) \xrightarrow{f^0} A_i \otimes_k \mathcal{O} \oplus A_{-i-1} \otimes_k \mathcal{O}(-1) \xrightarrow{(0, f^1)} A_{-i-2} \otimes_k \mathcal{O} \rightarrow 0 \rightarrow \cdots.$$

Noting that A_i has dimension $i+1$ when $i \geq 0$, we can deduce that $\mathcal{O}(-2) = \Omega$ is quasi-isomorphic to a complex of form given in (4). Likewise, $\mathcal{O}(1)$ is quasi-isomorphic to a complex of form

$$\cdots \rightarrow 0 \rightarrow \mathcal{O}(-1) \rightarrow \mathcal{O}^{\oplus 2} \rightarrow 0 \rightarrow \cdots,$$

which is again something we can explicitly confirm by considering twists of the Euler sequence.

General form of the decomposition. Building on the computation of Example 2.16, we can easily say what vector spaces underlie the decomposition of any sheaf $\mathcal{A} \in \mathcal{Coh} \mathbb{P}^n$ in terms of generators. Indeed, applying the $\Phi_-(\mathcal{A})$ functor to the exact triangles which resolve the diagonal yields a collection of exact triangles in $\mathbf{D}^b(\mathbb{P}^n)$, given by

$$\begin{aligned} \tau^* \mathbf{R}\Gamma(\mathcal{A}(-n)) \otimes \Omega^n(n) &\rightarrow \tau^* \mathbf{R}\Gamma(\mathcal{A}(-n+1)) \otimes \Omega^{n-1}(n-1) \rightarrow \mathcal{E}_1 \rightarrow \tau^* \mathbf{R}\Gamma(\mathcal{A}(-n)) \otimes \Omega^n(n)[1] \\ \mathcal{E}_1 &\rightarrow \tau^* \mathbf{R}\Gamma(\mathcal{A}(-n+2)) \otimes \Omega^{n-2}(n-2) \rightarrow \mathcal{E}_2 \rightarrow \mathcal{E}_1[1] \\ &\vdots \\ \mathcal{E}_{n-1} &\rightarrow \tau^* \mathbf{R}\Gamma(\mathcal{A}) \otimes \mathcal{O} \rightarrow \mathcal{A} \rightarrow \mathcal{E}_{n-1}[1]. \end{aligned}$$

Then the third complex in each exact triangle is isomorphic to the cone of the first triangle, so we have equalities of underlying complexes

$$\begin{aligned} \mathcal{E}_1 &= (\tau^* \mathbf{R}\Gamma(\mathcal{A}(-n+1)) \otimes \Omega^{n-1}(n-1)) \oplus (\tau^* \mathbf{R}\Gamma(\mathcal{A}(-n)) \otimes \Omega^n(n))[1] \\ \mathcal{E}_2 &= (\tau^* \mathbf{R}\Gamma(\mathcal{A}(-n+2)) \otimes \Omega^{n-2}(n-2)) \oplus \mathcal{E}_1[1] \\ &\vdots \\ \mathcal{A} &= (\tau^* \mathbf{R}\Gamma(\mathcal{A}) \otimes \mathcal{O}) \oplus \mathcal{E}_{n-1}[1]. \end{aligned}$$

Since the complexes of the form $\mathbf{R}\Gamma(-)$ split and are quasi-isomorphic to the complexes formed by homology, we can read off the vector spaces appearing in each differential degree in \mathcal{A} . The resulting computation is stated below.

Proposition 2.17. Given $\mathcal{A} \in \mathcal{Coh} \mathbb{P}^n$, there is a complex

$$\cdots \rightarrow \mathcal{B}^{-1} \rightarrow \mathcal{B}^0 \rightarrow \mathcal{B}^1 \rightarrow \cdots$$

where $\mathcal{B}^i = \bigoplus_j H^j(\mathcal{A}(i-j)) \otimes_k \Omega^{j-i}(j-i)$, which is exact everywhere except at \mathcal{B} where it has cohomology \mathcal{A} . This complex is the *Beilinson monad* for \mathcal{A} .

The maps in the Beilinson monad come from those in the Koszul complex, and can be computed explicitly. In Section 3.4, we state Eisenbud et al.'s (2003) result which computes the Beilinson monad from Koszul complexes of module categories, which are easier to manipulate and compute.

2.3.3 Equivalence with module categories

In fact Beilinson proves a stronger result than Theorem 2.15, showing that the category is generated by $\{\mathcal{O}, \mathcal{O}(-1), \dots, \mathcal{O}(-n)\}$ ‘as simply as possible’, i.e. by taking bounded complexes of finite direct sums of the sheaves. This is done by establishing an equivalence with the category $\mathbf{K}^b\{A, A\langle -1 \rangle, \dots, A\langle -n \rangle\}$ of graded A -modules, where $A = k[x_0, \dots, x_n]$ is the symmetric algebra on X and the category is as defined in Example 1.22. Since we know the generators of both the categories, we use Lemma 1.23.

Lemma 2.18. For any $0 \leq i, j \leq n$ and $m \in \mathbb{Z}$, we have

$$\begin{aligned} \mathrm{Hom}_{\mathbf{D}^b(\mathbb{P}^n)}(\Omega^i(i)[m], \Omega^j(j)) &= \mathrm{Hom}_{\mathbf{K}(A^!-\mathrm{grMod})}(A^!\langle i \rangle[m], A^!\langle j \rangle) \\ \mathrm{Hom}_{\mathbf{D}^b(\mathbb{P}^n)}(\mathcal{O}(-i)[m], \mathcal{O}(-j)) &= \mathrm{Hom}_{\mathbf{K}(A-\mathrm{grMod})}(A\langle i \rangle[m], A\langle j \rangle). \end{aligned}$$

Proof. Observe that we have

$$\mathrm{Hom}_{\mathbf{K}(A-\mathrm{grMod})}(A\langle -i \rangle[m], A\langle -j \rangle) = \begin{cases} A_{i-j}, & m = 0 \\ 0, & m \neq 0. \end{cases}$$

On the other hand, we have

$$\begin{aligned} \mathrm{Hom}_{\mathbf{D}^b(\mathbb{P}^n)}(\mathcal{O}(-i)[m], \mathcal{O}(-j)) &= \mathrm{Ext}_{\mathcal{O}(\mathbb{P}^n)}^m(\mathcal{O}(-i), \mathcal{O}(-j)) \\ &= \Gamma(\mathcal{E}xt^m(\mathcal{O}(-i), \mathcal{O}(-j))) \\ &= \Gamma(\mathcal{E}xt^m(\mathcal{O}, \mathcal{O}(-i)^\vee \otimes \mathcal{O}(-j))) \\ &= H^m(\mathcal{O}(i-j)) \\ &= \begin{cases} A_{i-j}, & m = 0 \\ 0, & m \neq 0. \end{cases} \end{aligned}$$

Likewise to prove the result for Ω , it suffices to show

$$\mathrm{Ext}_{\mathcal{O}(\mathbb{P}^n)}^m(\Omega^i(i), \Omega^j(j)) = \begin{cases} A_{j-i}^!, & m = 0 \\ 0, & m \neq 0. \end{cases}$$

This can be proven by induction, using the generalised Euler exact sequences (3) of Example 2.9. Note that we have $\bigwedge^i(\mathcal{O}(-1)^{\oplus(n+1)}) \cong \mathcal{O}(-i) \otimes_k \bigwedge^i V$ so the generalised Euler exact sequences become

$$0 \rightarrow \Omega^j(j) \rightarrow \mathcal{O} \otimes_k \bigwedge^j V \rightarrow \Omega^{j-1}(j) \rightarrow 0.$$

Then the complete induction argument can be found at Belmans (2015). \square

We can now prove the central result of Beilinson (1978).

Theorem 2.19 (Beilinson’s theorem). There are equivalences of categories

$$\mathbf{K}^b\{A\langle -n \rangle, \dots, A\langle -1 \rangle, A\} \cong \mathbf{D}^b(\mathbb{P}^n) \cong \mathbf{K}^b\{A^!\langle n \rangle, \dots, A^!\langle 1 \rangle, A^!\}.$$

Proof. Given Lemma 2.18, we can define additive functors

$$\begin{aligned} F_1 : \mathbf{K}^b\{A\langle -n \rangle, \dots, A\langle -1 \rangle, A\} &\longrightarrow \mathbf{D}^b(\mathbb{P}^n) \\ F_2 : \mathbf{K}^b\{A^!\langle n \rangle, \dots, A^!\langle 1 \rangle, A^!\} &\longrightarrow \mathbf{D}^b(\mathbb{P}^n) \end{aligned}$$

given by $A\langle -i \rangle \mapsto \mathcal{O}(-i)$ and $A^!\langle i \rangle \mapsto \Omega^i(i)$ respectively, the morphisms coming from the isomorphisms of Hom-groups. Since translations and cones are preserved, these are morphisms of triangulated categories. By Lemma 1.23, we immediately have that F_1 and F_2 are equivalences of categories. \square

3 The Bernstein-Gel'fand-Gel'fand correspondence

Beilinson's theorem hints at an underlying equivalence between the module categories of the symmetric algebra A and the exterior algebra $A^!$. This equivalence is proved in Bernstein et al. (1978), who define adjoint functors

$$\mathbf{C}(A\text{-grMod}) \xrightleftharpoons[F]{G} \mathbf{C}(A^!\text{-grMod})$$

which descend to equivalences of bounded derived categories. This is the so called 'BGG correspondence', named after the authors of the paper.

In this section we define the functors, and prove the adjunction between them comes from a $\otimes - \text{Hom}$ adjunction. Then the natural maps $G(F(\mathbf{M})) \rightarrow \mathbf{M}$ and $\mathbf{N} \rightarrow F(G(\mathbf{N}))$ turn out to be free and injective resolutions respectively, showing that F and G become equivalences after inverting quasi-isomorphisms. The material largely follows the treatment in Eisenbud et al. (2003), though we also provide a description of the functors using A^i (the coalgebra dual to $A^!$) by adapting the construction in Keller (2003). Keller's paper elaborates on this coalgebra construction, which is more succinct and applicable in broader contexts.

3.1 Twisted functors

We now define additive functors

$$\mathbf{C}(A\text{-grMod}) \xrightleftharpoons[F]{G} \mathbf{C}(A^!\text{-grMod})$$

on which the BGG correspondence is based. In the framework of \mathbb{Z}^2 -graded vector spaces described in Section 1.2.3, we have

$$\bigoplus_{i,j} F(\mathbf{M})_j^i \cong \text{Hom}_k \left(A^!, \bigoplus_{p,q} M_q^p \right) = A^i \otimes_k \left(\bigoplus_{p,q} M_q^p \right), \quad \bigoplus_{i,j} G(\mathbf{N})_j^i \cong A \otimes_k \left(\bigoplus_{p,q} N_q^p \right).$$

However, care is needed to define the gradings and differentials since, for example, naively applying the functor $\text{Hom}_k(A^!, -)$ would lose all A -module structure. The key is to modify the naïve differential by adding a 'twist' as in Example 0.1.

3.1.1 Defining the functor F

We first define F on the category $A\text{-grMod}$, seen as the full subcategory of $\mathbf{C}(A\text{-grMod})$ whose objects are complexes concentrated in differential degree 0. If M_\bullet^0 is a graded A -module, we define $F(M_\bullet^0)$ to be the chain complex of $A^!$ -modules given by a

$$\begin{aligned} \cdots \rightarrow A^i\langle -i \rangle \otimes_k M_i^0 &\xrightarrow{\partial} A^i\langle -i-1 \rangle \otimes_k M_{i+1}^0 \rightarrow \cdots \\ a \otimes m &\mapsto \sum_{\alpha} \xi_{\alpha} a \otimes x_{\alpha} m. \end{aligned}$$

The module $A^i\langle -i \rangle \otimes_k M_i^0$ is naturally isomorphic to $\text{Hom}_k(A^i\langle i \rangle, M_i^0)$ and inherits an Adam's grading from A^i with the vector space $A_{j-i}^i \otimes_k M_i^0$ forming the j th graded piece. These shifts in grading have been chosen precisely so that the differential ∂ has degree $(1, 0)$, while the graded commutativity of A^i implies $\partial \circ \partial = 0$. Thus we indeed have a chain complex of A^i -modules.

Given a morphism $M_\bullet^0 \rightarrow M_\bullet^1$ in $A\text{-grMod}$, the functoriality of tensor products induces A^i -module homomorphisms $A^i\langle -i \rangle \otimes_k M_i^0 \rightarrow A^i\langle -i \rangle \otimes_k M_i^1$ which are compatible with the differentials (i.e. the natural squares commute). Thus we have an additive functor $F : A\text{-grMod} \rightarrow \mathbf{C}(A^i\text{-grMod})$.

To extend F to arbitrary chain complexes $\mathbf{M} = (\bigoplus_{i,j} M_j^i, d) \in \mathbf{C}(A\text{-grMod})$, we observe that the functoriality of F gives us a (commuting) bicomplex

$$\begin{array}{ccccccc}
 \vdots & & \vdots & & \vdots & & \\
 \uparrow & & \uparrow & & \uparrow & & \\
 F(M_\bullet^{i+1}) & \cdots \longrightarrow & A^i\langle -j \rangle \otimes_k M_j^{i+1} & \longrightarrow & A^i\langle -j-1 \rangle \otimes_k M_{j+1}^{i+1} & \longrightarrow & \cdots \\
 \uparrow & & \uparrow & & \uparrow & & \\
 F(M_\bullet^i) & \cdots \longrightarrow & A^i\langle -j \rangle \otimes_k M_j^i & \longrightarrow & A^i\langle -j-1 \rangle \otimes_k M_{j+1}^i & \longrightarrow & \cdots \\
 \uparrow & & \uparrow & & \uparrow & & \\
 \vdots & & \vdots & & \vdots & &
 \end{array} \tag{5}$$

where the vertical maps are $1 \otimes d$. Define $F(\mathbf{M})$ to be the total complex of this bicomplex, i.e. $F(\mathbf{M})$ is given by

$$\begin{aligned}
 \cdots \rightarrow \bigoplus_{p+q=i} A^i\langle -q \rangle \otimes_k M_q^p &\xrightarrow{\partial} \bigoplus_{p+q=i+1} A^i\langle -q \rangle \otimes_k M_q^p \rightarrow \cdots, \\
 \partial : a \otimes m &\longmapsto a \otimes dm + (-1)^{\#m} \sum_{\alpha} \xi_{\alpha} a \otimes x_{\alpha} m
 \end{aligned} \tag{6}$$

where $\#m$ is the differential degree of $m \in \mathbf{M}$.

The twist using comodules. Observe that the differential ∂ differs from the naïve differential $1 \otimes d$ by the horizontal maps, which are the ‘twists’ we have been alluding to. These have a nice description using the fact that a graded module $N_\bullet \in A^i\text{-grMod}$ has the structure of a graded A^i -comodule via the map

$$\begin{aligned}
 \Delta : N_\bullet &\longrightarrow N_\bullet \otimes_k A^i \\
 n &\longmapsto \sum_{\alpha \subseteq \{0, \dots, n\}} \xi_{\alpha} n \otimes x_{\alpha}.
 \end{aligned}$$

Applying this idea to the A^i -modules $A^i\langle -i \rangle$, we get a commuting square

$$\begin{array}{ccc}
 \bigoplus_{u+v=j-q} A_u^i \otimes_k A_v^i \otimes_k M_q^{i-q} & \xrightarrow{1 \otimes \tau \otimes 1} & A_{j-q-1}^i \otimes_k A_1 \otimes_k M_q^{i-q} \\
 \Delta \otimes 1 \uparrow & & \downarrow 1 \otimes \nabla \\
 A_{j-q}^i \otimes_k M_q^{i-q} & \xrightarrow{(-1)^{i-q}(\partial - 1 \otimes d)} & A_{j-q-1}^i \otimes_k M_{q+1}^{i-q}
 \end{array}$$

where $\nabla : A \otimes_k M_{\bullet}^{i-q} \rightarrow M_{\bullet}^{i-q}$ defines the A -module structure on M , and $\tau : A^i \rightarrow A^!$ is the morphism defined in Section 1.2.2. This morphism identifies A_1^i with A_1 , annihilating other graded pieces.

In summary, $F(\mathbf{M})$ as a \mathbb{Z}^2 -graded vector space is simply $A^i \otimes_k \mathbf{M}$ with (i, j) th piece

$$F(\mathbf{M})_j^i = \bigoplus_{p+q=i} A_{j-q}^i \otimes_k M_q^p$$

and differential given on $A_{j-q}^i \otimes_k M_q^p$ by

$$1 \otimes d + (-1)^p (1 \otimes \nabla) \circ (1 \otimes \tau \otimes 1) \circ (\Delta \otimes 1).$$

3.1.2 The left adjoint to F

The functor $G : \mathbf{C}(A^!-\text{grMod}) \rightarrow \mathbf{C}(A-\text{grMod})$ is analogously defined, and maps the chain complex $\mathbf{N} = (\bigoplus_{i,j} N, \partial)$ to $G(\mathbf{N})$ given by

$$\begin{aligned} \cdots \rightarrow \bigoplus_{p-q=i} N_q^p \otimes_k A\langle -q \rangle &\xrightarrow{d} \bigoplus_{p-q=i+1} N_q^p \otimes_k A\langle -q \rangle \rightarrow \cdots \\ d : n \otimes a &\mapsto \partial n \otimes a + (-1)^{\#n} \sum_{\alpha} \xi_{\alpha} n \otimes x_{\alpha} a \end{aligned} \quad (7)$$

where $\#n$ is the differential degree of $n \in \mathbf{N}$. The Adam's grading on each $G(\mathbf{N})_{\bullet}^i$ is inherited from A , and is given by

$$G(\mathbf{N})_j^i = \bigoplus_{p-q=i} N_q^p \otimes_k A_{j-q}.$$

Recalling that every $A^!$ -module is a $A^!$ -comodule (see Section 3.1.1), we can use the comodule structure-map $\Delta : N_{\bullet}^i \rightarrow N_{\bullet}^i \otimes A^!$ to define the differential on $N_q^p \otimes_k A_{j-q}$ as

$$\partial \otimes 1 + (-1)^p (1 \otimes \nabla) \circ (1 \otimes \tau \otimes 1) \circ (\Delta \otimes 1).$$

The adjunction. Having defined the functors F and G , we show that G is left adjoint to F . Spelled out this means given $\mathbf{M} \in \mathbf{C}(A-\text{grMod})$ and $\mathbf{N} \in \mathbf{C}(A^!-\text{grMod})$, there is a natural isomorphism of abelian groups

$$\text{Hom}_{\mathbf{C}(A-\text{grMod})}(G(\mathbf{N}), \mathbf{M}) \cong \text{Hom}_{\mathbf{C}(A^!-\text{grMod})}(\mathbf{N}, F(\mathbf{M})).$$

At its heart this is a \otimes -Hom adjunction, as we shall illustrate in the special case of module categories below.

Lemma 3.1. Given modules $M \in A\text{-Mod}$ and $N \in A^!\text{-Mod}$, there are natural isomorphisms of abelian groups

$$\text{Hom}_A(A \otimes_k N, M) \cong \text{Hom}_k(N, M) \cong \text{Hom}_{A^!}(N, \text{Hom}_k(A^!, M)).$$

Proof. Consider the $(A, A^!)$ -bimodule $T = A \otimes_k A^!$. Then the standard \otimes -Hom adjunction for bimodules (Bourbaki 1989) gives us a natural isomorphism

$$\text{Hom}_A(T \otimes_{A^!} N, M) \cong \text{Hom}_{A^!}(N, \text{Hom}_A(T, M)).$$

Then observe that there are natural isomorphisms

$$T \otimes_{A^!} N \cong A \otimes_k A^! \otimes_{A^!} N \cong A \otimes_k N, \quad \text{Hom}_A(T, M) \cong \text{Hom}_A(A \otimes_k A^!, M) \cong \text{Hom}_k(A^!, M).$$

The isomorphism with $\text{Hom}_k(N, M)$ comes similarly from treating A as an (A, k) -bimodule. \square

We now exhibit the general adjunction for F and G , and it is here that the flexibility of interpreting a chain complex \mathbf{M} of graded modules as a single \mathbb{Z}^2 -graded module $\bigoplus_{i,j} M_j^i$ (see Section 1.2.3) really comes handy. Interpreting $\mathbf{C}(A\text{-grMod})$ as a subcategory of $A\text{-Mod}$ (likewise for A^\dagger), we use Lemma 3.1 to identify $\text{Hom}_{\mathbf{C}(A\text{-grMod})}(G(\mathbf{N}), \mathbf{M}) \subset \text{Hom}_A(\mathbf{N} \otimes_k A, \mathbf{M})$ and $\text{Hom}_{\mathbf{C}(A^\dagger\text{-grMod})}(\mathbf{N}, F(\mathbf{M})) \subset \text{Hom}_{A^\dagger}(\mathbf{N}, \text{Hom}_k(A^\dagger, \mathbf{M}))$ with the same subgroup of $\text{Hom}_k(\mathbf{N}, \mathbf{M})$.

Theorem 3.2 (Bernstein et al. (1978)). The functor G , from the category of complexes of graded A^\dagger -modules to the category of complexes of graded A -modules, is a left adjoint to the functor F .

Proof. Given $\bar{\varphi} \in \text{Hom}_A(G(\mathbf{N}), \mathbf{M})$, the corresponding map $\varphi \in \text{Hom}_k(\mathbf{N}, \mathbf{M})$ found in Lemma 3.1 is given by $\varphi(n) = \bar{\varphi}(n \otimes 1)$. Thus $\bar{\varphi}$ has degree $(0, 0)$ if and only if $\bar{\varphi}(N_j^i \otimes_k A_0) \subseteq M_j^{i-j}$, if and only if $\varphi(N_j^i) \subseteq M_j^{i-j}$. Moreover for $n \in N_j^i$, direct computation shows

$$(d_M \circ \bar{\varphi} - \bar{\varphi} \circ d_{G(\mathbf{N})})(n \otimes 1) = (d_M \circ \varphi - \varphi \circ \partial_N)(n) - (-1)^i \sum_{\alpha} x_{\alpha} \varphi(\xi_{\alpha} n),$$

thus $\bar{\varphi}$ is a morphism in $\mathbf{C}(A\text{-grMod})$ if and only if

$$\varphi(N_j^i) \subseteq M_j^{i-j}, \quad \text{and} \quad d_M \circ \varphi - \varphi \circ \partial_N = \sum_{\alpha} x_{\alpha} \varphi \xi_{\alpha} \quad (8)$$

where we write $\sum_{\alpha} x_{\alpha} \varphi \xi_{\alpha}$ for the map that takes $n \in N_j^i$ to $(-1)^i \sum_{\alpha} x_{\alpha} \varphi(\xi_{\alpha} n)$.

Likewise given $\varphi^\dagger \in \text{Hom}_{A^\dagger}(\mathbf{N}, F(\mathbf{M}))$, repeating the above argument shows φ^\dagger is an element of $\text{Hom}_{\mathbf{C}(A^\dagger\text{-grMod})}(\mathbf{N}, F(\mathbf{M}))$ if and only if the corresponding map $\varphi \in \text{Hom}_k(\mathbf{N}, \mathbf{M})$ satisfies (8). This shows that the isomorphisms given in Lemma 3.1 restrict to isomorphisms

$$\text{Hom}_{\mathbf{C}(A\text{-grMod})}(G(\mathbf{N}), \mathbf{M}) \cong \{\varphi \in \text{Hom}_k(\mathbf{N}, \mathbf{M}) \text{ satisfying (8)}\} \cong \text{Hom}_{\mathbf{C}(A^\dagger\text{-grMod})}(\mathbf{N}, F(\mathbf{M}))$$

thereby showing G is left adjoint to F . □

3.2 BGG resolutions

Given a complex $\mathbf{M} \in \mathbf{C}(A\text{-grMod})$, the adjunction $F \dashv G$ takes the identity morphism

$$1_{F(\mathbf{M})} \in \text{Hom}_{\mathbf{C}(A^\dagger\text{-grMod})}(F(\mathbf{M}), F(\mathbf{M}))$$

to a map

$$\varepsilon_{\mathbf{M}} \in \text{Hom}_{\mathbf{C}(A\text{-grMod})}(G(F(\mathbf{M})), \mathbf{M}).$$

The natural transformation $\varepsilon : G \circ F \rightarrow \mathbf{1}$ thus obtained is called the *counit* of the adjunction, and we say the morphism $\varepsilon_{\mathbf{M}}$ is the *component* of the transformation at \mathbf{M} . Likewise, there is the dual notion called the *unit* of the adjunction, which is a natural transformation $\eta : \mathbf{1} \rightarrow F \circ G$ giving, for any $\mathbf{N} \in \mathbf{C}(A^\dagger\text{-grMod})$, a morphism $\eta_{\mathbf{N}} : \mathbf{N} \rightarrow F(G(\mathbf{N}))$.

Begin with the following observation.

Proposition 3.3. The functor F maps elements of $\mathbf{C}(A\text{-grMod})$ to complexes of injective A^\dagger -modules, and the functor G maps elements of $\mathbf{C}(A^\dagger\text{-grMod})$ to complexes of free A -modules.

Proof. The statement for G is immediate from definition, so we prove that for any $\mathbf{M} \in \mathbf{C}(A\text{-grMod})$, the modules $F(\mathbf{M})_\bullet^i$ are injective over A^\dagger .

Recall from Weibel (2003) (Proposition 2.3.10) that if $R : \mathfrak{B} \rightarrow \mathfrak{A}$ is an additive functor which is right adjoint to an exact functor $L : \mathfrak{A} \rightarrow \mathfrak{B}$, then for any injective object $I \in \mathfrak{B}$ the object $R(I) \in \mathfrak{A}$ is injective. Applying this to the pair of adjoint functors

$$R = \text{Hom}_k(A^!, -) : k\text{-grMod} \rightarrow A^!\text{-grMod}, \quad L = (- \otimes_k A^!) : A^!\text{-grMod} \rightarrow k\text{-grMod}$$

and observing that L is exact since all k -vector spaces are flat over k , see that R preserves injectives. But every k -vector space is also injective, so the $A^!$ -modules $A^i\langle -q \rangle \otimes_k M_q^p \cong R(M_q^p)$ appearing in (6) are all injective.

To conclude, observe that the k -algebra $A^!$ is finite dimensional hence noetherian. By the theorem of Bass & Papp (see Lam (1999), Theorem 3.46) which asserts that a ring is (left) noetherian if and only if any direct sum of injective modules over it is injective, we are done. \square

We show that the component $\varepsilon_{\mathbf{M}} : G(F(\mathbf{M})) \rightarrow \mathbf{M}$ is, in fact, a free resolution of the complex \mathbf{M} and dually, the component $\eta_{\mathbf{N}}$ is an injective resolution of \mathbf{N} . A special but important case of this phenomenon is when the complex is

$$\cdots \rightarrow 0 \rightarrow k \rightarrow 0 \rightarrow \cdots,$$

and this will be central in proving the result for general complexes.

Example 3.4. The 1-dimensional vector space k can be considered a graded A -module concentrated degree 0, such that all $x_i \in A$ annihilate k . Then $F(k)$ is the complex $0 \rightarrow \Lambda^\bullet(X) \rightarrow 0$ concentrated in differential degree 0. We compute the complex $G(F(k))$ to be

$$0 \rightarrow A_{n+1}^i \otimes_k A\langle -n-1 \rangle \rightarrow A_n^i \otimes_k A\langle -n \rangle \rightarrow \cdots \rightarrow A_1^i \otimes_k A\langle -1 \rangle \rightarrow A_0^i \otimes_k A \rightarrow 0 \quad (9)$$

$$(x_{\alpha_1} \wedge \cdots \wedge x_{\alpha_i}) \otimes 1 \mapsto \sum_j (-1)^{i+j-1} (x_{\alpha_1} \wedge \cdots \wedge \hat{x}_{\alpha_j} \wedge \cdots \wedge x_{\alpha_i}) \otimes x_{\alpha_j}$$

where $\hat{}$ denotes omission of a term. This is precisely the Koszul complex associated to the regular sequence (x_0, \dots, x_n) in A , so by Proposition 2.7, we see that it is exact everywhere except in degree 0 where it has cohomology k . A similar result can be proven for the complex $F(G(k))$.

Thus we have resolutions

$$G(F(k)) \rightarrow k \rightarrow 0, \quad 0 \rightarrow k \rightarrow F(G(k))$$

of k by free A -modules and by injective $A^!$ -modules, respectively. It is not hard to see that these maps are precisely the ones given by the counit and the unit of adjunction.

3.2.1 Resolutions in general

We show that the counit (resp. unit) gives a free (resp. injective) resolution, by first showing this is the case for graded modules (i.e. complexes concentrated in a single degree). When \mathbf{M} is a graded A -module, we will show that the complex $G(F(\mathbf{M}))$ is ‘built up’ from the tensor product of \mathbf{M} and the Koszul complex of k .

Lemma 3.5 (Eisenbud et al. (2003)). If $\mathbf{M} \in \mathbf{C}(A\text{-grMod})$ is a chain complex concentrated in differential degree 0, then the natural map

$$\varepsilon_{\mathbf{M}} : G(F(\mathbf{M})) \rightarrow \mathbf{M}$$

is an epimorphism and induces an isomorphism on cohomology. Likewise, if $\mathbf{N} \in \mathbf{C}(A^!-\text{grMod})$ is a chain complex concentrated in differential degree 0, then the natural map

$$\eta_{\mathbf{N}} : \mathbf{N} \rightarrow F(G(\mathbf{N}))$$

is a monomorphism and induces an isomorphism on cohomology.

Proof. We first show that the complex $G(F(\mathbf{M}))$ has the same cohomology as the complex \mathbf{M} . Direct computation shows $G(F(\mathbf{M}))$ is given by

$$\begin{aligned} \cdots \rightarrow \bigoplus_p A_{-i}^i \otimes_k M_p^0 \otimes_k A\langle i-p \rangle &\longrightarrow \bigoplus_p A_{-i-1}^i \otimes_k M_p^0 \otimes_k A\langle i+1-p \rangle \rightarrow \cdots, \\ a \otimes m \otimes b &\longmapsto \sum_{\alpha} \xi_{\alpha} a \otimes x_{\alpha} m \otimes b + (-1)^{\deg m} \sum_{\alpha} \xi_{\alpha} a \otimes m \otimes x_{\alpha} b \end{aligned}$$

so the strand of this in Adam's degree r is seen to be the total complex of the (commuting) bicomplex

$$\begin{array}{ccccccc} & & \vdots & & \vdots & & \\ & & \downarrow & & \downarrow & & \\ \cdots & \longrightarrow & A_{-i}^i \otimes_k M_p^0 \otimes_k A_{i-p+r} & \longrightarrow & A_{-i-1}^i \otimes_k M_p^0 \otimes_k A_{i-p+r+1} & \longrightarrow & \cdots \\ & & \downarrow & & \downarrow & & \\ \cdots & \longrightarrow & A_{-i-1}^i \otimes_k M_{p+1}^0 \otimes_k A_{i-p+r} & \longrightarrow & A_{-i-2}^i \otimes_k M_{p+1}^0 \otimes_k A_{i-p+r+1} & \longrightarrow & \cdots \\ & & \downarrow & & \downarrow & & \\ & & \vdots & & \vdots & & \end{array} \quad . \quad (10)$$

It is clear that this bicomplex is bounded. Here p th row is obtained by applying $(- \otimes_k M_p^0)$ to the complex $G(F(k))_{r-p}$, so is exact from Proposition 2.7 unless $p = r$. Moreover, the r th row is

$$\cdots 0 \rightarrow M_r^0 \rightarrow 0 \rightarrow \cdots .$$

Thus first page of the spectral sequence (starting with horizontal cohomology) of (10) is

$$\begin{array}{ccccccc} & & \vdots & & \vdots & & \\ & & \downarrow & & \downarrow & & \\ \cdots & & M_r^0 & & 0 & & \cdots \\ & & \downarrow & & \downarrow & & \\ \cdots & & 0 & & 0 & & \cdots \\ & & \downarrow & & \downarrow & & \\ & & \vdots & & \vdots & & \end{array} .$$

By Proposition 1.7, we conclude that the spectral sequence converges and hence the total complex $G(F(\mathbf{M}))_r$ has cohomology

$$H^k(G(F(\mathbf{M}))_r) = \begin{cases} M_r^0, & k = 0 \\ 0, & \text{otherwise} \end{cases} .$$

Now it suffices to show that the map $\varepsilon_{\mathbf{M}} : G(F(\mathbf{M}))_r^0 \rightarrow M_r^0$ is the cokernel of $G(F(\mathbf{M}))_r^{-1} \rightarrow G(F(\mathbf{M}))_r^0$. But this is immediate because the sequence

$$\begin{aligned} \bigoplus_p X \otimes_k M_p^0 \otimes_k \text{Sym}^{r-p-1}(X) &\longrightarrow \bigoplus_p M_p^0 \otimes_k \text{Sym}^{r-p}(X) \longrightarrow M_r^0 \longrightarrow 0 \\ x_\alpha \otimes m \otimes a &\longmapsto m \otimes x_\alpha a + (-1)^{\deg m} x_\alpha m \otimes a \\ m \otimes a &\longmapsto (-1)^{\deg m} a m \end{aligned}$$

is (split) exact.

The analogous statement about graded $A^!$ -modules follows from a similar calculation. \square

The argument to extend this result to all chain complexes is purely formal.

Theorem 3.6 (Eisenbud et al. (2003)). For any complex $\mathbf{M} \in \mathbf{C}(A\text{-grMod})$, the complex $G(F(\mathbf{M}))$ is a free resolution of \mathbf{M} which surjects onto \mathbf{M} , and for any complex $\mathbf{N} \in \mathbf{C}(A^!\text{-grMod})$, the complex $F(G(\mathbf{N}))$ is an injective resolution of \mathbf{N} which \mathbf{N} injects into.

Proof. Given $\mathbf{M} \in \mathbf{C}(A\text{-grMod})$, the surjectivity of $\varepsilon_{\mathbf{M}} : G(F(\mathbf{M})) \rightarrow \mathbf{M}$ can be checked on the level of underlying \mathbb{Z}^2 -graded modules. The map is given on the (i, j) th component by

$$\begin{aligned} \bigoplus_{p,q} \text{Hom}_k(A_{q-i}^!, M_{p-q}^q) \otimes_k A_{j-p+i} &\longrightarrow M_j^i \\ f \otimes a &\mapsto af(1) \end{aligned}$$

hence any $m \in M_j^i$ can be written $\varepsilon_{\mathbf{M}}(f_m \otimes 1)$ where $f_m : A_0^! \rightarrow M_j^i$ is the function $f_m(1) = m$.

To prove that the induced map on cohomology is an isomorphism we first reduce the problem to bounded complexes using formal properties of the functors, and then induct on the length of the complex to reduce our problem to Lemma 3.5. The key properties we use are naturality of ε , and the fact that G , F and cohomology functors all preserve direct limits.

Any complex $\mathbf{M} \in \mathbf{C}(A\text{-grMod})$ can be written as the direct limit of bounded complexes $(\mathbf{M}^b)_{b \in B}$, giving us commuting diagrams

$$\begin{array}{ccc} G(F(\mathbf{M}^b)) & \longrightarrow & G(F(\mathbf{M})) \xlongequal{\quad} \varinjlim G(F(\mathbf{M}^b)) \\ \downarrow \varepsilon_{\mathbf{M}^b} & & \downarrow \varepsilon_{\mathbf{M}} \\ \mathbf{M}^b & \longrightarrow & \mathbf{M} \xlongequal{\quad} \varinjlim \mathbf{M}^b \end{array} \quad . \quad (11)$$

Then applying the i th cohomology functor H^i to (11) then shows that the map $H^i(\varepsilon_{\mathbf{M}})$ is the limit of the maps $\varepsilon_{\mathbf{M}^b}$, so to show $H^i(\varepsilon_{\mathbf{M}})$ is an isomorphism it suffices to show all $H^i(\varepsilon_{\mathbf{M}^b})$ are. Thus without loss of generality the complex \mathbf{M} is bounded. Since F and G respect translation in differential degree, say \mathbf{M} has form

$$0 \rightarrow M_\bullet^0 \rightarrow \dots \rightarrow M_\bullet^d \rightarrow 0. \quad (12)$$

Let \mathbf{M}^d be the chain complex with M_\bullet^d in degree d , and 0 elsewhere. We have a short exact sequence

$$0 \longrightarrow \ker(\varphi) \longrightarrow \mathbf{M} \xrightarrow{\varphi} \mathbf{M}^d \longrightarrow 0$$

where φ is the obvious map. The complex $\ker(\varphi)$ is concentrated in degrees $0, \dots, d-1$. Applying the exact functor H^i to the diagram formed by the naturality squares of ε on (12) gives us a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^i(G(F(\ker \varphi))) & \longrightarrow & H^i(G(F(\mathbf{M}))) & \longrightarrow & H^i(G(F(\mathbf{M}^d))) \longrightarrow 0 \\ & & \downarrow H^i(\varepsilon_{\ker \varphi}) & & \downarrow H^i(\varepsilon_{\mathbf{M}}) & & \downarrow (H^i \varepsilon_{\mathbf{M}^d}) \\ 0 & \longrightarrow & H^i(\ker(\varphi)) & \longrightarrow & H^i(\mathbf{M}) & \longrightarrow & H^i(\mathbf{M}^d) \longrightarrow 0 \end{array}$$

where the rows are exact. By Lemma 3.5, $H^i(\varepsilon_{\mathbf{M}^d})$ is an isomorphism. By the Five lemma, $H^i(\varepsilon_{\mathbf{M}})$ is an isomorphism if and only if $H^i(\varepsilon_{\ker(\varphi)})$ is. Since $\ker(\varphi)$ is a strictly shorter complex than \mathbf{M} , we are done.

The analogous statement for $F \circ G$ follows from a similar calculation. \square

Thus we have a formulaic (albeit inefficient– the free A -module A is resolved to an n -term free complex) method to compute resolutions of complexes. As an application we will see how this can be used to compute the projective dimension of the polynomial ring.

Syzygies and regularity of modules. We use the resolutions produced in Theorem 3.6 to prove a classical result of commutative algebra– Hilbert’s syzygy theorem, and provide a way to compute the Castelnuovo-Mumford regularity of modules. We briefly discuss the notions involved, and refer to Eisenbud (1995) for details.

Writing a graded A -module M in terms of generators and relations produces a short exact sequence

$$0 \rightarrow S \rightarrow F \rightarrow M \rightarrow 0,$$

where F is a free module. The module S is unique up to direct sum with a free module (i.e. if $0 \rightarrow S' \rightarrow F' \rightarrow M \rightarrow 0$ is another such resolution then there are free modules L and L' such that $L \oplus S \cong L' \oplus S'$), and is called the *first syzygy* of M . Continuing the process, we can write S in terms of generators and relations and define the second syzygy of M to be the first syzygy of S . Thus the j th syzygy of M is the module S_j (up to direct sum with a free module) such that there is an exact sequence

$$0 \rightarrow S_j \rightarrow F_{j-1} \rightarrow \dots \rightarrow F_0 \rightarrow M \rightarrow 0$$

where F_0, \dots, F_{j-1} are free modules. Note that if the j th syzygy of M is free then M has a free resolution of length $j+1$ – thus syzygies form a measure of the ‘complexity’ of M . This is made precise using the notion of *projective dimension*, defined as

$$\text{pd}(M) = \min\{j \mid \text{the } j\text{th syzygy module of } M \text{ is free or projective}\}.$$

Hilbert showed that the projective dimension of A -modules is bounded. The resolution produced using Theorem 3.6 provides an immediate constructive proof of this result.

Corollary 3.7 (Hilbert Syzygy Theorem). If M is a graded module over $k[x_0, \dots, x_n]$, then the $n+1$ st syzygy module of M is free.

In fact, this bound is strict– for instance, the A -module k has projective dimension $n+1$. To see this, observe that (9) allows us to compute $\text{Ext}_A^{n+1}(k, k) \cong k$ but by Lemma 4.1.6 of Weibel (2003),

an A -module M has a projective resolution of length $\leq d$ if and only if $\text{Ext}_A^d(M, N) = 0$ for all A -modules N . Defining the *graded global dimension* of a graded ring R to be

$$\text{gr.gl.dim}(R) = \sup\{\text{pd}(M) \mid M \in R\text{-grMod}\},$$

we have thus shown that $\text{gr.gl.dim}(k[x_0, \dots, x_n]) = n + 1$.

The notion of *Castelnuovo-Mumford regularity* builds upon this, putting a bound on the degrees of generators and relations of a finitely generated graded A -module M . We say M is m -regular if the j th syzygy of m is generated in degrees $\leq m + j$. We state a homological characterisation of regularity, referring to Eisenbud & Goto (1984) for a proof.

Theorem 3.8 (Eisenbud & Goto (1984)). For a finitely generated graded A -module M , the following conditions are equivalent.

1. M is m -regular.
2. $M_{\geq m} = \bigoplus_{i \geq m} M_i$ is generated by M_m and has a *linear free resolution* (a free resolution in which the differentials are represented by matrices whose entries have degree ≤ 1 .)
3. $M_{\geq m}$ is generated by M_m and $\text{Tor}_A(M, k)_j^{j-i} = 0$ for all j and all $i > m$.

Using the Koszul complex (9), we extend the result above to the following.

Corollary 3.9 (Eisenbud et al. (2003)). A finitely generated graded A -module M is m -regular if and only if $M_{\geq m}$ is generated by M_m and the complex $F(M)$ is exact at degrees $> m$.

Proof. It suffices to show that the complex $F(M)$ has cohomology $H^i(F(M))_j = \text{Tor}_A(M, k)_j^{j-i}$. To see this, note that the Koszul complex $G(F(k))$ given by (9) is a free resolution of k . Then the complex $M \otimes_A G(F(k))$ is given in differential degree $i - j$ by

$$M \otimes_A A_{j-i}^! \otimes_k A\langle i - j \rangle \cong A_{j-i}^! \otimes_k M\langle i - j \rangle.$$

The component in Adam's degree j is $A_{j-i}^! \otimes_k M_i$, which occurs as the degree (i, j) component of $F(M)$. Moreover, the differentials in both complexes coincide, hence we are done. \square

3.3 Descending to triangulated categories

Theorem 3.6 shows that the functors F and G preserve cohomology, so it is reasonable to ask whether they descend to the homotopy and derived categories which are the natural setting for formulating statements about cohomology. We show that the answer is positive for homotopy categories.

Theorem 3.10 ((Eisenbud et al. 2003)). The functors F and G descend to adjoint functors

$$\mathbf{K}(A\text{-grMod}) \xrightleftharpoons[\bar{F}]{\bar{G}} \mathbf{K}(A^!\text{-grMod})$$

between the triangulated homotopy categories of chain complexes.

Proof. An easy explicit check shows that the functors F and G take cones to cones. Thus we are done by Lemma 1.24.

The adjunction between \bar{F} and \bar{G} follows immediately from the adjunction in Theorem 3.2 which identifies subgroups of nullhomotopic morphisms. \square

The adjunction is not an equivalence of homotopy categories– for instance, consider the graded $k[x]$ -module $M = k[x]/(x^2)$. We have shown $G(F(M)) \rightarrow M \rightarrow 0$ is a free resolution, but clearly there is no map $M \rightarrow G(F(M))$ which is a homotopy inverse to this (any such map would have to send x to 0.)

One hopes that inverting quasi-isomorphisms (i.e. passing to the derived category) gives an equivalence. However, the two functors don't descend to the derived categories, as shown by multiple examples throughout the literature– for instance (Keller 2003) argues that the complex $A \in \mathbf{D}(A\text{-grMod})$ is a compact object (i.e. the functor $\text{Hom}_{\mathbf{D}(A\text{-grMod})}(A, -)$ commutes with infinite direct sums) but the object $F(A) \cong k\langle n+1 \rangle[-n-1] \in \mathbf{D}(A^1\text{-grMod})$ is not compact. Below we exhibit explicitly the failure of our functors to descend to the derived category.

Example 3.11 (G does not preserve quasi-isomorphisms). Let $n = 0$, so that $A = k[x]$ and $A^1 = k[\xi]/(\xi^2)$. Consider the complex of graded A^1 -modules

$$\cdots \rightarrow A^1\langle 2 \rangle \xrightarrow{\xi} A^1\langle 1 \rangle \xrightarrow{\xi} A^1 \xrightarrow{\xi} A^1\langle -1 \rangle \xrightarrow{\xi} A^1\langle -2 \rangle \rightarrow \cdots$$

which is exact hence isomorphic to the zero complex in $\mathbf{D}(A^1\text{-grMod})$. The functor G maps this to

$$\cdots \rightarrow 0 \rightarrow \bigoplus_q A\langle -q \rangle \xrightarrow{1+x} \bigoplus_q A\langle -q \rangle \rightarrow 0 \rightarrow \cdots,$$

which is not acyclic (the only non-zero differential is not surjective), hence non-zero in $\mathbf{D}(A\text{-grMod})$.

Bernstein-Gel'fand-Gel'fand equivalence. To work around this apparent problem, we restrict to the full subcategory of bounded complexes and use Proposition 1.25. In this case, a simple spectral sequence argument shows F and G preserve acyclicity. This gives well-defined functors between the bounded derived categories which form an adjoint equivalence– the so-called ‘BGG correspondence’.

Lemma 3.12. If \mathbf{M} is a bounded acyclic complex of finitely generated A -modules, then the complex $F(\mathbf{M})$ is acyclic. Likewise, if \mathbf{N} is a bounded acyclic complex of finitely generated A^1 -modules, then the complex $G(\mathbf{N})$ is acyclic.

Proof. Given such an \mathbf{M} , the double complex (5) has exact columns. Then the first page of the spectral sequence (starting with vertical cohomology) vanishes everywhere. Since $M_p^p = 0$ for large p , the double complex is bounded and the convergence theorem holds, indicating the total complex is acyclic.

The argument for G is similar. □

Thus by Proposition 1.25, F and G descend to functors between derived categories

$$F_D : \mathbf{D}^b(A\text{-grMod}) \rightarrow \mathbf{D}(A^1\text{-grMod}), \quad G_D : \mathbf{D}^b(A^1\text{-grMod}) \rightarrow \mathbf{D}(A\text{-grMod}).$$

To conclude, we show that F_D in fact has image $\mathbf{D}^b(A^1\text{-grMod})$ and likewise for G_D .

Lemma 3.13. If \mathbf{M} is a bounded complex of finitely generated A -modules, then the complex $F(\mathbf{M})$ has bounded cohomology and is quasi-isomorphic to a bounded complex of finitely generated A^1 -modules.

Likewise, if \mathbf{N} is a bounded complex of finitely generated $A^!$ -modules, then the complex $G(\mathbf{N})$ has bounded cohomology and is quasi-isomorphic to a bounded complex of finitely generated $A^!$ -modules.

Proof. If \mathbf{N} is as given, then the complex $G(\mathbf{N})$ is bounded by definition— for any p , we have that the module N_{\bullet}^p is finitely generated hence has only finitely many graded components. Then for sufficiently large i , we have $N_{p-i}^p = 0$ for all p .

For \mathbf{M} as given the double complex (5) which computes M is bounded, and by Corollary 3.9 the first page of the corresponding spectral sequence (starting with horizontal cohomology) has finite support. Thus by the convergence theorem for spectral sequences, the cohomology of $F(\mathbf{M})$ is bounded. The existence of the quasi-isomorphic bounded complex of finitely generated modules follows from Hartshorne (2008), III Lemma 12.3. \square

Theorem 3.14 (Bernstein et al. (1978)). The functors F and G induce an equivalence of derived categories

$$\mathbf{D}^b(A\text{-grMod}) \xrightleftharpoons[F_D]{G_D} \mathbf{D}^b(A^!\text{-grMod}).$$

Proof. From Lemma 3.13, the functors given are well-defined. Then Theorem 3.6 shows that $F_D \circ G_D$ and $G_D \circ F_D$ are naturally equivalent to the identity morphism, hence we have an equivalence of categories. \square

3.4 The Tate resolution and Beilinson monads

Eisenbud et al. (2003) exhibits a connection between Beilinson’s theorem and the work of Bernstein-Gel’fand-Gel’fand, showing that the BGG functors can be used to extract the resolution of any sheaf in terms of the generators of $\mathbf{D}^b(\mathbb{P}^n)$ given in Theorem 2.19. We state the result, and look at some computations.

A sheaf $\mathcal{A} \in \mathcal{Coh} \mathbb{P}^n$ corresponds to a finitely generated graded A -module M , for example $M = \bigoplus_{i \geq 0} H^0(\mathcal{A}(i))$. Any two such modules must agree in large degrees. Since finitely generated modules have finite regularity, by Corollary 3.9 there is an $m \geq 0$ such that the complex $F(M)$ is exact in degrees $> m$. Taking the minimal free resolution of $\ker(F(M)^m \rightarrow F(M)^{m+1})$, we obtain an exact complex

$$\mathbf{T}(\mathcal{A}) : \dots \rightarrow T^{m-2} \rightarrow T^{m-1} \rightarrow F(M)^m \rightarrow F(M)^{m+1} \rightarrow \dots$$

called the *Tate resolution* of \mathcal{A} . Eisenbud et al. (2003) argues this is independent of choice of M and m , and computes the modules T^i appearing in the Tate resolution as

$$T^i = A^i \otimes_k \bigoplus_j H^j(\mathcal{A}(i-j))$$

where $H^j(\mathcal{A}(i-j))$ is regarded as a vector space of internal degree $i-j$.

Note each module in $\mathbf{T}(\mathcal{A})$ is a finite direct sum of modules of the form $A^i\langle i \rangle \cong A^!\langle i-n-1 \rangle$, the dimension controlled by the cohomologies of twists of \mathcal{A} . Thus we can adapt the proof of Lemma 2.18 to show

$$\mathrm{Hom}_{A^!\text{-grMod}}(A^i\langle i \rangle, A^j\langle j \rangle) \cong \mathrm{Hom}_{\mathcal{Coh} \mathbb{P}^n}(\Omega^i(i), \Omega^j(j))$$

allowing us to functorially replace each copy of $A^i(i)$ in $T(\mathcal{A})$ by $\Omega^i(i)$, giving a new complex $\Omega(T(\mathcal{A})) \in \mathbf{C}(\mathcal{Coh} \mathbb{P}^n)$. Note $\Omega^i(i) = 0$ for $i < 0$ or $i > n$ so the complex is bounded. Then Eisenbud et al. (2003) show that $\Omega(T(\mathcal{A}))$ is quasi-isomorphic to \mathcal{A} . In fact, it is precisely the Beilinson monad for \mathcal{A} which we computed in Proposition 2.17.

Theorem 3.15. The complex $\Omega(T(\mathcal{A}))$ is the Beilinson monad for the coherent sheaf \mathcal{A} .

Proof. Observe that the underlying vector spaces do agree with those computed in Proposition 2.17. A full proof of the result is beyond the scope of this exposition, see Section 6 of Eisenbud et al. (2003). \square

This gives an algorithm to compute sheaf cohomology, see for Appendix A of Decker (2006).

3.5 Koszul duality

The work of Bernstein et al. (1978) is the first of a series of equivalences of increasing generalities. Say A is a *Koszul algebra* if the trivial module $k \in A\text{-grMod}$ has a projective resolution

$$\dots \rightarrow P^2 \rightarrow P^1 \rightarrow P^0 \rightarrow k \rightarrow 0$$

such that P^i is generated in degree i . Then the algebra $A^! = \text{Ext}_A^\bullet(k, k)$ is called its *Koszul dual*, and Beilinson, Ginzburg & Soergel (1996) shows that the bounded derived categories of A and $A^!$ are equivalent whenever the graded components of A are finite dimensional. The Koszul complex (9) shows that the polynomial algebra and exterior algebra satisfy the hypotheses, and the proof in the general case is similar to the one for the BGG correspondence.

Keller (2003) argues that this formulation of Koszul duality is restrictive for three reasons—

1. A is a Koszul algebra.
2. A is a graded algebra with finite dimensional components.
3. There is an equivalence only between subcategories of the derived category.

Fløystad (2005) provides a framework in which the algebra A is no longer required to be graded, but the equivalence lies between certain quotients of the homotopy categories that are strictly bigger than the derived categories. This approach is perfected in Keller (2003), where the duality is phrased in terms of differentially graded algebras and coalgebras (which are chain complexes with multiplication and comultiplication respectively given by chain-maps) and corresponding categories of differentially graded modules and comodules. This has the effect of allowing a broader class of algebras into the framework. In the symmetric-exterior case, this simply corresponds to considering A as being concentrated in a single differential degree, and considering the coalgebra $A^!$ as a chain complex with graded components lying in successive differential degrees, the differentials being zero. Then the derived category of differentially graded A -modules is quasi-isomorphic to the *coderived category* of differentially graded $A^!$ -comodules, where the coderived category is defined by localising a class of morphisms smaller than that of quasi-isomorphisms. This is closely related to the Bar and Cobar constructions in algebraic topology.

References

- Beilinson, A. A. (1978), 'Coherent sheaves on \mathbb{P}^n and problems of linear algebra', *Functional Analysis and Its Applications* **12**(3), 214–216.
URL: <http://link.springer.com/10.1007/BF01681436>
- Beilinson, A., Ginzburg, V. & Soergel, W. (1996), 'Koszul Duality Patterns in Representation Theory', *Journal of the American Mathematical Society* **9**(2), 473–527.
URL: <https://www.ams.org/jams/1996-9-02/S0894-0347-96-00192-0/>
- Belmans, P. (2015), 'Lemma 2 from beilinson's "coherent sheaves on \mathbb{P}^n and problems of linear algebra", intuition?', MathOverflow. (<https://mathoverflow.net/users/6263/pbelmans>).
URL: <https://mathoverflow.net/q/223418>
- Bernstein, I. N., Gel'fand, I. M. & Gel'fand, S. I. (1978), 'Algebraic bundles over \mathbb{P}^n and problems of linear algebra', *Functional Analysis and Its Applications* **12**(3), 212–214.
URL: <http://link.springer.com/10.1007/BF01681435>
- Bourbaki, N. (1989), *Algebra*, Elements of mathematics, reprint edn, Springer, Berlin.
- Caldararu, A. (2005), 'Derived categories of sheaves: a skimming', *arXiv:math/0501094* . arXiv: math/0501094.
URL: <http://arxiv.org/abs/math/0501094>
- Carbone, R. (2016), 'The resolution of the diagonal and the derived category on some homogeneous spaces'.
URL: <https://www.math.u-bordeaux.fr/~ybilu/algant/documents/theses/CARBONE.pdf>
- Decker, W. (2006), *Computing in algebraic geometry: a quick start using SINGULAR*, number v. 16 in 'Algorithms and computation in mathematics', Springer ; Hindustan Book Agency, Berlin ; New York : New Delhi.
- Eisenbud, D. (1995), *Commutative algebra with a view toward algebraic geometry*, number 150 in 'Graduate texts in mathematics', Springer-Verlag, New York.
- Eisenbud, D., Floystad, G. & Schreyer, F.-O. (2003), 'Sheaf Cohomology and Free Resolutions over Exterior Algebras', *Transactions of the American Mathematical Society* **355**(11), 4397–4426. arXiv: math/0104203.
URL: <http://arxiv.org/abs/math/0104203>
- Eisenbud, D. & Goto, S. (1984), 'Linear free resolutions and minimal multiplicity', *Journal of Algebra* **88**(1), 89–133.
URL: <https://linkinghub.elsevier.com/retrieve/pii/0021869384900929>
- Fløystad, G. (2005), 'Koszul duality and equivalences of categories', *Transactions of the American Mathematical Society* **358**(6), 2373–2398.
URL: <http://www.ams.org/tran/2006-358-06/S0002-9947-05-04035-3/>
- Hartshorne, R. (2008), *Algebraic geometry*, number 52 in 'Graduate texts in mathematics', 14 edn, Springer, New York, NY.
- Huybrechts, D. (2006), *Fourier-Mukai transforms in algebraic geometry*, Oxford mathematical monographs, Clarendon, Oxford ; New York. OCLC: ocm64097295.

- Keller, B. (2003), ‘Koszul Duality and Coderived Categories (After K. Lefevre)’.
- Lam, T. Y. (1999), *Lectures on modules and rings*, number 189 in ‘Graduate texts in mathematics’, Springer, New York.
- Loday, J. L. (2012), *Algebraic operads*, number 346 in ‘Grundlehren der mathematischen wissenschaften’, Springer, New York.
- The Stacks project authors (2022), ‘The stacks project’.
URL: <https://stacks.math.columbia.edu>
- Thomas, R. P. (2001), ‘Derived categories for the working mathematician’, *arXiv:math/0001045* .
 arXiv: math/0001045.
URL: <http://arxiv.org/abs/math/0001045>
- Weibel, C. A. (2003), *An introduction to homological algebra*, number 38 in ‘Cambridge studies in advanced mathematics’, reprint. 1997, transf. to digital print edn, Cambridge Univ. Press, Cambridge.