

3.3 Derivatives of Trigonometric Functions

1. $f(x) = x^2 \sin x \xRightarrow{\text{PR}} f'(x) = x^2 \cos x + (\sin x)(2x) = x^2 \cos x + 2x \sin x$
2. $f(x) = x \cos x + 2 \tan x \Rightarrow f'(x) = x(-\sin x) + (\cos x)(1) + 2 \sec^2 x = \cos x - x \sin x + 2 \sec^2 x$
3. $f(x) = e^x \cos x \Rightarrow f'(x) = e^x(-\sin x) + (\cos x)e^x = e^x(\cos x - \sin x)$
4. $y = 2 \sec x - \csc x \Rightarrow y' = 2(\sec x \tan x) - (-\csc x \cot x) = 2 \sec x \tan x + \csc x \cot x$
5. $y = \sec \theta \tan \theta \Rightarrow y' = \sec \theta (\sec^2 \theta) + \tan \theta (\sec \theta \tan \theta) = \sec \theta (\sec^2 \theta + \tan^2 \theta)$. Using the identity $1 + \tan^2 \theta = \sec^2 \theta$, we can write alternative forms of the answer as $\sec \theta (1 + 2 \tan^2 \theta)$ or $\sec \theta (2 \sec^2 \theta - 1)$.
6. $g(\theta) = e^\theta (\tan \theta - \theta) \Rightarrow g'(\theta) = e^\theta (\sec^2 \theta - 1) + (\tan \theta - \theta)e^\theta = e^\theta (\sec^2 \theta - 1 + \tan \theta - \theta)$

$$7. y = c \cos t + t^2 \sin t \Rightarrow y' = c(-\sin t) + t^2(\cos t) + \sin t(2t) = -c \sin t + t(t \cos t + 2 \sin t)$$

$$8. f(t) = \frac{\cot t}{e^t} \Rightarrow f'(t) = \frac{e^t(-\csc^2 t) - (\cot t)e^t}{(e^t)^2} = \frac{e^t(-\csc^2 t - \cot t)}{(e^t)^2} = -\frac{\csc^2 t + \cot t}{e^t}$$

$$9. y = \frac{x}{2 - \tan x} \Rightarrow y' = \frac{(2 - \tan x)(1) - x(-\sec^2 x)}{(2 - \tan x)^2} = \frac{2 - \tan x + x \sec^2 x}{(2 - \tan x)^2}$$

$$10. y = \sin \theta \cos \theta \Rightarrow y' = \sin \theta(-\sin \theta) + \cos \theta(\cos \theta) = \cos^2 \theta - \sin^2 \theta \quad [\text{or } \cos 2\theta]$$

$$11. f(\theta) = \frac{\sin \theta}{1 + \cos \theta} \Rightarrow$$

$$f'(\theta) = \frac{(1 + \cos \theta) \cos \theta - (\sin \theta)(-\sin \theta)}{(1 + \cos \theta)^2} = \frac{\cos \theta + \cos^2 \theta + \sin^2 \theta}{(1 + \cos \theta)^2} = \frac{\cos \theta + 1}{(1 + \cos \theta)^2} = \frac{1}{1 + \cos \theta}$$

$$12. y = \frac{\cos x}{1 - \sin x} \Rightarrow$$

$$y' = \frac{(1 - \sin x)(-\sin x) - \cos x(-\cos x)}{(1 - \sin x)^2} = \frac{-\sin x + \sin^2 x + \cos^2 x}{(1 - \sin x)^2} = \frac{-\sin x + 1}{(1 - \sin x)^2} = \frac{1}{1 - \sin x}$$

$$13. y = \frac{t \sin t}{1 + t} \Rightarrow$$

$$y' = \frac{(1 + t)(t \cos t + \sin t) - t \sin t(1)}{(1 + t)^2} = \frac{t \cos t + \sin t + t^2 \cos t + t \sin t - t \sin t}{(1 + t)^2} = \frac{(t^2 + t) \cos t + \sin t}{(1 + t)^2}$$

$$14. y = \frac{\sin t}{1 + \tan t} \Rightarrow$$

$$y' = \frac{(1 + \tan t) \cos t - (\sin t) \sec^2 t}{(1 + \tan t)^2} = \frac{\cos t + \sin t - \frac{\sin t}{\cos^2 t}}{(1 + \tan t)^2} = \frac{\cos t + \sin t - \tan t \sec t}{(1 + \tan t)^2}$$

$$15. \text{ Using Exercise 3.2.61(a) , } f(\theta) = \theta \cos \theta \sin \theta \Rightarrow$$

$$\begin{aligned} f'(\theta) &= 1 \cos \theta \sin \theta + \theta(-\sin \theta) \sin \theta + \theta \cos \theta(\cos \theta) = \cos \theta \sin \theta - \theta \sin^2 \theta + \theta \cos^2 \theta \\ &= \sin \theta \cos \theta + \theta(\cos^2 \theta - \sin^2 \theta) = \frac{1}{2} \sin 2\theta + \theta \cos 2\theta \quad [\text{using double-angle formulas}] \end{aligned}$$

$$16. \text{ Using Exercise 3.2.61(a) , } f(t) = te^t \cot t \Rightarrow$$

$$f'(t) = 1e^t \cot t + te^t \cot t + te^t(-\csc^2 t) = e^t(\cot t + t \cot t - t \csc^2 t)$$

$$17. \frac{d}{dx}(\csc x) = \frac{d}{dx} \left(\frac{1}{\sin x} \right) = \frac{(\sin x)(0) - 1(\cos x)}{\sin^2 x} = \frac{-\cos x}{\sin^2 x} = -\frac{1}{\sin x} \cdot \frac{\cos x}{\sin x} = -\csc x \cot x$$

$$18. \frac{d}{dx}(\sec x) = \frac{d}{dx} \left(\frac{1}{\cos x} \right) = \frac{(\cos x)(0) - 1(-\sin x)}{\cos^2 x} = \frac{\sin x}{\cos^2 x} = \frac{1}{\cos x} \cdot \frac{\sin x}{\cos x} = \sec x \tan x$$

$$19. \frac{d}{dx}(\cot x) = \frac{d}{dx} \left(\frac{\cos x}{\sin x} \right) = \frac{(\sin x)(-\sin x) - (\cos x)(\cos x)}{\sin^2 x} = -\frac{\sin^2 x + \cos^2 x}{\sin^2 x} = -\frac{1}{\sin^2 x} = -\csc^2 x$$

20. $f(x) = \cos x \Rightarrow$

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{\cos(x+h) - \cos x}{h} = \lim_{h \rightarrow 0} \frac{\cos x \cos h - \sin x \sin h - \cos x}{h} \\ &= \lim_{h \rightarrow 0} \left(\cos x \frac{\cos h - 1}{h} - \sin x \frac{\sin h}{h} \right) = \cos x \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} - \sin x \lim_{h \rightarrow 0} \frac{\sin h}{h} \\ &= (\cos x)(0) - (\sin x)(1) = -\sin x \end{aligned}$$


21. $y = \sin x + \cos x \Rightarrow y' = \cos x - \sin x$, so $y'(0) = \cos 0 - \sin 0 = 1 - 0 = 1$. An equation of the tangent line to the curve $y = \sin x + \cos x$ at the point $(0, 1)$ is $y - 1 = 1(x - 0)$ or $y = x + 1$.

22. $y = e^x \cos x \Rightarrow y' = e^x(-\sin x) + (\cos x)e^x = e^x(\cos x - \sin x) \Rightarrow$ the slope of the tangent line at $(0, 1)$ is $e^0(\cos 0 - \sin 0) = 1(1 - 0) = 1$ and an equation is $y - 1 = 1(x - 0)$ or $y = x + 1$.


23. $y = \cos x - \sin x \Rightarrow y' = -\sin x - \cos x$, so $y'(\pi) = -\sin \pi - \cos \pi = 0 - (-1) = 1$. An equation of the tangent line to the curve $y = \cos x - \sin x$ at the point $(\pi, -1)$ is $y - (-1) = 1(x - \pi)$ or $y = x - \pi - 1$.

24. $y = x + \tan x \Rightarrow y' = 1 + \sec^2 x$, so $y'(\pi) = 1 + (-1)^2 = 2$. An equation of the tangent line to the curve $y = x + \tan x$ at the point (π, π) is $y - \pi = 2(x - \pi)$ or $y = 2x - \pi$.


25. (a) $y = 2x \sin x \Rightarrow y' = 2(x \cos x + \sin x \cdot 1)$. At $(\frac{\pi}{2}, \pi)$,
 $y' = 2(\frac{\pi}{2} \cos \frac{\pi}{2} + \sin \frac{\pi}{2}) = 2(0 + 1) = 2$, and an equation of the tangent line is $y - \pi = 2(x - \frac{\pi}{2})$, or $y = 2x$.



26. (a) $y = 3x + 6 \cos x \Rightarrow y' = 3 - 6 \sin x$. At $(\frac{\pi}{3}, \pi + 3)$,
 $y' = 3 - 6 \sin \frac{\pi}{3} = 3 - 6 \frac{\sqrt{3}}{2} = 3 - 3\sqrt{3}$, and an equation of the tangent line is $y - (\pi + 3) = (3 - 3\sqrt{3})(x - \frac{\pi}{3})$, or
 $y = (3 - 3\sqrt{3})x + 3 + \pi\sqrt{3}$.




27. (a) $f(x) = \sec x - x \Rightarrow f'(x) = \sec x \tan x - 1$



Note that $f' = 0$ where f has a minimum. Also note that f' is negative when f is decreasing and f' is positive when f is increasing.

28. (a) $f(x) = e^x \cos x \Rightarrow f'(x) = e^x(-\sin x) + (\cos x)e^x = e^x(\cos x - \sin x) \Rightarrow$

$$f''(x) = e^x(-\sin x - \cos x) + (\cos x - \sin x)e^x = e^x(-\sin x - \cos x + \cos x - \sin x) = -2e^x \sin x$$



Note that $f' = 0$ where f has a minimum and $f'' = 0$ where f' has a minimum. Also note that f' is negative when f is decreasing and f'' is negative when f' is decreasing.

29. $H(\theta) = \theta \sin \theta \Rightarrow H'(\theta) = \theta (\cos \theta) + (\sin \theta) \cdot 1 = \theta \cos \theta + \sin \theta \Rightarrow$

$H''(\theta) = \theta (-\sin \theta) + (\cos \theta) \cdot 1 + \cos \theta = -\theta \sin \theta + 2 \cos \theta$

30. $f(t) = \sec t \Rightarrow f'(t) = \sec t \tan t \Rightarrow f''(t) = (\sec t) \sec^2 t + (\tan t) \sec t \tan t = \sec^3 t + \sec t \tan^2 t$, so

$f''\left(\frac{\pi}{4}\right) = (\sqrt{2})^3 + \sqrt{2}(1)^2 = 2\sqrt{2} + \sqrt{2} = 3\sqrt{2}$.

31. (a) $f(x) = \frac{\tan x - 1}{\sec x} \Rightarrow$

$f'(x) = \frac{\sec x (\sec^2 x) - (\tan x - 1)(\sec x \tan x)}{(\sec x)^2} = \frac{\sec x (\sec^2 x - \tan^2 x + \tan x)}{\sec^2 x} = \frac{1 + \tan x}{\sec x}$

(b) $f(x) = \frac{\tan x - 1}{\sec x} = \frac{\frac{\sin x}{\cos x} - 1}{\frac{1}{\cos x}} = \frac{\sin x - \cos x}{\frac{1}{\cos x}} = \sin x - \cos x \Rightarrow f'(x) = \cos x - (-\sin x) = \cos x + \sin x$

(c) From part (a), $f'(x) = \frac{1 + \tan x}{\sec x} = \frac{1}{\sec x} + \frac{\tan x}{\sec x} = \cos x + \sin x$, which is the expression for $f'(x)$ in part (b).

32. (a) $g(x) = f(x) \sin x \Rightarrow g'(x) = f(x) \cos x + \sin x \cdot f'(x)$, so

$g'\left(\frac{\pi}{3}\right) = f\left(\frac{\pi}{3}\right) \cos \frac{\pi}{3} + \sin \frac{\pi}{3} \cdot f'\left(\frac{\pi}{3}\right) = 4 \cdot \frac{1}{2} + \frac{\sqrt{3}}{2} \cdot (-2) = 2 - \sqrt{3}$

(b) $h(x) = \frac{\cos x}{f(x)} \Rightarrow h'(x) = \frac{f(x) \cdot (-\sin x) - \cos x \cdot f'(x)}{[f(x)]^2}$, so

$h'\left(\frac{\pi}{3}\right) = \frac{f\left(\frac{\pi}{3}\right) \cdot (-\sin \frac{\pi}{3}) - \cos \frac{\pi}{3} \cdot f'\left(\frac{\pi}{3}\right)}{[f\left(\frac{\pi}{3}\right)]^2} = \frac{4\left(-\frac{\sqrt{3}}{2}\right) - \left(\frac{1}{2}\right)(-2)}{4^2} = \frac{-2\sqrt{3} + 1}{16} = \frac{1 - 2\sqrt{3}}{16}$

33. $f(x) = x + 2 \sin x$ has a horizontal tangent when $f'(x) = 0 \Leftrightarrow 1 + 2 \cos x = 0 \Leftrightarrow \cos x = -\frac{1}{2} \Leftrightarrow$

$x = \frac{2\pi}{3} + 2\pi n$ or $\frac{4\pi}{3} + 2\pi n$, where n is an integer. Note that $\frac{4\pi}{3}$ and $\frac{2\pi}{3}$ are $\pm \frac{\pi}{3}$ units from π . This allows us to write the solutions in the more compact equivalent form $(2n + 1)\pi \pm \frac{\pi}{3}$, n an integer.

34. $f(x) = e^x \cos x$ has a horizontal tangent when $f'(x) = 0$. $f'(x) = e^x(-\sin x) + (\cos x)e^x = e^x(\cos x - \sin x)$.

$f'(x) = 0 \Leftrightarrow \cos x - \sin x = 0 \Leftrightarrow \cos x = \sin x \Leftrightarrow \tan x = 1 \Leftrightarrow x = \frac{\pi}{4} + n\pi$, n an integer.

35. (a) $x(t) = 8 \sin t \Rightarrow v(t) = x'(t) = 8 \cos t \Rightarrow a(t) = x''(t) = -8 \sin t$

(b) The mass at time $t = \frac{2\pi}{3}$ has position $x\left(\frac{2\pi}{3}\right) = 8 \sin \frac{2\pi}{3} = 8\left(\frac{\sqrt{3}}{2}\right) = 4\sqrt{3}$, velocity $v\left(\frac{2\pi}{3}\right) = 8 \cos \frac{2\pi}{3} = 8\left(-\frac{1}{2}\right) = -4$,


and acceleration $a\left(\frac{2\pi}{3}\right) = -8 \sin \frac{2\pi}{3} = -8\left(\frac{\sqrt{3}}{2}\right) = -4\sqrt{3}$. Since $v\left(\frac{2\pi}{3}\right) < 0$, the particle is moving to the left.

36. (a) $s(t) = 2 \cos t + 3 \sin t \Rightarrow v(t) = -2 \sin t + 3 \cos t \Rightarrow$
 $a(t) = -2 \cos t - 3 \sin t$


(c) $s = 0 \Rightarrow t_2 \approx 2.55$. So the mass passes through the equilibrium position for the first time when $t \approx 2.55$ s.

(d) $v = 0 \Rightarrow t_1 \approx 0.98, s(t_1) \approx 3.61$ cm. So the mass travels a maximum of about 3.6 cm (upward and downward) from its equilibrium position.

(e) The speed $|v|$ is greatest when $s = 0$, that is, when $t = t_2 + n\pi$, n a positive integer.



37.




From the diagram we can see that $\sin \theta = x/10 \Leftrightarrow x = 10 \sin \theta$. We want to find the rate of change of x with respect to θ , that is, $dx/d\theta$. Taking the derivative of $x = 10 \sin \theta$, we get $dx/d\theta = 10(\cos \theta)$. So when $\theta = \frac{\pi}{3}$, $\frac{dx}{d\theta} = 10 \cos \frac{\pi}{3} = 10(\frac{1}{2}) = 5$ ft/rad.

38. (a) $F = \frac{\mu W}{\mu \sin \theta + \cos \theta} \Rightarrow \frac{dF}{d\theta} = \frac{(\mu \sin \theta + \cos \theta)(0) - \mu W(\mu \cos \theta - \sin \theta)}{(\mu \sin \theta + \cos \theta)^2} = \frac{\mu W(\sin \theta - \mu \cos \theta)}{(\mu \sin \theta + \cos \theta)^2}$

(b) $\frac{dF}{d\theta} = 0 \Leftrightarrow \mu W(\sin \theta - \mu \cos \theta) = 0 \Leftrightarrow \sin \theta = \mu \cos \theta \Leftrightarrow \tan \theta = \mu \Leftrightarrow \theta = \tan^{-1} \mu$

(c)



From the graph of $F = \frac{0.6(50)}{0.6 \sin \theta + \cos \theta}$ for $0 \leq \theta \leq 1$, we see that

$\frac{dF}{d\theta} = 0 \Rightarrow \theta \approx 0.54$. Checking this with part (b) and $\mu = 0.6$, we

calculate $\theta = \tan^{-1} 0.6 \approx 0.54$. So the value from the graph is consistent with the value in part (b).

39. $\lim_{x \rightarrow 0} \frac{\sin 5x}{3x} = \lim_{x \rightarrow 0} \frac{5}{3} \left(\frac{\sin 5x}{5x} \right) = \frac{5}{3} \lim_{x \rightarrow 0} \frac{\sin 5x}{5x} = \frac{5}{3} \lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} \quad [\theta = 5x] = \frac{5}{3} \cdot 1 = \frac{5}{3}$

40. $\lim_{x \rightarrow 0} \frac{\sin x}{\sin \pi x} = \lim_{x \rightarrow 0} \frac{\sin x}{x} \cdot \frac{\pi x}{\sin \pi x} \cdot \frac{1}{\pi} = \lim_{x \rightarrow 0} \frac{\sin x}{x} \cdot \lim_{\theta \rightarrow 0} \frac{\theta}{\sin \theta} \cdot \frac{1}{\pi} \quad [\theta = \pi x]$
 $= 1 \cdot \lim_{\theta \rightarrow 0} \frac{1}{\frac{\sin \theta}{\theta}} \cdot \frac{1}{\pi} = 1 \cdot 1 \cdot \frac{1}{\pi} = \frac{1}{\pi}$

41. $\lim_{t \rightarrow 0} \frac{\tan 6t}{\sin 2t} = \lim_{t \rightarrow 0} \left(\frac{\sin 6t}{t} \cdot \frac{1}{\cos 6t} \cdot \frac{t}{\sin 2t} \right) = \lim_{t \rightarrow 0} \frac{6 \sin 6t}{6t} \cdot \lim_{t \rightarrow 0} \frac{1}{\cos 6t} \cdot \lim_{t \rightarrow 0} \frac{2t}{2 \sin 2t}$
 $= 6 \lim_{t \rightarrow 0} \frac{\sin 6t}{6t} \cdot \lim_{t \rightarrow 0} \frac{1}{\cos 6t} \cdot \frac{1}{2} \lim_{t \rightarrow 0} \frac{2t}{\sin 2t} = 6(1) \cdot \frac{1}{1} \cdot \frac{1}{2}(1) = 3$

42. $\lim_{\theta \rightarrow 0} \frac{\cos \theta - 1}{\sin \theta} = \lim_{\theta \rightarrow 0} \frac{\frac{\cos \theta - 1}{\theta}}{\frac{\sin \theta}{\theta}} = \frac{\lim_{\theta \rightarrow 0} \frac{\cos \theta - 1}{\theta}}{\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta}} = \frac{0}{1} = 0$

$$43. \lim_{x \rightarrow 0} \frac{\sin 3x}{5x^3 - 4x} = \lim_{x \rightarrow 0} \left(\frac{\sin 3x}{3x} \cdot \frac{3}{5x^2 - 4} \right) = \lim_{x \rightarrow 0} \frac{\sin 3x}{3x} \cdot \lim_{x \rightarrow 0} \frac{3}{5x^2 - 4} = 1 \cdot \left(\frac{3}{-4} \right) = -\frac{3}{4}$$

$$44. \lim_{x \rightarrow 0} \frac{\sin 3x \sin 5x}{x^2} = \lim_{x \rightarrow 0} \left(\frac{\sin 3x}{3x} \cdot \frac{5 \sin 5x}{5x} \right) = \lim_{x \rightarrow 0} \frac{\sin 3x}{3x} \cdot \lim_{x \rightarrow 0} \frac{5 \sin 5x}{5x} \\ = 3 \lim_{x \rightarrow 0} \frac{\sin 3x}{3x} \cdot 5 \lim_{x \rightarrow 0} \frac{\sin 5x}{5x} = 3(1) \cdot 5(1) = 15$$

45. Divide numerator and denominator by θ . ($\sin \theta$ also works.)

$$\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta + \tan \theta} = \lim_{\theta \rightarrow 0} \frac{\frac{\sin \theta}{\theta}}{1 + \frac{\sin \theta}{\theta} \cdot \frac{1}{\cos \theta}} = \frac{\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta}}{1 + \lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} \lim_{\theta \rightarrow 0} \frac{1}{\cos \theta}} = \frac{1}{1 + 1 \cdot 1} = \frac{1}{2}$$

$$46. \lim_{x \rightarrow 0} \csc x \sin(\sin x) = \lim_{x \rightarrow 0} \frac{\sin(\sin x)}{\sin x} = \lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} \quad [\text{As } x \rightarrow 0, \theta = \sin x \rightarrow 0.] = 1$$

$$47. \lim_{\theta \rightarrow 0} \frac{\cos \theta - 1}{2\theta^2} = \lim_{\theta \rightarrow 0} \frac{\cos \theta - 1}{2\theta^2} \cdot \frac{\cos \theta + 1}{\cos \theta + 1} = \lim_{\theta \rightarrow 0} \frac{\cos^2 \theta - 1}{2\theta^2(\cos \theta + 1)} = \lim_{\theta \rightarrow 0} \frac{-\sin^2 \theta}{2\theta^2(\cos \theta + 1)} \\ = -\frac{1}{2} \lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} \cdot \frac{\sin \theta}{\theta} \cdot \frac{1}{\cos \theta + 1} = -\frac{1}{2} \lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} \cdot \lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} \cdot \lim_{\theta \rightarrow 0} \frac{1}{\cos \theta + 1} \\ = -\frac{1}{2} \cdot 1 \cdot 1 \cdot \frac{1}{1 + 1} = -\frac{1}{4}$$

$$48. \lim_{x \rightarrow 0} \frac{\sin(x^2)}{x} = \lim_{x \rightarrow 0} \left[x \cdot \frac{\sin(x^2)}{x \cdot x} \right] = \lim_{x \rightarrow 0} x \cdot \lim_{x \rightarrow 0} \frac{\sin(x^2)}{x^2} = 0 \cdot \lim_{y \rightarrow 0^+} \frac{\sin y}{y} \quad [\text{where } y = x^2] = 0 \cdot 1 = 0$$

$$49. \lim_{x \rightarrow \pi/4} \frac{1 - \tan x}{\sin x - \cos x} = \lim_{x \rightarrow \pi/4} \frac{\left(1 - \frac{\sin x}{\cos x}\right) \cdot \cos x}{(\sin x - \cos x) \cdot \cos x} = \lim_{x \rightarrow \pi/4} \frac{\cos x - \sin x}{(\sin x - \cos x) \cos x} = \lim_{x \rightarrow \pi/4} \frac{-1}{\cos x} = \frac{-1}{1/\sqrt{2}} = -\sqrt{2}$$

$$50. \lim_{x \rightarrow 1} \frac{\sin(x-1)}{x^2 + x - 2} = \lim_{x \rightarrow 1} \frac{\sin(x-1)}{(x+2)(x-1)} = \lim_{x \rightarrow 1} \frac{1}{x+2} \lim_{x \rightarrow 1} \frac{\sin(x-1)}{x-1} = \frac{1}{3} \cdot 1 = \frac{1}{3}$$

$$51. \frac{d}{dx}(\sin x) = \cos x \Rightarrow \frac{d^2}{dx^2}(\sin x) = -\sin x \Rightarrow \frac{d^3}{dx^3}(\sin x) = -\cos x \Rightarrow \frac{d^4}{dx^4}(\sin x) = \sin x.$$

The derivatives of $\sin x$ occur in a cycle of four. Since $99 = 4(24) + 3$, we have $\frac{d^{99}}{dx^{99}}(\sin x) = \frac{d^3}{dx^3}(\sin x) = -\cos x$.

52. Let $f(x) = x \sin x$ and $h(x) = \sin x$, so $f(x) = xh(x)$. Then $f'(x) = h(x) + xh'(x)$,

$$f''(x) = h'(x) + h'(x) + xh''(x) = 2h'(x) + xh''(x),$$

$$f'''(x) = 2h''(x) + h''(x) + xh'''(x) = 3h''(x) + xh'''(x), \dots, f^{(n)}(x) = nh^{(n-1)}(x) + xh^{(n)}(x).$$

Since $34 = 4(8) + 2$, we have $h^{(34)}(x) = h^{(2)}(x) = \frac{d^2}{dx^2}(\sin x) = -\sin x$ and $h^{(35)}(x) = -\cos x$.

$$\text{Thus, } \frac{d^{35}}{dx^{35}}(x \sin x) = 35h^{(34)}(x) + xh^{(35)}(x) = -35 \sin x - x \cos x.$$

53. $y = A \sin x + B \cos x \Rightarrow y' = A \cos x - B \sin x \Rightarrow y'' = -A \sin x - B \cos x$. Substituting these expressions for y , y' , and y'' into the given differential equation $y'' + y' - 2y = \sin x$ gives us

$$(-A \sin x - B \cos x) + (A \cos x - B \sin x) - 2(A \sin x + B \cos x) = \sin x \Leftrightarrow$$

$$-3A \sin x - B \sin x + A \cos x - 3B \cos x = \sin x \Leftrightarrow (-3A - B) \sin x + (A - 3B) \cos x = 1 \sin x, \text{ so we must have}$$

$-3A - B = 1$ and $A - 3B = 0$ (since 0 is the coefficient of $\cos x$ on the right side). Solving for A and B , we add the first



equation to three times the second to get $B = -\frac{1}{10}$ and $A = -\frac{3}{10}$.

54. (a) Let $\theta = \frac{1}{x}$. Then as $x \rightarrow \infty$, $\theta \rightarrow 0^+$, and $\lim_{x \rightarrow \infty} x \sin \frac{1}{x} = \lim_{\theta \rightarrow 0^+} \frac{1}{\theta} \sin \theta = \lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1$.

(b) Since $-1 \leq \sin(1/x) \leq 1$, we have (as illustrated in the figure)

$$-|x| \leq x \sin(1/x) \leq |x|. \text{ We know that } \lim_{x \rightarrow 0} (|x|) = 0 \text{ and}$$

$$\lim_{x \rightarrow 0} (-|x|) = 0; \text{ so by the Squeeze Theorem, } \lim_{x \rightarrow 0} x \sin(1/x) = 0.$$



55. (a) $\frac{d}{dx} \tan x = \frac{d}{dx} \frac{\sin x}{\cos x} \Rightarrow \sec^2 x = \frac{\cos x \cos x - \sin x (-\sin x)}{\cos^2 x} = \frac{\cos^2 x + \sin^2 x}{\cos^2 x}$. So $\sec^2 x = \frac{1}{\cos^2 x}$.

(b) $\frac{d}{dx} \sec x = \frac{d}{dx} \frac{1}{\cos x} \Rightarrow \sec x \tan x = \frac{(\cos x)(0) - 1(-\sin x)}{\cos^2 x}$. So $\sec x \tan x = \frac{\sin x}{\cos^2 x}$.

(c) $\frac{d}{dx} (\sin x + \cos x) = \frac{d}{dx} \frac{1 + \cot x}{\csc x} \Rightarrow$

$$\begin{aligned} \cos x - \sin x &= \frac{\csc x (-\csc^2 x) - (1 + \cot x)(-\csc x \cot x)}{\csc^2 x} = \frac{\csc x [-\csc^2 x + (1 + \cot x) \cot x]}{\csc^2 x} \\ &= \frac{-\csc^2 x + \cot^2 x + \cot x}{\csc x} = \frac{-1 + \cot x}{\csc x} \end{aligned}$$


$$\text{So } \cos x - \sin x = \frac{\cot x - 1}{\csc x}.$$

56. We get the following formulas for r and h in terms of θ :

$$\sin \frac{\theta}{2} = \frac{r}{10} \Rightarrow r = 10 \sin \frac{\theta}{2} \quad \text{and} \quad \cos \frac{\theta}{2} = \frac{h}{10} \Rightarrow h = 10 \cos \frac{\theta}{2}$$

Now $A(\theta) = \frac{1}{2} \pi r^2$ and $B(\theta) = \frac{1}{2} (2r)h = rh$. So

$$\begin{aligned} \lim_{\theta \rightarrow 0^+} \frac{A(\theta)}{B(\theta)} &= \lim_{\theta \rightarrow 0^+} \frac{\frac{1}{2} \pi r^2}{rh} = \frac{1}{2} \pi \lim_{\theta \rightarrow 0^+} \frac{r}{h} = \frac{1}{2} \pi \lim_{\theta \rightarrow 0^+} \frac{10 \sin(\theta/2)}{10 \cos(\theta/2)} \\ &= \frac{1}{2} \pi \lim_{\theta \rightarrow 0^+} \tan(\theta/2) = 0 \end{aligned}$$




57. By the definition of radian measure, $s = r\theta$, where r is the radius of the circle. By drawing the bisector of the angle θ , we can

see that $\sin \frac{\theta}{2} = \frac{d/2}{r} \Rightarrow d = 2r \sin \frac{\theta}{2}$. So $\lim_{\theta \rightarrow 0^+} \frac{s}{d} = \lim_{\theta \rightarrow 0^+} \frac{r\theta}{2r \sin(\theta/2)} = \lim_{\theta \rightarrow 0^+} \frac{2 \cdot (\theta/2)}{2 \sin(\theta/2)} = \lim_{\theta \rightarrow 0} \frac{\theta/2}{\sin(\theta/2)} = 1$.

[This is just the reciprocal of the limit $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$ combined with the fact that as $\theta \rightarrow 0$, $\frac{\theta}{2} \rightarrow 0$ also.]

58. (a)



It appears that $f(x) = \frac{x}{\sqrt{1 - \cos 2x}}$ has a jump discontinuity at $x = 0$.

(b) Using the identity $\cos 2x = 1 - \sin^2 x$, we have $\frac{x}{\sqrt{1 - \cos 2x}} = \frac{x}{\sqrt{1 - (1 - \sin^2 x)}} = \frac{x}{\sqrt{2 \sin^2 x}} = \frac{x}{\sqrt{2} |\sin x|}$.

Thus,

$$\begin{aligned} \lim_{x \rightarrow 0^-} \frac{x}{\sqrt{1 - \cos 2x}} &= \lim_{x \rightarrow 0^-} \frac{x}{\sqrt{2} |\sin x|} = \frac{1}{\sqrt{2}} \lim_{x \rightarrow 0^-} \frac{x}{-(\sin x)} \\ &= -\frac{1}{\sqrt{2}} \lim_{x \rightarrow 0^-} \frac{1}{\sin x / x} = -\frac{1}{\sqrt{2}} \cdot \frac{1}{1} = -\frac{\sqrt{2}}{2} \end{aligned}$$

Evaluating $\lim_{x \rightarrow 0^+} f(x)$ is similar, but $|\sin x| = +\sin x$, so we get $\frac{1}{2}\sqrt{2}$. These values appear to be reasonable values for the graph, so they confirm our answer to part (a).

Another method: Multiply numerator and denominator by $\sqrt{1 + \cos 2x}$.

3.4 The Chain Rule

1. Let $u = g(x) = 1 + 4x$ and $y = f(u) = \sqrt[3]{u}$. Then $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = (\frac{1}{3}u^{-2/3})(4) = \frac{4}{3\sqrt[3]{(1+4x)^2}}$.

2. Let $u = g(x) = 2x^3 + 5$ and $y = f(u) = u^4$. Then $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = (4u^3)(6x^2) = 24x^2(2x^3 + 5)^3$.

3. Let $u = g(x) = \pi x$ and $y = f(u) = \tan u$. Then $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = (\sec^2 u)(\pi) = \pi \sec^2 \pi x$.

4. Let $u = g(x) = \cot x$ and $y = f(u) = \sin u$. Then $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = (\cos u)(-\csc^2 x) = -\cos(\cot x) \csc^2 x$.

5. Let $u = g(x) = \sqrt{x}$ and $y = f(u) = e^u$. Then $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = (e^u)(\frac{1}{2}x^{-1/2}) = e^{\sqrt{x}} \cdot \frac{1}{2\sqrt{x}} = \frac{e^{\sqrt{x}}}{2\sqrt{x}}$.

6. Let $u = g(x) = 2 - e^x$ and $y = f(u) = \sqrt{u}$. Then $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = (\frac{1}{2}u^{-1/2})(-e^x) = -\frac{e^x}{2\sqrt{2 - e^x}}$.

7. $F(x) = (5x^6 + 2x^3)^4 \Rightarrow F'(x) = 4(5x^6 + 2x^3)^3 \cdot \frac{d}{dx}(5x^6 + 2x^3) = 4(5x^6 + 2x^3)^3(30x^5 + 6x^2)$.

We can factor as follows: $4(x^3)^3(5x^3 + 2)^3 6x^2(5x^3 + 1) = 24x^{11}(5x^3 + 2)^3(5x^3 + 1)$