Grain spoilage detection methods

"Provide scientific evidence regarding the correlation between CO2 and VOC (Volatile Organic Compounds) levels and the early stages of grain spoilage. Compare the 'detection lead time' of CO2/VOC sensors versus traditional temperature sensors. Does the chemical signature of spoilage appear before the thermal signature?" *Generated by LeapSpace*

CO2 and VOC Signatures in Early Grain Spoilage: Quick Insights

Key Findings Table

Key finding	Grain / context	What the evidence shows	Citations
CO2 is the strongest supported early chemical marker	Wheat, maize, barley, mixed cereal/oilseed bulks	Elevated intergranular CO2 repeatedly correlates with early biological activity and later spoilage outcomes including germination loss, quality decline, ergosterol increase, mycotoxin risk, and dry-matter loss.	[1] [2] [3] [4] [5] [6] [7]
VOC profiles also track deterioration, but timing is less well quantified	Wheat, maize, rice, barley; fungal and insect damage	VOC classes and marker compounds can distinguish healthy from deteriorating grain, including fungal and insect-related changes, but most studies emphasize discrimination/classification rather than exact time-to-detection.	[8] [9] [10] [11] [12] [13] [14] [15] [16] [17] [18] [19]
Chemical sensing usually outperforms temperature for early warning	Stored grain hotspots, wet pockets, localized deterioration	CO2 and VOC changes arise from respiration/metabolism before enough heat accumulates for a clear bulk-temperature rise; temperature cables can miss localized spoilage unless close to the active zone.	[20] [21] [22] [23] [24] [25] [26] [27] [28]

Key finding	Grain / context	What the evidence shows	Citations
Direct chemical-versus-thermal lead-time data are scarce	Same-bulk comparisons across grains	Most retrieved studies do not report exact hour/day/week differences between first CO2/VOC rise and first measurable temperature increase in the same bulk.	[20] [21] [22] [23] [24] [27] [29] [30] [31]
Best explicit CO2 timing example is still approximate	Wheat at 18% moisture	One review reports CO2 increases before a noticeable temperature rise, with the CO2 increase developing over about 10 days; hotspots can form in as little as 5 days at high moisture.	[20]
Performance depends strongly on storage conditions	Commodity, moisture, temperature, aeration, airflow, placement	Gas production and detectability change with moisture, temperature, airflow, commodity type, and sensor location; a signal strategy that works in wheat may not transfer directly to rapeseed or other systems.	[3] [4] [29] [32] [33] [34] [35] [36] [37]

Direct Answer

Yes—on balance, the chemical signature of spoilage appears before the thermal signature in many practical grain-storage scenarios. Rising intergranular CO2 has the strongest scientific support as an early spoilage indicator, with repeated links to fungal activity, insect activity, grain respiration, and later quality loss across multiple grains [1] [2] [3] [4] [5] [6] [7]. VOC changes also correlate with early deterioration, including fungal and insect damage, and specific markers such as 1-octen-3-ol in Aspergillus-infected wheat and ethyl acetate plus 3-hydroxybutan-2-one in early moldy maize have been proposed [9] [10] [11] [14] [15].

Compared with traditional temperature monitoring, **CO2 sensing is generally reported as earlier or more sensitive**, mainly because biological respiration produces gases before enough heat accumulates to create a measurable bulk-temperature rise, and because temperature probes often miss localized hotspots unless positioned very near them [20] [22] [23] [24] [28]. VOC sensing is promising for the same reason, but the evidence base is less direct on timing and stronger on spoilage-state classification than on lead-time quantification [8] [12] [13] [16] [17] [18].

However, **the exact detection lead time is rarely measured rigorously head-to-head**. The retrieved literature mostly supports the sequence qualitatively rather than with precise same-bulk timing benchmarks in hours or days [22] [23] [24] [27] [29] [30]. The clearest explicit timing statement is that CO2 increased before a noticeable temperature rise in wheat at 18% moisture, with the increase developing over about 10 days [20].

Bottom line: chemical sensing, especially CO2, is scientifically supported as an earlier warning layer than temperature sensing, but the literature summarized here does not yet provide a robust universal numeric lead-time range for CO2/VOC versus temperature [20] [21] [31].

Study Scope

This report synthesizes evidence from stored-grain studies and reviews covering wheat, maize/corn, rice, barley, mixed cereal/oilseed bulks, and some broader stored-product contexts, with emphasis on CO2 monitoring, VOC profiling, electronic noses, and comparisons with conventional temperature-based monitoring [1] [2] [8] [12] [14] [18] [20] [38]. The included evidence spans controlled-storage experiments, pilot-scale silo studies, farm-storage observations, sensor-development studies, and review/mechanistic papers [5] [6] [21] [31] [38].

Assumptions & Limitations

The main limitation is that **few studies directly measured first chemical signal and first thermal signal in the same grain bulk**, so most conclusions about lead time are inferential or approximate rather than benchmarked experimentally [22] [23] [24] [27] [29] [30]. VOC evidence is also less mature than CO2 evidence for temporal onset, and engineering-grade validation metrics such as drift, false alarms, and long-term robustness are often missing [15] [31] [39]. Performance is context-dependent: moisture, temperature, aeration, airflow, commodity type, and placement can all alter detectability [29] [32] [34] [36] [37].

Suggested Further Research

- Conduct same-bulk studies that quantify **CO2, VOC, and temperature lead times** in hours/days under controlled and field conditions [23] [24] [31].
- Standardize **alert thresholds and baseline normalization** for chemical sensing across storage conditions [39] [40].
- Validate VOC onset timing against **microbiology, mycotoxins, germination loss, and grade loss**, not only classification accuracy [16] [17] [18].
- Report engineering metrics including **false positives, false negatives, drift, humidity cross-sensitivity, and seasonal robustness** [31] [38] [39].
- Test **sensor fusion** strategies combining CO2, VOC, temperature, humidity, and airflow modeling for earlier and more specific hotspot detection [21] [36] [37].

Full Report

Introduction

Scope and central question

The core question is whether CO2 and VOC signals correlate with the **earliest stages of grain spoilage**, and whether they provide earlier warning than traditional temperature monitoring [1] [20] [31]. The evidence base is strongest for intergranular CO2 as an early marker of biological activity in stored grain, while VOC profiling offers richer diagnostic information but weaker temporal quantification [1] [2] [8] [12] [14].

The practical significance is straightforward: if chemical signatures emerge before measurable heat buildup, then gas sensing can extend the intervention window before spoilage becomes visible, widespread, or commercially significant [20] [21] [23].

Confidence level: High - The existence of an early-warning role for CO2 is well supported across multiple grain contexts, though the exact timing advantage is less precisely quantified [1] [2] [20].

Foundational Context

Early spoilage in stored grain is fundamentally a **biological and physicochemical process** driven by grain respiration, microbial metabolism, and insect activity [30] [41] [42] [43]. These processes generate **CO2**, consume oxygen, and produce **volatile metabolites**, while also eventually generating heat and moisture [41] [42] [44] [45].

This matters for signal timing. A grain mass does not warm uniformly at the onset of deterioration; localized biological activity can produce a detectable gas signal before enough total heat accumulates to change **bulk temperature** or to reach a distant thermistor [23] [24] [26] [27]. That is why gas sensing and temperature sensing are not interchangeable: CO2/VOC measurements can capture **metabolic onset**, while temperature often captures **later escalation or spatially mature hotspots** [20] [22] [28].

VOCs add a second layer of information. Whereas CO2 is a broad respiration marker, VOC profiles can reflect the **type of spoilage process**—for example fungal contamination, insect infestation, or broader storage-related degradation [9] [11] [14] [46]. Taken together, these findings suggest a functional division: CO2 is useful for early warning, VOCs for discrimination, and temperature for confirmation/localization once deterioration intensifies [19] [21] [26] [46].

Confidence level: High - The biological basis for early gas production is consistent across mechanistic and applied studies [30] [41] [42] [43].

CO2 and VOC as Early Biomarkers of Grain Spoilage

CO2 accumulation as an incipient spoilage signal

Across the retrieved literature, intergranular CO2 is the most consistently supported chemical indicator of early spoilage [1] [2] [4]. In stored wheat, cumulative CO2 production was linked to spoilage variables and the onset of germination loss, indicating that the gas signal rises during deterioration rather than merely after severe damage appears [1]. Farm-stored cereal and oilseed bulks also showed that elevated interseed-air CO2 aligned with spoiled grain in most bins [2].

The mechanism is biologically plausible and repeatedly observed: higher CO2 in wheat reflected fungal and insect activity, with fungi more important in high-moisture wheat and insects more important in safe-moisture wheat [3] [4]. Controlled studies in durum wheat showed CO2 rising substantially over storage while tracking ergosterol and ochratoxin A, linking the gas signal not just to respiration but to fungal colonization and mycotoxin-relevant deterioration [5]. Corn studies likewise found elevated CO2 associated with later germination loss, dry-matter loss, altered composition, and increased mycotoxin contamination [6].

These studies support a strong conclusion: **CO2 is not merely correlated with late spoilage; it is a biologically meaningful early-warning marker that rises during incipient deterioration** [1] [2] [5] [6].

VOC profiles and marker compounds

VOC evidence is also substantial, though differently structured [8] [10] [11] [12] [14]. In stored wheat, VOC profiles changed with storage time and temperature, and some VOC sets were proposed as indicators of storage quality or germination rate [8] [10] [11]. VOC profiling also distinguished healthy wheat from wheat infested with *Rhyzopertha dominica*, supporting its use for insect-related deterioration [9].

In maize, VOC emissions increased early during storage and consolidation and then declined, indicating that volatile output can be dynamic rather than monotonic during deterioration [12] [13]. For rice, mold infestation during paddy storage was associated with characteristic VOCs, suggesting utility for early mildew detection [14].

Specific candidate marker compounds were identified in some studies. The evidence cited here highlights **1-octen-3-ol** in *Aspergillus flavus*-infected wheat and **ethyl acetate** plus **3-hydroxybutan-2-one** as proposed early moldy-stage markers in maize kernels [15]. Broader fungal spoilage work also identified alcohols, ketones, aldehydes, acids, esters, and hydrocarbons associated with deterioration-related headspaces [47] [48] [49] [50].

The main caveat is temporal precision: most VOC studies identify discriminatory compounds or classification performance, but not exact first-detection timing relative to visible mold or temperature rise [15] [16] [17].

Biological interpretation of chemical changes

Microbial respiration, insect metabolism, and grain respiration all contribute to CO2 production in stored grain, and several papers explicitly treat these processes as the basis for early spoilage monitoring [30] [41] [42] [43] [44] [51]. VOCs reflect metabolic by-products of these same processes and may appear before obvious visual or sensory rejection [25] [26] [52].

This supports the central causal logic of the field: **chemical signatures arise at the metabolic stage of deterioration, whereas thermal signatures become useful after activity has intensified enough to create detectable heat accumulation** [26] [27] [33].

Confidence level: High for CO2, Medium for VOCs - CO2 evidence is repeated across grains and endpoints; VOC evidence is strong for association and classification but weaker for exact onset timing [1] [2] [4] [8] [12] [14].

Detection Lead Time: Chemical Signatures Versus Thermal Signatures

Evidence that CO2 can precede temperature rise

The strongest general conclusion from the retrieved evidence is that CO2 monitoring can detect spoilage earlier than temperature monitoring, especially for localized hotspots [20] [23] [24] [28]. One review states that CO2 increases were reported before a noticeable temperature rise in wheat stored at 18% moisture, with the CO2 increase developing over about 10 days [20]. The same source notes that hotspots can form in as little as 5 days at high moisture, underscoring how narrow the intervention window can be [20].

Pilot-scale silo work found that CO2 responded earlier and more sensitively than temperature or relative humidity to a localized wet pocket, even though the excerpt does not provide a precise day-by-day offset [21]. More broadly, several sources state that temperature systems are less sensitive because they only detect spoilage well when the sensor lies close to the active zone [23] [24].

Limits of direct timing evidence

Despite that consistent directional conclusion, the literature gap is important: the retrieved studies rarely provide **same-bulk, side-by-side first-detection timestamps** for CO2, VOC, and temperature [22] [23] [24] [29]. For VOCs in particular, the retrieved work mostly emphasizes pattern recognition or spoilage classification, not lead time against thermal monitoring [15] [16] [17].

So the field supports a sequence—chemical first, thermal later—but usually not a robust universal numerical lead time [20] [27] [30].

Does the chemical signature appear first?

The balance of evidence says **usually yes**, particularly for early, localized, biologically driven spoilage [20] [22] [23] [26] [28]. This is supported by mechanism, pilot evidence, and repeated engineering critiques of thermal monitoring in nonuniform bulks [21] [23] [27].

A careful formulation is warranted: **chemical signatures probably appear before useful thermal signatures**, rather than before any conceivable microscale heat generation [27]. Bulk temperature may lag not because no heat exists, but because the heat is too localized or too weak to be detected by typical temperature cables before gas changes are measurable [23] [24].

Confidence level: High for direction, Low-to-Medium for exact lead-time magnitude - Multiple sources support earlier chemical detection, but numeric lead times are sparse [20] [21] [23] [31].

Sensor Performance, Thresholds, and Validation Against Conventional Monitoring

Calibration, sensitivity, and threshold design

A CO2 sensor-development study reported detection over roughly 380-2400 ppm and tested performance across temperature and relative-humidity ranges, with stronger temperature effects than humidity and no cross-sensitivity to alcohols or ketones in the reported tests [38]. Another study indicated that an instrument capable of sensing 0.1% CO2 could detect spoilage in 80% of deteriorating bulks and recommended placement near the bin center when deterioration location is unknown [39].

Threshold concepts are beginning to emerge. One review described stable grain environments as roughly **400-600 ppm CO2**, with persistent values above **800 ppm** treated as early indicators of increased biological activity [40]. A pilot wet-pocket study linked **<~3000 ppm** with mycotoxins below detection limits and **>~6000 ppm** peaks with regions above legal mycotoxin limits, suggesting that thresholds may be made operationally actionable when tied to quality endpoints [21].

Cross-sensitivity and environmental robustness

The evidence consistently indicates that CO2 and VOC sensing are not plug-and-play across all conditions [35] [36] [37] [38] [53]. Temperature, humidity, and airflow affect gas accumulation and transport [36] [38]. Convection and fan airflow can move CO2 vertically or vent it from bins, meaning poor placement can mask local spoilage even when the chemistry is informative [36] [37].

VOC sensor work also suggests sensitivity and selectivity can be improved by using sensor arrays operated at different temperatures or frequencies, reducing sensitivity to ambient fluctuations [53]. Taken together, these findings imply that **placement and environmental compensation are central to real-world performance**, not secondary engineering details [36] [38].

Validation against microbiology and quality loss

VOC and CO2 evidence is strongest when validated against objective endpoints rather than odor alone. Barley studies explicitly related volatile patterns to CFU, ergosterol, and mycotoxin outcomes, showing that electronic-nose and GC-MS approaches can track biologically meaningful deterioration [16] [17] [18]. Another study showed volatile-production profiles could distinguish mycotoxigenic from non-mycotoxigenic *Fusarium* strains, strengthening the relevance of VOC patterns to grain safety rather than just odor class [54].

For CO2, controlled studies in wheat and corn linked rising gas levels to ergosterol, mycotoxins, germination loss, and dry-matter or compositional changes [5] [6]. This gives CO2 a stronger causal footing than a simple nuisance-gas interpretation [1] [6].

The gap is that relatively few papers report full operational metrics such as drift, false alarms, missed alarms, or cross-season robustness [31] [39].

Confidence level: Medium - The biological validation is credible, but engineering-grade validation remains incomplete [16] [17] [31] [38].

Grain-Type and Storage-Condition Dependence

Commodity dependence

Evidence spans wheat, maize, rice, barley, and mixed bulks, but performance is not uniform [1] [2] [5] [6] [12] [14] [55]. Wheat has the strongest support base for CO2 monitoring [1] [4], while maize and barley also show useful chemical signatures linked to spoilage or mycotoxin risk [6] [12] [18]. Rice evidence supports VOC-based mold detection [14].

A notable contrast is rapeseed: one study reported CO2 was not suitable for early spoilage detection under that model system, despite positive results in wheat [34]. This reinforces that no single threshold or sensor strategy should be assumed universal [34] [40].

Moisture and temperature effects on signal timing

Moisture and temperature strongly accelerate spoilage chemistry [29] [32] [33] [35]. In wheat, tests across multiple moisture contents and temperatures explicitly used CO2 as the primary early-warning signal in place of visual or olfactory checks [35]. In maize silo bags, CO2 reached its maximum limit within the first 26 days at 30°C, while storage at lower temperatures altered that timing [29].

Higher moisture also relates to later increases in intergranular temperature and respiration, fitting the idea that chemical signatures can emerge during the buildup phase before strong thermal manifestations [33].

Aeration, silo geometry, and sensor placement

Aeration and airflow can redistribute gas signals and complicate interpretation [21] [36] [37] [39]. In practice, this means chemical sensing may outperform temperature sensing for early warning, yet still fail to localize hotspots if airflow vents or redistributes CO2 [36] [37]. The recommendation to place a CO2 sensor near the bin center when spoilage location is unknown reflects this operational reality [39].

Taken together, these findings suggest that improved early warning may depend as much on **multi-point placement and airflow-aware design** as on the intrinsic sensitivity of the sensor chemistry [36] [39].

Confidence level: High - Context dependence is repeatedly emphasized across commodity and engineering studies [29] [32] [34] [36].

Synthesis and Practical Interpretation

The evidence supports three increasingly specific conclusions.

1. **CO2 is the most established early spoilage biomarker in stored grain** [1] [2] [4] [5] [6].
2. **VOCs provide useful spoilage signatures and can add causal specificity**, especially for fungal or insect problems, but their lead time versus temperature is less well quantified [9] [12] [14] [15] [19].
3. **Chemical sensing generally precedes useful thermal detection**, especially for localized spoilage, because gas generation begins at the metabolic stage whereas detectable bulk heating typically requires more accumulated activity or favorable sensor proximity [20] [21] [23] [27].

Inferred: a practical monitoring hierarchy emerges from the evidence—**CO2 as sentinel, VOCs as diagnostic layer, temperature as escalation/confirmation layer** [19] [21] [26] [46].
Confidence level: Low-to-Medium - This is a synthesis across studies, not a directly tested unified architecture.

Conclusion

The scientific evidence supports a clear qualitative answer: **yes, chemical spoilage signatures—especially rising intergranular CO2—typically appear before thermal signatures become practically detectable in stored grain** [1] [2] [20] [23]. VOC changes also correlate with early deterioration and can help identify spoilage type, but the temporal evidence for VOC lead time is less direct than for CO2 [8] [12] [14] [15].

What the literature does **not** yet provide is a robust, universal, side-by-side numeric lead-time benchmark for CO2/VOC sensors versus temperature sensors across commodities and storage conditions [22] [23] [24] [29] [31]. So the best evidence-based statement is:

Chemical sensing is earlier in principle and usually earlier in practice, but the exact number of hours or days of warning depends on moisture, commodity, airflow, sensor placement, and the biology of the spoilage event [20] [21] [34].

Overall confidence level: High for the direction of effect; Medium-Low for exact lead-time magnitude - The sequence is well supported, but precise timing remains undermeasured [20] [27] [31].

References

1. White, N.D.G., Sinha, R.N., Muir, W.E. (1982) INTERGRANULAR CARBON DIOXIDE AS AN INDICATOR OF BIOLOGICAL ACTIVITY ASSOCIATED WITH THE SPOILAGE OF STORED WHEAT.. *CAN AGRIC ENG*. https://www.scopus.com/pages/publications/0020145479
2. Muir, W.E., Waterer, D., Sinha, R.N. (1985) CARBON DIOXIDE AS AN EARLY INDICATOR OF STORED CEREAL AND OILSEED SPOILAGE.. *Transactions of the American Society of Agricultural Engineers*. https://www.scopus.com/pages/publications/0022119036
3. Raudienė, E., Rušinskas, D., Balčiūnas, G., (...), Gailius, D. (2015) Early detection of fungi in grain using CO2 level measurement. *XXI IMEKO World Congress "Measurement in Research and Industry"*. https://www.scopus.com/pages/publications/84951196875
4. Guo, L.-H., Zhai, H.-C., Cai, J.-P. (2015) Characteristics of carbon dioxide production by insect and fungus growth in stored wheat. *Modern Food Science and Technology*. https://doi.org/10.13982/j.mfst.1673-9078.2015.4.025
5. Abramson, D., Hulasare, R., York, R.K., (...), Jayas, D.S. (2005) Mycotoxins, ergosterol, and odor volatiles in durum wheat during granary storage at 16% and 20% moisture content. *Journal of Stored Products Research*. https://doi.org/10.1016/j.jspr.2003.11.002
6. Dos Santos Bilhalva, N., Coradi, P.C., Biduski, B., (...), Muller, E.I. (2023) Early detection of quality alterations in corn grains stored in vertical prototype silos using real-time monitoring of carbon dioxide and equilibrium moisture content. *Food and Bioproducts Processing*. https://doi.org/10.1016/j.fbp.2023.06.008
7. Dos Santos Bilhalva, N., Coradi, P.C., Biduski, B., (...), Muller, E.I. (2023) Early detection of quality alterations in corn grains stored in vertical prototype silos using real-time monitoring of carbon dioxide and equilibrium moisture content. *Food and Bioproducts Processing*. https://doi.org/10.1016/j.fbp.2023.06.008
8. Liu, Y., Ren, S., Sun, Q., (...), Zhang, Y. (2024) Analysis of Characteristics Volatile Organic Compounds and Quality Evaluation of Wheat Kernels Storage. *Journal of Nuclear Agricultural Sciences*. https://doi.org/10.11869/j.issn.1000-8551.2024.11.2155
9. Niu, Y., Hua, L., Hardy, G., (...), Ren, Y. (2016) Analysis of volatiles from stored wheat and Rhyzopertha dominica (F.) with solid phase microextraction-gas chromatography mass spectrometry. *Journal of the Science of Food and Agriculture*. https://doi.org/10.1002/jsfa.7274

10. Liu, Y., Ren, S., Sun, Q., (...), Zhang, R. (2024) GC-IMS determination of volatile organic compounds as potential indicators of wheat germination rate. *Food Bioscience*. https://doi.org/10.1016/j.fbio.2023.103535

11. Lu, Y., Wang, J., Mao, H., (...), Yu, M. (2026) Effects of different storage conditions on volatile components of wheat. *Food and Fermentation Industries*. https://doi.org/10.13995/j.cnki.11-1802/ts.044141

12. Weiß, K., Kroschewski, B., Auerbach, H. (2020) Formation of volatile organic compounds during the fermentation of maize as affected by sealing time and silage additive use. *Archives of Animal Nutrition*. https://doi.org/10.1080/1745039X.2019.1694357

13. Żytek, A., Rusinek, R., Oniszczuk, A., Gancarz, M. (2023) Effect of the Consolidation Level on Organic Volatile Compound Emissions from Maize during Storage. *Materials*. https://doi.org/10.3390/ma16083066

14. Guo, J., Qiu, M., Li, L., (...), Liu, X. (2025) Comparative transcriptomic analysis and volatile compound characterization of Aspergillus tubingensis and Penicillium oxalicum during their infestation of Japonica rice. *Food Microbiology*. https://doi.org/10.1016/j.fm.2024.104626

15. Ma, J., Guan, Y., Xing, F., (...), Tai, B. (2023) Accurate and non-destructive monitoring of mold contamination in foodstuffs based on whole-cell biosensor array coupling with machine-learning prediction models. *Journal of Hazardous Materials*. https://doi.org/10.1016/j.jhazmat.2023.131030

16. Olsson, J., Börjesson, T., Lundstedt, T., Schnürer, J. (2000) Volatiles for mycological quality grading of barley grains: Determinations using gas chromatography-mass spectrometry and electronic nose. *International Journal of Food Microbiology*. https://doi.org/10.1016/S0168-1605(00)00355-X

17. Olsson, J., Börjesson, T., Lundstedt, T., Schnürer, J. (2002) Detection and quantification of ochratoxin A and deoxynivalenol in barley grains by GC-MS and electronic nose. *International Journal of Food Microbiology*. https://doi.org/10.1016/S0168-1605(01)00685-7

18. Olsson, J., Börjesson, T., Lundstedt, T., Schnürer, J. (2002) Detection and quantification of ochratoxin A and deoxynivalenol in barley grains by GC-MS and electronic nose. *International Journal of Food Microbiology*. https://doi.org/10.1016/S0168-1605(01)00685-7

19. Olsson, J., Börjesson, T., Lundstedt, T., Schnürer, J. (2000) Volatiles for mycological quality grading of barley grains: Determinations using gas chromatography-mass spectrometry and electronic nose. *International Journal of Food Microbiology*. https://doi.org/10.1016/S0168-1605(00)00355-X

20. Ramachandran, R.P. (2022) Integrated approach on stored grain quality management with CO2 monitoring-A review. *Journal of Stored Products Research*. https://doi.org/10.1016/j.jspr.2022.101950

21. Ma, W., Ma, T., Ji, D., Tsuchikawa, S. (2026) Non-destructive diagnostics and early-warning systems for Penicillium rot in fresh produce: A review of emerging technologies. *Trends in Food Science and Technology*. https://doi.org/10.1016/j.tifs.2026.105793

22. Yin, Q., Li, J., Zhou, L., (...), Li, H. (2024) Classification and recognition method of rice situation based on gray correlation degree. *Journal of Stored Products Research*. https://doi.org/10.1016/j.jspr.2024.102448

23. Cui, H., Wang, S., Yang, X., (...), Wang, S. (2022) Predictive models for assessing the risk of Fusarium pseudograminearum mycotoxin contamination in post-harvest wheat with multi-parameter integrated sensors. *Food Chemistry: X*. https://doi.org/10.1016/j.fochx.2022.100472

24. Huang, H., Danao, M.G.C., Rausch, K.D., Singh, V. (2013) Diffusion and production of carbon dioxide in bulk corn at various temperatures and moisture contents. *Journal of Stored Products Research*. https://doi.org/10.1016/j.jspr.2013.07.002

25. Cai, Lijun, Macfadyen, Sarina, Hua, Baozhen, (...), Ren, Yonglin (2022) *Insects*. https://doi.org/10.3390/insects13050478

26. Dos Santos Bilhalva, N., Coradi, P.C., Biduski, B., (...), Muller, E.I. (2023) Early detection of quality alterations in corn grains stored in vertical prototype silos using real-time monitoring of carbon dioxide and equilibrium moisture content. *Food and Bioproducts Processing*. https://doi.org/10.1016/j.fbp.2023.06.008

27. Liu, C., Chen, G., Zhou, Y., (...), Zhang, Z. (2022) Element tests and simulation of effects of vertical pressure on compression and mildew of wheat. *Computers and Electronics in Agriculture*. https://doi.org/10.1016/j.compag.2022.107447

28. Kim, Y., Kang, S., Ajani, O.S., (...), Ha, Y. (2024) Predicting early mycotoxin contamination in stored wheat using machine learning. *Journal of Stored Products Research*. https://doi.org/10.1016/j.jspr.2024.102294

29. https://www.sciencedirect.com/science/article/pii/S1936976X23000920

30. Neethirajan, S., Freund, M.S., Jayas, D.S., (...), White, N.D.G. (2010) Development of carbon dioxide (CO2) sensor for grain quality monitoring. *Biosystems Engineering*. https://doi.org/10.1016/j.biosystemseng.2010.05.002

31. Wu, Y., Zhu, Y., Lin, J., (...), Zhang, X. (2026) Stage-aware, gas-led early warning of cotton smoldering: A stored-product safety review. *Journal of Stored Products Research*. https://doi.org/10.1016/j.jspr.2026.103139

32. Burrell, N.J., Knight, G.P., Armitage, D.M., Hill, S.T. (1980) Determination of the time available for drying rapeseed before the appearance of surface moulds. *Journal of Stored Products Research*. https://doi.org/10.1016/0022-474X(80)90007-7

33. Oluwakayode, A., Sulyok, M., Krska, R., Medina, A. (2024) The effect of the interactions of water activity, and temperature on OTA, OTB, and OTα produced by Penicillium verrucosum in a mini silo of natural and inoculated wheat using CO2 production as fungal activity sentinel. *Food Chemistry*. https://doi.org/10.1016/j.foodchem.2024.140590

34. Raudienė, E., Gailius, D., Balčiūnas, G., (...), Dobiliene, J. (2018) Precocious rapeseed spoilage detection. *Journal of Physics: Conference Series*. https://doi.org/10.1088/1742-6596/1065/23/232001

35. Lutz, É., Coradi, P.C. (2022) Applications of new technologies for monitoring and predicting grains quality stored: Sensors, Internet of Things, and Artificial Intelligence. *Measurement: Journal of the International Measurement Confederation*. https://doi.org/10.1016/j.measurement.2021.110609

36. Ramachandran, R.P. (2022) Integrated approach on stored grain quality management with CO2 monitoring-A review. *Journal of Stored Products Research*. https://doi.org/10.1016/j.jspr.2022.101950

37. Liu, Jingyun, Li, Ping (2021) Control and Real-Time Data Acquisition of an Experimental Platform for Stored Grain Aeration Study. *Sensors (Basel, Switzerland)*. https://doi.org/10.3390/s21165403

38. Neethirajan, S., Freund, M.S., Jayas, D.S., (...), White, N.D.G. (2010) Development of carbon dioxide (CO2) sensor for grain quality monitoring. *Biosystems Engineering*. https://doi.org/10.1016/j.biosystemseng.2010.05.002

39. Neethirajan, S., Freund, M.S., Jayas, D.S., (...), White, N.D.G. (2010) Development of carbon dioxide (CO2) sensor for grain quality monitoring. *Biosystems Engineering*. https://doi.org/10.1016/j.biosystemseng.2010.05.002

40. Kononets, Y., Rabenseifer, R., Kuneš, R., (...), Babenko, M. (2027) Spatial control of warehouse microclimate heterogeneity to preserve grain quality. *Food Control*. https://doi.org/10.1016/j.foodcont.2026.112486

41. Jian, F., Jayas, D.S., White, N.D.G., Fields, P.G. (2014) Carbon dioxide sorption by stored canola under different storage conditions. *Journal of Stored Products Research*. https://doi.org/10.1016/j.jspr.2014.05.009

42. Bartosik, R., Urcola, H., Cardoso, L., (...), Busato, P. (2023) Silo-bag system for storage of grains, seeds and by-products: A review and research agenda. *Journal of Stored Products Research*. https://doi.org/10.1016/j.jspr.2022.102061

43. Marcos Valle, F.J., Gastón, A., Abalone, R.M., (...), Bartosik, R.E. (2021) Study and modelling the respiration of corn seeds (Zea mays L.) during hermetic storage. *Biosystems Engineering*. https://doi.org/10.1016/j.biosystemseng.2021.05.009

44. Ubhi, G.S., Sadaka, S. (2015) Temporal valuation of corn respiration rates using pressure sensors. *Journal of Stored Products Research*. https://doi.org/10.1016/j.jspr.2015.02.004

45. Arias-Barreto, A., Abalone, R., Gastón, A., Bartosik, R. (2013) Analysis of storage conditions of a wheat silo-bag for different weather conditions by computer simulation. *Biosystems Engineering*. https://doi.org/10.1016/j.biosystemseng.2013.10.012

46. Magan, N., Evans, P. (2000) Volatiles as an indicator of fungal activity and differentiation between species, and the potential use of electronic nose technology for early detection of grain spoilage. *Journal of Stored Products Research*. https://doi.org/10.1016/S0022-474X(99)00057-0

47. Paolesse, R., Alimelli, A., Martinelli, E., (...), Fanelli, C. (2006) Detection of fungal contamination of cereal grain samples by an electronic nose. *Sensors and Actuators, B: Chemical*. https://doi.org/10.1016/j.snb.2005.12.047

48. Dong, L., Liu, R., Dong, H., (...), Zhao, C. (2015) Volatile metabolite profiling of malt contaminated by Fusarium poae during malting. *Journal of Cereal Science*. https://doi.org/10.1016/j.jcs.2015.09.006

49. Hashem, A.S., Guedes, R.N.C., Alhayyani, S., (...), Aljameeli, M.M. (2025) Flour beetle responses to volatile organic compounds from uninfested and previously infested wheat flour. *Journal of Stored Products Research*. https://doi.org/10.1016/j.jspr.2025.102766

50. Buśko, Maciej, Stuper, Kinga, Jeleń, Henryk, (...), Perkowski, Juliusz (2016) Comparison of Volatiles Profile and Contents of Trichothecenes Group B, Ergosterol, and ATP of Bread Wheat, Durum Wheat, and Triticale Grain Naturally Contaminated by Mycobiota. *Frontiers in Plant Science*. https://doi.org/10.3389/fpls.2016.01243

51. Arias-Barreto, A., Abalone, R., Gastón, A., (...), Bartosik, R. (2017) Validation of a heat, moisture and gas concentration transfer model for soybean (Glycine max) grains stored in plastic bags (silo bags). *Biosystems Engineering*. https://doi.org/10.1016/j.biosystemseng.2017.03.009

52. https://www.sciencedirect.com/science/article/pii/S142482202604118X

53. Emadi, T.A., Shafai, C., Thomson, D.J., (...), Jayas, D.S. (2013) Polymer-based chemicapacitor sensor for 1-octanol and relative humidity detections at different temperatures and frequencies. *IEEE Sensors Journal*. https://doi.org/10.1109/JSEN.2012.2220760

54. Keshri, G., Magan, N. (2000) *Journal of Applied Microbiology*. https://doi.org/10.1046/j.1365-2672.2000.01185.x

55. Wang, Shiming, Zhao, Yan, Lv, Haoxin, (...), Song, Yongling (2025) *Foods*. https://doi.org/10.3390/foods14183217