


A

Shared primary warm-start-region selection

The joint workflow uses the saved k=2 primary partition.
80% subsample silhouette maxima: k=2: 100/100.


Average silhouette



B

Selected-family joint-objective eligibility

Blue region: lowest 10%; vertical lines mark 5%, 10%, and 20% sets.




C

Context-specific top-50 endpoint robustness

Text gives the modal result and exact support; a red cross marks a change after collapsing duplicate parameter endpoints.

in vivo	Both ploidy regimes present at 0% O2	yes 100%		yes 100%
	Both ploidy regimes present at 1% O2	yes 100%		no 100%
	Both ploidy regimes present at 5% O2	no 100%		yes 100%
	Unmodified trajectory: O2 response direction	decrease 100%		decrease 100%
	Surface O2 response at low p_misseg	approximately flat 100%		increase 100%
	Surface O2 response at intermediate p_misseg	decrease 100%		approximately flat 100%
	Surface O2 response at high p_misseg	decrease 100%		decrease 100%
	At least half of the prespecified diagnostic union has spectral gap >= 0.005	no 100%		no 100%
in vitro	Both ploidy regimes present at 0% O2	yes 100%		yes 100%
	Both ploidy regimes present at 1% O2	yes 100%		yes 100%
	Both ploidy regimes present at 5% O2	yes 100%		yes 100%
	Unmodified trajectory: O2 response direction	increase 100%		increase 100%
	Surface O2 response at low p_misseg	approximately flat 100%		approximately flat 100%
	Surface O2 response at intermediate p_misseg	increase 100%		increase 100%
	Surface O2 response at high p_misseg	decrease 100%		decrease 100%
	At least half of the prespecified diagnostic union has spectral gap >= 0.005	yes 100%		yes 100%
		C01	Selected warm-start family	C02



A records the 2 primary warm-start regions. B-C include C01, C02, the pairs displayed in Figure 7; optimizer seeds are not biological replicates.