



A



B



C

Context-specific top-50 endpoint robustness

Text gives the modal result and exact support; a red cross marks a change after collapsing duplicate parameter endpoints.

	Modal support within the top-50 ensemble		
	1.0	0.9	0.8
In vivo	Both ploidy regimes present at 0% O2	yes 100%	yes 100%
	Both ploidy regimes present at 1% O2	no 100%	yes 100%
	Both ploidy regimes present at 5% O2	no 100%	no 100%
	Unmodified trajectory: O2 response direction	decrease 100%	increase 100%
	Surface O2 response at low effective missegregation	increase 100%	increase 100%
	Surface O2 response at intermediate effective missegregation	increase 100%	increase 100%
	Surface O2 response at high effective missegregation	increase 100%	increase 100%
	At least half of the prespecified diagnostic union has spectral gap ≥ 0.005	no 100%	no 100%
In vitro	Both ploidy regimes present at 0% O2	yes 100%	yes 100%
	Both ploidy regimes present at 1% O2	yes 100%	yes 100%
	Both ploidy regimes present at 5% O2	yes 100%	yes 100%
	Unmodified trajectory: O2 response direction	approximately flat 100%	approximately flat 100%
	Surface O2 response at low effective missegregation	approximately flat 100%	approximately flat 100%
	Surface O2 response at intermediate effective missegregation	increase 100%	increase 100%
	Surface O2 response at high effective missegregation	decrease 100%	decrease 100%
	At least half of the prespecified diagnostic union has spectral gap ≥ 0.005	yes 100%	yes 100%
		C01	C02
		Selected warm-start family	
		C03	

A defines the three primary warm-start regions. B-C retain only the C01, C02, and C03 pairs displayed in Figure 6; optimizer seeds are not biological replicates.