

Sobol sensitivity | N4096, five repeats, 201 oxygen points
Cross-scope row order: neighborhood ploidy; directly ordered by high/low window scores.


A Global full bounds | S1
Dominant mean ploidy




B 500 neighborhoods | S1
Dominant mean ploidy




C Global full bounds | S1
Net live-cell growth rate




D 500 neighborhoods | S1
Net live-cell growth rate




E Global full bounds | ST
Dominant mean ploidy




F 500 neighborhoods | ST
Dominant mean ploidy




G Global full bounds | ST
Net live-cell growth rate




H 500 neighborhoods | ST
Net live-cell growth rate




I Global full bounds | S2: 36 focal pairs
Dominant mean ploidy




J 500 neighborhoods | S2: 36 focal pairs
Dominant mean ploidy



K Global full bounds | S2: 36 focal pairs
Net live-cell growth rate



L 500 neighborhoods | S2: 36 focal pairs
Net live-cell growth rate



High ≥ Low: sort by High descending; Low > High: sort by Low ascending. No middle sorting group.
S2 annotations 1/2: left/right half = first/second named parameter. Annotation 3 is shown only for S1/ST and never sets row order.
Cells: global five-repeat mean; neighborhood median of 500 five-repeat means. No diagnostic exclusions.
O_2: $\log_{10}$ above 0.025%; linear from 0 to 0.025% to retain zero. Low: 0–1%; intermediate: >1–<3%; high: 3–5%.

1 Process
Division + genome change · Stress-linked missegregation · Oxygen response · Ploidy-linked growth + death · Post-missegregation survival

2 Figure 4B O_2 correlation 3 Sobol O_2 sensitivity (S1/ST only; ploidy-based)
High O_2 · Low O_2 · No resolved O_2 preference