Sobol sensitivity | N4096, five repeats, 201 oxygen points
Row order: ploidy high/low window scores; growth follows the same rows.

A 500 neighborhoods | S2: 91 pairs
Dominant mean ploidy

B 500 neighborhoods | S2: 91 pairs
Net live-cell growth rate

1 2

buffer_smax × buffer_beta
p_mis_base × buffer_beta
p_mis_base × buffer_smax
p_mis_base × O2_crit
p_mis_base × n_O
buffer_beta × n_O
O2_crit × n_O
buffer_beta × buffer_n_exp
buffer_smax × n_O
buffer_beta × gamma_mu
buffer_beta × p_wgd
buffer_beta × alpha_o2
buffer_beta × gamma_growth
p_mis_base × mu_hp
buffer_smax × buffer_n_exp
p_mis_base × p_misseg
p_mis_base × k_o_mis
p_mis_base × buffer_n_exp
p_misseg × n_O
p_mis_base × gamma_mu
p_mis_base × p_wgd
gamma_mu × n_O
buffer_n_exp × O2_crit
buffer_n_exp × n_O
p_mis_base × alpha_o2
buffer_n_exp × gamma_mu
buffer_n_exp × alpha_o2
p_mis_base × gamma_growth
buffer_n_exp × p_wgd
buffer_n_exp × gamma_growth
buffer_n_exp × mu_hp
lam_max × p_wgd
lam_max × k_o_mis
lam_max × alpha_o2
lam_max × buffer_n_exp
lam_max × gamma_growth
lam_max × gamma_mu
lam_max × n_O
lam_max × p_mis_base
p_misseg × p_wgd
p_wgd × O2_crit
p_wgd × gamma_growth
p_wgd × gamma_mu
p_wgd × alpha_o2
p_wgd × n_O
p_wgd × mu_hp
lam_max × p_misseg
alpha_o2 × gamma_mu
lam_max × buffer_smax
p_misseg × buffer_n_exp
alpha_o2 × gamma_growth
alpha_o2 × n_O
k_o_mis × p_wgd
lam_max × mu_hp
lam_max × O2_crit
k_o_mis × alpha_o2
p_misseg × gamma_growth
k_o_mis × gamma_growth
gamma_growth × n_O
k_o_mis × buffer_n_exp
p_misseg × alpha_o2
gamma_growth × gamma_mu
alpha_o2 × mu_hp
gamma_growth × mu_hp
k_o_mis × gamma_mu
buffer_smax × p_wgd
gamma_growth × O2_crit
alpha_o2 × O2_crit
lam_max × buffer_beta
k_o_mis × n_O
p_misseg × k_o_mis
p_misseg × gamma_mu
mu_hp × n_O
mu_hp × gamma_mu
buffer_smax × gamma_growth
gamma_mu × O2_crit
buffer_smax × alpha_o2
k_o_mis × mu_hp
buffer_smax × gamma_mu
k_o_mis × buffer_smax
p_misseg × mu_hp
k_o_mis × buffer_beta
p_misseg × O2_crit
p_misseg × buffer_smax
buffer_smax × mu_hp
k_o_mis × O2_crit
p_misseg × buffer_beta
mu_hp × O2_crit
buffer_beta × mu_hp
buffer_beta × O2_crit
buffer_smax × O2_crit

Low Mid High

0 0.025 0.1 0.5 1 3 5
Fixed oxygen (%)

0 0.5 1
Diagnostic pass fraction

High ≥ Low: sort by High descending; Low > High: sort by Low ascending. No middle sorting group.
S2 annotations 1/2: left/right half = first/second named parameter. S2 has only two annotation columns.
Cells: global five-repeat mean; neighborhood median of 500 five-repeat means. No diagnostic exclusions.
O₂: $\log_{10}$ above 0.025%; linear from 0 to 0.025% to retain zero. Low: 0–1%; intermediate: >1–<3%; high: 3–5%.

1 Process
Division + genome change
Stress-linked missegregation
Oxygen response
Ploidy-linked growth + death
Post-missegregation survival

2 Figure 4B O₂ correlation
High O₂
Low O₂
No resolved O₂ preference