

Sobol sensitivity | N4096, five repeats, 201 oxygen points
Row order and sensitivity annotations: ploidy-based; growth follows the same rows.


A Global full bounds | S1
Dominant mean ploidy




B Global full bounds | S1
Net live-cell growth rate




C Global full bounds | ST
Dominant mean ploidy




D Global full bounds | ST
Net live-cell growth rate



E Global full bounds | S2: 36 focal pairs
Dominant mean ploidy



F Global full bounds | S2: 36 focal pairs
Net live-cell growth rate



High group: high-O_2 score descending; Low group: low-O_2 score ascending; unresolved last. No 0.3 sensitivity gate.
S2 annotations 1/2: left/right half = first/second named parameter. Gray = no resolved O_2 preference, not no output effect.
Cells: global five-repeat mean; neighborhood median of 500 five-repeat means. No diagnostic exclusions.
O_2: $\log_{10}$ above 0.025%; linear from 0 to 0.025% to retain zero. Low: 0–1%; intermediate: >1–<3%; high: 3–5%.

1 Process
Division + genome change · Stress-linked missegregation · Oxygen response · Ploidy-linked growth + death · Post-missegregation survival

2 Figure 4B O_2 correlation 3 Sobol O_2 sensitivity (ploidy-based)
High O_2 · Low O_2 · No resolved O_2 preference