Global full bounds | 36 focal parameter pairs | N4096, five repeats

Dominant mean ploidy

Net live-cell growth rate

Low
Mid
High

p_mis_base × lam_max
p_mis_base × p_misseg
p_mis_base × k_o_mis
p_mis_base × buffer_smax
p_mis_base × buffer_beta
p_mis_base × buffer_n_exp
p_mis_base × p_wgd
p_mis_base × alpha_o2
p_mis_base × gamma_growth
p_mis_base × mu_hp
p_mis_base × gamma_mu
p_mis_base × O2_crit
p_mis_base × n_O
mu_hp × lam_max
mu_hp × p_misseg
mu_hp × k_o_mis
mu_hp × buffer_smax
mu_hp × buffer_beta
mu_hp × buffer_n_exp
mu_hp × p_wgd
mu_hp × alpha_o2
mu_hp × gamma_growth
mu_hp × gamma_mu
mu_hp × O2_crit
mu_hp × n_O
gamma_mu × lam_max
gamma_mu × p_misseg
gamma_mu × k_o_mis
gamma_mu × buffer_smax
gamma_mu × buffer_beta
gamma_mu × buffer_n_exp
gamma_mu × p_wgd
gamma_mu × alpha_o2
gamma_mu × gamma_growth
gamma_mu × O2_crit
gamma_mu × n_O

0
0.025
0.1
0.5
1
3
5

Fixed oxygen (%)

-0.0614
0
0.0614

S2 estimate

-0.141
0
0.141

S2 estimate

Global: five-repeat mean; neighborhoods: median of 500 five-repeat means. All estimates retained.
Positive estimates: white to purple. Negative estimates: blue, not antagonism.
O_2: $\log_{10}$ above 0.025%; linear from 0 to 0.025% to retain zero. Low: 0–1%; intermediate: >1–<3%; high: 3–5%.