

Machine Learning

Probability Review for Discrete Random Variables

Fabio Vandin

October 9th, 2023

Theorem

Let $g(X)$ be a function of a discrete random variable X . Then $\mathbb{E}[g(X)] = \sum_x g(x)p_X(x)$.

For a random variable X we define:

- **Mean:** $m_X \doteq \mathbb{E}[X]$
- **Variance:** $\sigma_X^2 \doteq \mathbb{E}[(X - m_X)^2] = \mathbb{E}[X^2] - m_X^2 = \mathbf{Var}[X]$
- k -th moment: $\mathbb{E}[X^k]$

Example die rally

\underline{Y} = outcome of a die

mean: $m_{\underline{Y}} = E[\underline{Y}] = \frac{1}{6}(1+2+3+4+5+6) = 3.5$

variance: $\sigma_{\underline{Y}}^2 = E[\underline{Y}^2] - m_{\underline{Y}}^2$

$$= \frac{1}{6}(1+4+9+16+25+36) - (3.5)^2$$

$$= \frac{35}{12} \approx 2.916 \dots$$

For a vector valued r.v. $\mathbf{X} \in \mathbb{R}^n$

Expectation:

$$\mathbb{E}[\mathbf{X}] = \begin{bmatrix} m_{X_1} \\ \vdots \\ m_{X_n} \end{bmatrix}$$

$$\vec{X} = \begin{bmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{bmatrix}$$
$$m_{X_i} = \mathbb{E}[X_i]$$

Instead of the variance, we have the *covariance matrix*:

$$\Sigma = \mathbb{E}[(\mathbf{X} - m_{\mathbf{X}})(\mathbf{X} - m_{\mathbf{X}})^T] = \begin{bmatrix} \sigma_{X_1}^2 & \sigma_{X_1, X_2} & \cdots & \sigma_{X_1, X_n} \\ \sigma_{X_2, X_1} & \sigma_{X_2}^2 & \vdots & \sigma_{X_2, X_n} \\ \vdots & \vdots & \ddots & \vdots \\ \sigma_{X_n, X_1} & \sigma_{X_n, X_2} & \cdots & \sigma_{X_n}^2 \end{bmatrix}$$

where

$$\sigma_{X_i, X_j} = \text{Cov}(X_i, X_j) \doteq \mathbb{E}[(X_i - m_{X_i})(X_j - m_{X_j})]$$

σ_{X_i, X_j} is the covariance of X_i and X_j

Theorem

If X_1, X_2 are independent then $\sigma_{X_1, X_2} = 0$.

e) $\vec{X} = \begin{bmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{bmatrix}$, X_1, X_2, \dots, X_n are mutually independent
r. v.'s

$$\Sigma = \begin{bmatrix} \sigma_{X_1}^2 & 0 & \dots & 0 \\ 0 & \sigma_{X_2}^2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \sigma_{X_n}^2 \end{bmatrix}$$

•) If $\sigma_{X_1, X_2} = 0$ then X_1, X_2 are independent? No!

Theorem

If X_1, X_2 are independent then $\sigma_{X_1, X_2} = 0$.

The other direction is not true!

Counter example: $X_1 = \begin{cases} 1 & \text{with prob. } \frac{1}{2} \\ -1 & \text{with prob. } \frac{1}{2} \end{cases} \quad \mathbb{E}[X_1] = 0$

$$X_2 = \begin{cases} 0 & \text{if } X_1 = -1 \\ -1 & \text{with prob. } \frac{1}{2} \text{ if } X_1 = 1 \\ +1 & \text{with prob. } \frac{1}{2} \text{ if } X_1 = 1 \end{cases}$$

$$\mathbb{E}[X_2] = -1 \cdot \frac{1}{2} + 1 \cdot \frac{1}{2} = 0$$

But:

$\Rightarrow X_1, X_2$ are not independent

HW: prove it

$$\begin{aligned} \sigma_{X_1, X_2} &= \mathbb{E}[(X_1 - m_{X_1})(X_2 - m_{X_2})] \\ &= \mathbb{E}[X_1 X_2 - \underbrace{m_{X_1} X_2}_0 - \underbrace{m_{X_2} X_1}_0 + \underbrace{m_{X_1} m_{X_2}}_0] \\ &= \mathbb{E}[X_1 X_2] = 0 \cdot \dots + 1 \cdot \frac{1}{2} + (-1) \cdot \frac{1}{2} = 0 \end{aligned}$$

constants

Theorem (Properties of Mean, Variance, etc.)

- $\mathbb{E}[X_1 + X_2] = \mathbb{E}[X_1] + \mathbb{E}[X_2]$ (linearity of expectation)
- $\mathbf{Var}(aX + b) = a^2 \mathbf{Var}[X]$
- $\mathbf{Var}[X_1 + X_2] = \mathbf{Var}[X_1] + \mathbf{Var}[X_2] + 2\sigma_{X_1, X_2}$

Corollary

If $\sigma_{X_1, X_2} = 0$ then $\mathbf{Var}[X_1 + X_2] = \mathbf{Var}[X_1] + \mathbf{Var}[X_2]$

Corollary

If X_1, X_2 (r.v.'s) are independent
 $\Rightarrow \mathbf{Var}[X_1 + X_2] = \mathbf{Var}[X_1] + \mathbf{Var}[X_2]$

Conditional Probability

Definition (Conditional probability)

A, B are events: $\mathbb{P}[A|B] \doteq \frac{\mathbb{P}[A \cap B]}{\mathbb{P}[B]}$. Well defined only if $\mathbb{P}[B] > 0$.

Example (Relative frequency, convergence, and conditional probability)

Consider an event A . X_1, \dots, X_n that are independent and identically distributed (i.i.d.) random variables that are indicator functions:

$$X_i(z) = \begin{cases} 1 & \text{if } z \in A \\ 0 & \text{if } z \notin A \end{cases}$$

$$X_i = \begin{cases} 1 & \text{if } A \text{ happens} \\ 0 & \text{otherwise} \end{cases}$$

Example (Relative frequency, convergence, and conditional probability)

Consider an event A . X_1, \dots, X_n that are independent and identically distributed (i.i.d.) random variables that are indicator functions:

$$X_i(z) = \begin{cases} 1 & \text{if } z \in A \\ 0 & \text{if } z \notin A \end{cases}$$

$$S_n = \sum_{i=1}^n X_i$$

Relative frequency of event A : $f_n(A) \doteq \frac{S_n}{n}$ \rightarrow proportion (ratio) of the n trials in which event A happened

Example (Relative frequency, convergence, and conditional probability)

Consider an event A . X_1, \dots, X_n that are independent and identically distributed (i.i.d.) random variables that are indicator functions:

$$X_i(z) = \begin{cases} 1 & \text{if } z \in A \\ 0 & \text{if } z \notin A \end{cases}$$

$$S_n = \sum_{i=1}^n X_i$$

Relative frequency: $f_n(A) \doteq \frac{S_n}{n}$

Example

Coin flips, event A = “the result of the coin flip is head”

Each X_i is a **Bernoulli r.v.** of parameter p : $X_i \sim B(p)$

Bernoulli: r.v. of
parameter p

Each X_i is a **Bernoulli r.v.** of parameter p : $X_i \sim B(p)$

$$p = \mathbb{P}[X_i = 1] = \mathbb{P}[z \in A]$$


Then $S_n = \sum_{i=1}^n X_i$ is a **Binomial r.v.** of parameters n, p :

$$S_n \sim \text{Bin}(n, p)$$

$$\begin{aligned} & \mathbb{P}_t[S_n = k] \\ &= \binom{n}{k} p^k \cdot (1-p)^{n-k} \end{aligned}$$

$$\mathbb{E}[S_n] = np$$

$$\text{Var}[S_n] = np(1-p)$$



Each X_i is a **Bernoulli r.v.** of parameter p : $X_i \sim B(p)$

$$p = \mathbb{P}[X_i = 1] = \mathbb{P}[z \in A]$$

Then $S_n = \sum_{i=1}^n X_i$ is a **Binomial r.v.** of parameters n, p :

$$S_n \sim \text{Bin}(n, p)$$

$$\mathbb{P}[S_n = k] = \binom{n}{k} p^k (1-p)^{n-k}$$

Then

$$\mathbb{E}[S_n] = np$$

$$\text{Var}[S_n] = np(1-p)$$

Exercise

Derive $\mathbb{E}[S_n]$ and $\text{Var}[S_n]$.

Let's go back to the relative frequency $f_n(A) \doteq \frac{S_n}{n}$:

$$S_n \sim \text{Bin}(n, p)$$

$$\mathbb{E}[f_n(A)] = \mathbb{E}\left[\frac{S_n}{n}\right]$$

$$= \frac{1}{n} \mathbb{E}[S_n] = \frac{1}{n} \cdot np = p$$

Exercise

Derive $\mathbb{E}[S_n]$ and $\text{Var}[S_n]$.

Let's go back to the relative frequency $f_n(A) \doteq \frac{S_n}{n}$:

$$\mathbb{E}[f_n(A)] = p$$

and

$$\begin{aligned}\text{Var}[f_n(A)] &= \text{Var}\left[\frac{S_n}{n}\right] \\ &= \left(\frac{1}{n}\right)^2 \text{Var}[S_n] \\ &= \frac{1}{n^2} np(1-p) \\ &= \frac{p(1-p)}{n}\end{aligned}$$

Exercise

Derive $\mathbb{E}[S_n]$ and $\mathbf{Var}[S_n]$.

Let's go back to the relative frequency $f_n(A) \doteq \frac{S_n}{n}$:

$$\mathbb{E}[f_n(A)] = p$$

and

$$\mathbf{Var}[f_n(A)] = \frac{p(1-p)}{n}$$

Theorem (Chebyshev's inequality)

Let X be a r.v. with $\mathbb{E}[X] = \mu$ and $\mathbf{Var}[X] = \sigma^2$. Then:

$$\mathbb{P}[|X - \mu| > \epsilon] \leq \frac{\sigma^2}{\epsilon^2}.$$

Therefore

$$\mathbb{P}[|f_n(A) - p| > \varepsilon] \leq \frac{p(1-p)}{n\varepsilon^2}$$

and

$$\lim_{n \rightarrow +\infty} f_n(A) = p$$

Note: there are tighter bounds than Chebyshev's, like Chernoff's and Hoeffding's - we will see them later.

Intermission

Theorem (Law of Large Numbers)

Let $X_i, i = 1, \dots, n$ be i.i.d. with $\mathbb{E}[X_i] = \mu$ and $\text{Var}[X] = \sigma^2 < +\infty$.

Then

$$\lim_{n \rightarrow +\infty} \mathbb{P} \left[\left| \frac{1}{n} \sum X_i - \mu \right| > \varepsilon \right] = 0.$$

Example (continue)

Remark 1:

$$\lim_{n \rightarrow +\infty} f_n(A) = \mathbb{P}[A]$$

Remark 2:

$$\mathbb{P}[A|B] = \frac{\mathbb{P}[A \cap B]}{\mathbb{P}[B]} = \lim_{n \rightarrow +\infty} \frac{f_n(A \cap B)}{f_n(B)}$$

$$\frac{f_n(A \cap B)}{f_n(B)} = \frac{S_n(A \cap B)}{S_n(B)}$$

it's the fraction of times $A \cap B$ happens among those in which B happens.

Computing Conditional Probabilities

Definition (Conditional probability)

A, B are events: $\mathbb{P}[A|B] \doteq \frac{\mathbb{P}[A \cap B]}{\mathbb{P}[B]}$. Well defined only if $\mathbb{P}[B] > 0$.

Theorem (Bayes Rule)

$$\mathbb{P}[A|B] = \frac{\mathbb{P}[B|A]\mathbb{P}[A]}{\mathbb{P}[B]}$$


Theorem (Law of Total Probability)

Let C_1, C_2, \dots, C_n be a partition of Ω :

- $\bigcup_{i=1}^n C_i = \Omega$
- $C_i \cap C_j = \emptyset$

For all $A \subset \Omega$:

$$\mathbb{P}[A] = \sum_{i=1}^n \mathbb{P}[A|C_i]\mathbb{P}[C_i]$$



Theorem (Law of Total Probability)


Let C_1, C_2, \dots, C_n be a partition of Ω :

- $\bigcup_{i=1}^n C_i = \Omega$
- $C_i \cap C_j = \emptyset$

For all $A \subset \Omega$:

$$\mathbb{P}[A] = \sum_{i=1}^n \mathbb{P}[A|C_i]\mathbb{P}[C_i]$$

Example: $\mathbb{P}[B] = \mathbb{P}[B|A]\mathbb{P}[A] + \mathbb{P}[B|A^c]\mathbb{P}[A^c]$


"not A"

Example

M = “have a rare disease”, with $\mathbb{P}[M] = 10^{-9}$

T = “test for the disease is positive” with:

- $\mathbb{P}[T|M] = 0.99$ (1% false negatives)
- $\mathbb{P}[T|M^c] = 0.001$ (0.1% false positives)

If you test positive, what is the probability that you have the disease?

High : 27

Low : 13

We should take a break. Not enough for
my taste
~

Example

M = “have a rare disease”, with $\mathbb{P}[M] = 10^{-9}$

T = “test for the disease is positive” with:

- $\mathbb{P}[T|M] = 0.99$ (1% false negatives)
- $\mathbb{P}[T|M^c] = 0.001$ (0.1% false positives)

If you test positive, what is the probability that you have the disease?

$$\mathbb{P}[M|T] = \frac{\mathbb{P}[T|M]\mathbb{P}[M]}{\mathbb{P}[T]}\mathbb{P}[M] = \frac{0.99 * 10^{-9}}{0.99 * 10^{-9} + 0.001(1 - 10^{-9})}$$
$$\approx \frac{1}{1 + 10^6} \approx 10^{-6}$$

low of total Prob.:

$$= \mathbb{P}[T|M] \cdot \mathbb{P}[M] + \mathbb{P}[T|M^c] \cdot \mathbb{P}[M^c]$$