

AFTER VISIT SUMMARY

Berdia M. Monson MRN: 004349698 CEID: UAM-XZBS-03ZZ-B7TR

UAMS Hospital Carcinosarcoma of uterus (HCC) 1/14/2026 - 1/15/2026 H7 - Medical Oncology

Instructions



Talk with your provider about your medications

- ASK how to take:
cyanocobalamin 500 MCG tablet
iron 325 mg (65 mg elemental iron) tablet (ferrous sulfate)
rosuvastatin 5 MG tablet (Crestor)

- CONTINUE taking your other medications
Review details of your medication changes below.

Your Next Steps



Go

FEB
6

Postop 11:20 AM
Lisa Bazzett-Matabele, MD
Women's Cancer Clinic -- Gynecology
501-686-8522

You are allergic to the following

Allergen	Reactions
Codeine	Other (See Comments)
Penicillins	Other (See Comments)

Date Reviewed: Jan 14, 2026

You are intolerant to the following

No active intolerances/contraindications

Date Reviewed: Jan 14, 2026

Your Doctors

Provider	Service	Role	Specialty
Bazzett-Matabele, Lisa, MD	—	Attending	Gynecologic Oncology
RESOURCE, CARDIOLOGY	—	Consulting Physician	Cardiology

Providers seen during hospitalization

Provider	Service	Role	Specialty	Primary Office Phone
Attending				
Bazzett-Matabele, Lisa, MD	—	Attending	Gynecologic Oncology	501-686-8000
Others who treated you				
RESOURCE, CARDIOLOGY	—	Consulting Physician	Cardiology	—

What's Next

FEB
6

Postop with Lisa Bazzett-Matabele, MD
Friday Feb 6, 2026 11:20 AM

Women's Cancer Clinic --
Gynecology
4018 W Capitol Ave.
Little Rock AR 72205
501-686-8522

UAMS CARES

If you feel you or a family member/friend is at risk of suicide please call the National Suicide Prevention Lifeline call, text or chat at 988.

APPOINTMENT CONTACT INFORMATION

To cancel or reschedule an appointment please call the clinic number listed in your clinic appointment.

If you are interested in making a new appointment with a different specialty or service at UAMS please call (501) 686-8000.

Sign up for Text Message Appointment Reminders

To sign up for text message appointment reminders from UAMS, please log into MyChart and make sure your mobile phone number is correct. If you don't have MyChart, you can sign up today using the instructions above or below.

After you log into MyChart, click on the **"Menu"** button. Under **"Account Settings,"** click on **"Communication Preferences."**

Under the Contact Information section at the bottom, make sure your mobile phone number is correct. If it is not correct, click **"Edit,"** type in the correct number, and click **"Save Changes."**

If you are not signed up for text messages from UAMS, click the **"Sign up now"** link in the Communication Preferences section. You can choose the type of information you'd like to receive by text message.

****Standard text message rates apply.****

Sign up for Text Message Appointment Reminders (continued)

UAMS Text Message Reminder Disclaimer

Warning: UAMS does not guarantee information sent or received via text message is secure. There are security risks associated with sending information via text message. For example, an unauthorized person or entity may access or use the information. By requesting that UAMS send you information via text message, you acknowledge you have been warned of and accept such risks.

We Want to Hear from You!

Thank you for choosing UAMS! We would like to hear about your experience. Soon you may receive a survey with questions about your care. Please complete the survey. Your feedback helps us continue to provide the best possible experience for you and other patients and families.

Your Medication List - The checkmarks below show the times you should take your medicines.

CONTINUE taking these medications

	Morning	Around Noon	Afternoon	Evening	Bedtime	As Needed
 albuterol 90 mcg/actuation inhaler Commonly known as: HFA every 4 (four) hours.	every 4 (four) hours.					
 amLODIPine 10 MG tablet Commonly known as: Norvasc Take one tablet (10 mg) by mouth daily.	 1 tablet					
 aspirin 81 MG EC tablet Take one tablet (81 mg) by mouth daily.	 1 tablet					
 BLACK SEED ORAL Take by mouth.	Take by mouth.					
 calcium carbonate-vitamin D3 600 mg-10 mcg (400 unit) per tablet Take one tablet by mouth daily.	 1 tablet					
 losartan-hydroCHLOROthiazide 100-25 mg per tablet Commonly known as: Hyzaar Take one tablet by mouth daily.	 1 tablet					
 megestrol 40 MG tablet Commonly known as: Megace Take one tablet (40 mg) by mouth daily.	 1 tablet					
 metoprolol tartrate 100 MG tablet Commonly known as: Lopressor Take one tablet (100 mg) by mouth.	Take one tablet (100 mg) by mouth.					

Your Medication List - The checkmarks below show the times you should take your medicines. (continued)

CONTINUE taking these medications (continued)

	Morning	Around Noon	Afternoon	Evening	Bedtime	As Needed
 omeprazole 40 MG capsule Commonly known as: PrilOSEC Take one capsule (40 mg) by mouth daily.	 1 capsule					

ASK your doctor about these medications

	Morning	Around Noon	Afternoon	Evening	Bedtime	As Needed
 cyanocobalamin 500 MCG tablet You are not taking this medication. If you have questions about this medication, ask your provider. Take one tablet (500 mcg) by mouth daily.						
 iron 325 mg (65 mg elemental iron) tablet You are not taking this medication. If you have questions about this medication, ask your provider. Take one tablet (325 mg) by mouth daily with breakfast. Generic drug: ferrous sulfate						
 rosuvastatin 5 MG tablet You are not taking this medication. If you have questions about this medication, ask your provider. Commonly known as: Crestor Take one tablet (5 mg) by mouth daily.						

January 2026

Sunday	Monday	Tuesday	Wednesday	Thursday	Friday	Saturday
				1	2	3
4	5	6	7	8	9	10
11	12	13	14 Admission 6:07 AM Bazzett-Matabele, Lisa, MD H7 - Medical Oncology (Discharge: 1/15/2026) ROBOTIC TOTAL LAPAROSCOPIC HYSTERECTOMY WITH SALPINGOOPHORECTOMY (XI) 8:12 AM Bazzett-Matabele, Lisa, MD UAMS Main OR	15	16	17
18	19	20	21	22	23	24
25	26	27	28	29	30	31

February 2026

Sunday	Monday	Tuesday	Wednesday	Thursday	Friday	Saturday
1	2	3	4	5	6 Postop 11:20 AM (20 min.) Bazzett- Matabele, Lisa, MD Women's Cancer Clinic -- Gynecology	7
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28

Active Treatment & Therapy Days for Monson, Berdia M (until 2/14/2026)

Monson, Berdia M does not have any active plans of the following types: INFUSION TREATMENT, OB/GYN Therapy Plans, Therapy Plan 2, Therapy Plan 3, Therapy Plan 4, Therapy Plan 5, Therapy Plan 6

Active Treatment & Therapy Days for Monson, Berdia M (until 4/15/2026)

Monson, Berdia M does not have any active plans of the following types: ONCOLOGY TREATMENT, ONCOLOGY SUPPORTIVE CARE, INFUSION TREATMENT, ONCOLOGY RESEARCH, ONCOLOGY TREATMENT 2, ONCOLOGY RESEARCH 2, RESEARCH

MyChart

View your After Visit Summary and more online at <https://mychart.uamshealth.com/MyChart/>.

Instructions

Alternating schedules for post-operative pain management without opioids

6 AM ☀️ Ibuprofen 600mg (e.g. Motrin, Advil) 	9 AM ☀️ Acetaminophen 1,000mg (e.g. Tylenol) 	12 PM ☀️ Ibuprofen 600mg 	3 PM ☀️ Acetaminophen 1,000mg 
6 PM 🌙 Ibuprofen 600mg 	9 PM 🌙 Acetaminophen 1,000mg 	12 AM 🌙 Acetaminophen 1,000mg + Ibuprofen 600mg 	Do not take more than 4,000 mg of Acetaminophen or 3,200 mg of Ibuprofen in a 24-hour period

Alternating schedule for post-operative pain management with opioids

6 AM ☀️ Ibuprofen 600mg (e.g. Motrin, Advil) 	9 AM ☀️ Oxycodone 5mg +/- Acetaminophen 1,000mg 	12 PM ☀️ Ibuprofen 600mg 	3 PM ☀️ Oxycodone 5mg +/- Acetaminophen 1,000mg 
6 PM 🌙 Ibuprofen 600mg 	9 PM 🌙 Oxycodone 5mg +/- Acetaminophen 1,000mg 	12 AM 🌙 Acetaminophen 1,000mg + Ibuprofen 600mg 	Do not take more than 4,000 mg of Acetaminophen or 3,200 mg of Ibuprofen in a 24-hour period

If your doctor prescribes an opioid pain medication you can replace one of the acetaminophen or ibuprofen doses with a dose of the opioid. If your pain is manageable, avoid taking opioids.

What pain should you expect after surgery?

- It is normal to have some pain after surgery. Post-operative surgical pain is typically the worst 24-48 hours after your procedure. After that, you should be able to get up, get dressed, and perform small tasks around the home.

What are your pain relief options?

- Heating packs

Instructions (continued)

- Ice packs
- Rest
- Scheduled ibuprofen and acetaminophen
- Opioids (if prescribed)

What if your doctor prescribed opioids for post-operative pain?

- If your doctor prescribed an opioid (such as Oxycodone or Hydrocodone), you may use it for "break-through pain," which is pain that is not completely controlled with scheduled ibuprofen and acetaminophen. Do not take more than prescribed dose (usually one pill) in a 4-6 hour period.
- Note: some opioids contain acetaminophen! Check your label to see if your opioid medication contains acetaminophen and include that amount in the total daily intake. You should not take more than 4000 mg of acetaminophen in a 24 hour period.

Our nurse will call you to arrange for a postoperative visit. However, you can always contact us to schedule a visit sooner if you have concerns about your surgery.

Do not lift any objects over 10lbs for 6 weeks.

The stitches are absorbable in the skin and do not have to be removed. They will dissolve in 6-12 weeks. The glue on the incisions will come off in 1-2 weeks.

A small amount of clear or pink discharge from the incision can be normal. You can place a clean dry towel to absorb the drainage. If your incision itches, you can use a thin layer of KY jelly, petroleum jelly, olive oil, or A&D cream.

You are encouraged to shower daily, but avoid soaking in baths.

Nothing per vagina. No douching, tub baths, sexual intercourse, or tampons.

Do not submerge incision in water for 14 days after surgery. Showers are allowed anytime and should occur daily. Pat, do not rub, the incision.

You may eat or drink as usual. Plenty of fluids are encouraged.

To Prevent Constipation

You may have trouble with bowel movements after you go home. Drink plenty of liquids. Avoid caffeine drinks as they may dehydrate you. Being up and about is helpful as well. Narcotic pain pills will cause constipation. Take a stool softener (Docusate Sodium/Colace®) 100 mg twice daily and Miralax® 17 gm once daily while on narcotics. You can buy this without a prescription at the drugstore.

You can take ginger tea, mint tea, Gas-X, or Tums if you feel bloated. Cultured yogurt or probiotics can also help. Walking will also help.

Some swelling/bruising of the incision is normal. Call our gyn specialty nurses during business hours at (501) 526-7425/6990 if there is unusual chest pain, shortness of breath, significant redness/swelling/tenderness/drainage of the incision sites, fever greater than 100.4°F, persistent vomiting, uncontrolled pain, if you are not able to eat or drink liquids, vaginal bleeding more than 1 pad/hr or for any other concerns. After 4:30 pm, call ANGELS at (501) 526-7425 and ask for the gynecology resident on call.

Instructions (continued)

We cannot completely assess you over the phone. If you are concerned or feel it is an emergency, you should proceed to the nearest ER or UAMS.

COVID-19 VACCINATION RECORD

January 15, 2026

Berdia M. Monson Date of Birth: 5/1/1937

No record of COVID-19 immunization