

Trauma Assessment/Treatment Guidelines 705.01

- I. Purpose: To establish a consistent approach to the care of the trauma patient
 - A. Rapid trauma survey
 1. Airway
 - a. Maintain inline cervical stabilization
 - 1) Follow spinal motion restriction guidelines per VCEMS Policy 614
 - b. Open airway as needed
 - 2) Utilize a trauma jaw thrust to maintain inline cervical stabilization if indicated
 - c. Suction airway if indicated
 - d. Insert appropriate airway adjunct if indicated
 2. Breathing
 - a. Assess rate, depth, and quality of respirations
 - b. If respiratory effort inadequate, assist ventilations with BVM
 - c. Assess lung sounds
 - d. Initiate airway management and oxygen therapy as indicated
 - 1) Maintain SpO₂ ≥ 94%
 3. Circulation
 - a. Assess skin color, temperature, and condition
 - b. Check distal/central pulses and capillary refill time
 - c. Control major bleeding
 - d. Initiate shock management as indicated
 4. Disability
 - a. Determine level of consciousness (Glasgow Coma Scale)
 - b. Assess pupils
 5. Exposure
 - a. If indicated, remove clothing for proper assessment/treatment of injury location, always maintaining patient dignity
 - b. Always maintain patient body temperature
 - B. Detailed physical examination
 1. Head
 - a. Inspect/palpate skull
 - b. Inspect eyes, ears, nose and throat
 2. Neck
 - a. Palpate cervical spine
 - b. Check position of trachea
 - c. Assess for jugular vein distention (JVD)

Effective Date: April 10, 2025
Next Review Date: April 30, 2027

Date Revised: April 10, 2025
Last Reviewed: April 10, 2025



VCEMS Medical Director

3. Chest
 - a. Visualize, palpate, and auscultate chest wall
 4. Abdomen/Pelvis
 - a. Inspect/palpate abdomen
 - b. Assess pelvis, including genitalia/perineum if pertinent
 5. Extremities
 - a. Visualize, inspect, and palpate
 - b. Assess Circulation, Sensory, Motor (CSM)
 6. Back
 - a. Visualize, inspect, and palpate thoracic, and lumbar spines
- C. Trauma care guidelines
1. Fluid Administration
 - a. Maintain SBP of ≥ 90 mmHg
 - 1) Patients 65 years and older, maintain SBP of ≥ 100 mmHg
 - 2) Isolated head injuries, maintain SBP of ≥ 100 mmHg
 - b. Pediatric patients, maintain minimum SBP for respective age in Handtevy
 2. Transfusion of Blood Products
 - a. Warm and transfuse one unit (Approximately 500 mL) of whole blood or packed RBC when indications are met.
 - b. Repeat x1 to a total max of 2 units of blood transfused when indications continue to be met.
 - c. Inclusion Criteria
 - 1) Adult patient ≥ 14 Y.O.
 - 2) Patient consent obtained (informed or implied)
 - d. Indications
 - 1) Life threatening hemorrhage
 - 2) Vital sign criteria met (1 or more required)
 - a. SBP < 70 mmHg
 - b. SBP < 90 mmHg AND HR > 110
 - c. EMS witnessed traumatic cardiac arrest
 - e. Contraindications
 - 1) Ground level fall
 - 2) Isolated head injury
 - 3) Patient refusal
 - 4) Patient < 14 Y.O.
 - 5) Traumatic arrest not witnessed by EMS

3. Tranexamic Acid (TXA) Administration
 - a. As indicated in VCEMS Policy 734
3. Head injuries
 - a. General treatments
 - 1) Evaluate head and face – maintain high index of suspicion for injury if significant mechanism of injury is present or physical examination is remarkable for findings
 - 2) Elevate head 30° unless contraindicated
 - 3) Do not attempt to intubate head injured patients unless unable to manage with BLS airway measures
 - 4) Do not delay transport if significant airway compromise
 - b. Penetrating injuries
 - 1) DO NOT REMOVE IMPALED OBJECT (unless airway obstruction is present)
 - 2) Stabilize object manually or with bulky dressings
 - c. Facial injuries
 - 1) Assess airway and suction as needed
 - 2) Remove loose teeth or dentures if present
 - d. Eye injuries
 - 1) Remove contact lenses
 - 2) Irrigate eye thoroughly with suspected acid/alkali burns
 - 3) Avoid direct pressure
 - 4) Place eye shield over injured eye only
 - 5) Ask patient to keep eyes closed
 - 6) Stabilize any impaled object manually or with bulky dressing
4. Spinal cord injuries
 - a. General treatments
 - 1) Evaluate spinal column – maintain high index of suspicion for injury if significant mechanism of injury is present or physical examination is remarkable for findings
 - 2) Place patient in supine position if hypotension is present
 - b. Penetrating injuries – DO NOT REMOVE IMPALED OBJECT
 - 1) Stabilize object manually or with bulky dressings
 - 2) Control bleeding if present
 - 3) In the presence of isolated penetrating injuries, spinal motion restriction is contraindicated

- c. Neck injuries
 - 1) Monitor airway
 - 2) Control bleeding if present
- 5. Thoracic Trauma
 - a. General treatments
 - 1) Evaluate chest – maintain high index of suspicion for internal injury if significant mechanism of injury is present or physical examination is remarkable for findings
 - 2) Keep patients sitting high-fowlers
 - i. In the presence of isolated penetrating injuries, spinal motion restriction is CONTRAINDICATED
 - b. Penetrating injuries – DO NOT REMOVE IMPALED OBJECT
 - 1) Remove object if CPR is interfered
 - 2) Stabilize object manually or with bulky dressings
 - 3) Control bleeding if present
 - c. Flail Chest/Rib injuries
 - 1) Assist ventilations if respiratory status deteriorates
 - d. Pneumothorax/Hemothorax
 - 1) Keep patient sitting high-fowlers
 - 2) Assist ventilations if respiratory status deteriorates
 - 3) Suspected tension pneumothorax should be managed per VCEMS Policy 715
 - e. Open (Sucking) Chest Wound
 - 1) Place an occlusive dressing to wound site, secure on 3 sides only or place a vented chest seal.
 - 2) Assist ventilations if respiratory status deteriorates
 - f. Cardiac Tamponade – If suspected, expedite transport
 - 1) Beck's Triad
 - i. Muffled heart tones
 - ii. JVD
 - iii. Hypotension
 - g. Traumatic Aortic Disruption
 - 1) Assess for quality of radial and femoral pulses
 - 2) If suspected, expedite transport
- 6. Abdominal/Pelvic Trauma
 - a. General Treatments

- 1) Evaluate abdomen and pelvis – maintain high index of suspicion for internal injury if significant mechanism of injury is present or physical examination is remarkable for findings
- b. Blunt injuries
 - 1) Place patient in supine position if hypotension is present
- c. Penetrating injuries – DO NOT REMOVE IMPALED OBJECT
 - 1) Stabilize object manually or with bulky dressings
 - 2) Control bleeding if present
- d. Eviscerations
 - 1) DO NOT REPLACE ABDOMINAL CONTENTS
 - 2) Cover wound with saline-soaked dressings
 - 3) Control bleeding if present
- e. Pregnancy
 - 1) Place patient in left-lateral position to prevent supine hypotensive syndrome
- f. Pelvic injuries
 - 1) Assessment of pelvis should be only performed **ONCE** to limit additional injury
 - 2) Control external bleeding if present
 - 3) Place a commercial binder or sheet if pelvic injury is suspected and patient is hemodynamically unstable (see step one for parameters)
 - 4) Empirically place a binder or sheet if patient is in cardiac arrest due to a blunt or blast injury
 - 5) **Consider** applying a binder or sheet in patients with suspected pelvic injury **without** hemodynamic instability
7. Extremity Trauma
 - a. General Treatments
 - 1) Evaluate CSM distal to injury
 - i. If decrease or absence in CSM is present:
 - a) Attempt to reposition extremity into anatomical position
 - b) Re-evaluate CSM
 - c) If no change in CSM after repositioning, splint and expedite transport
 - d) Cover open wounds with sterile dressings
 - e) Place ice pack on injury area (if closed wound)
 - f) Splint/elevate extremity with appropriate equipment
 - b. Dislocations

- 1) Splint in position found with appropriate equipment
- c. Penetrating injuries – DO NOT REMOVE IMPALED OBJECTS
 - 1) Stabilize object manually or with bulky dressings
 - 2) Control bleeding if present
- d. Femur fractures
 - 1) Utilize traction splint only if isolated mid-shaft femur fracture is suspected
 - 2) Assess CSM before and after traction splint application
- e. Amputations
 - 1) Clean the amputated extremity with NS
 - 2) Wrap in moist sterile gauze
 - 3) Place in plastic bag
 - 4) Place bag with amputated extremity into a separate bag containing ice packs
 - 5) Prevent direct tissue contact with the ice pack