

Pain Control	
ADULT	PEDIATRIC
BLS Procedures	
Administer oxygen as indicated Assess numerical or faces pain scale Control pain using basic measures such as patient positioning, ice packs, and splinting	
ALS Standing Orders	
IV/IO access Cardiac Monitor <u>Mild to Moderate Pain</u> Acetaminophen <ul style="list-style-type: none">IV/IO – 1g in 100 mL NS over 15 minutes. <u>Severe Pain</u> - Utilize opiate analgesic in addition to, or in place of, acetaminophen. Fentanyl <ul style="list-style-type: none">IM/IN - 1 mcg/kg, Max 100 mcg<ul style="list-style-type: none">Repeat q 10 minutes for persistent pain to a max total dose of 200 mcg.IV/IO - 1 mcg/kg over 1 minute, Max 100 mcg<ul style="list-style-type: none">Repeat q 5 minutes for persistent pain to a max total dose of 200 mcg.Repeat doses should be administered IV/IO if vascular access obtained.Withhold for systolic < 90 mmHg. If Fentanyl unavailable; Ondansetron - Per 705.15 Nausea/Vomiting Policy <ul style="list-style-type: none">Repeat x 1 in 10 minutes for nausea or > 2 doses of Morphine Morphine <ul style="list-style-type: none">IM - 0.1 mg/kg, Max 10 mg<ul style="list-style-type: none">Repeat ½ initial dose q 10 minutes x 2 for persistent pain.IV/IO - 0.1 mg/kg, Max 10 mg, over 1 minute<ul style="list-style-type: none">Repeat ½ initial dose q 5 minutes x 2 for persistent pain.	IV/IO access Cardiac Monitor <u>Mild to Moderate Pain</u> Acetaminophen <ul style="list-style-type: none">IV/IO – 15 mg/kg in 100 mL NS over 15 minutes. <u>Severe Pain</u> – Utilize opiate analgesic in addition to, or in place of, acetaminophen. Fentanyl <ul style="list-style-type: none">IM/IN - 1 mcg/kg, Max 100 mcg<ul style="list-style-type: none">Repeat q 10 minutes for persistent pain to a max total dose of 200 mcg.IV/IO - 1 mcg/kg over 1 minute, Max 100 mcg<ul style="list-style-type: none">Repeat q 5 minutes for persistent pain to a max total dose of 200 mcg.Repeat doses should be administered IV/IO if vascular access obtained.Withhold for systolic less than minimum SBP in Handtevy for respective age. If Fentanyl unavailable; Ondansetron - Per 705.15 Nausea/Vomiting Policy <ul style="list-style-type: none">Repeat x 1 in 10 minutes for nausea or > 2 doses of Morphine Morphine <ul style="list-style-type: none">IM - 0.1 mg/kg, Max 10 mg<ul style="list-style-type: none">Repeat ½ initial dose q 10 minutes x 2 for persistent pain.IV/IO - 0.1 mg/kg, Max 10 mg, over 1 minute<ul style="list-style-type: none">Repeat ½ initial dose q 5 minutes x 2 for persistent pain.
Base Hospital Orders only	
Consult with ED Physician for further treatment measures	