

21-701: Discrete Mathematics

Lecturer: Wes Pegden

Scribe: Sidharth Hariharan

Carnegie Mellon University - Fall 2026

This version of the notes was last compiled on August 26, 2026.

For the latest version, visit [https:](https://thefundamentaltheor3m.github.io/DiscreteMathNotes/main.pdf)

[//thefundamentaltheor3m.github.io/DiscreteMathNotes/main.pdf](https://thefundamentaltheor3m.github.io/DiscreteMathNotes/main.pdf).

For comments, errors or suggestions, please raise an issue/make a pull request at

<https://github.com/thefundamentaltheor3m/DiscreteMathNotes>.

Contents

1	Graphs and Colorings	2
1.1	Colorings	3
1.1.1	2-Colorings of Graphs	3
1.1.2	Colorings of Hypergraphs	4
1.1.3	Bounds on $m(k)$	7
1.1.4	Maker-Breaker Games	8
1.2	Ramsey Numbers	10
	Bibliography	12

Chapter 1

Graphs and Colorings

Much of this course will be about graphs, and a surprising amount of what we want to know about a graph can be phrased as a question about coloring it. We begin with the vocabulary, and then settle the first such question completely: which graphs admit a coloring with only two colors. We then ask the same question of hypergraphs, where it becomes considerably harder—even pinning down how many edges a 3-uniform hypergraph needs before two colors are no longer enough takes real work. Turning the question around—how large must a graph be before every coloring of it is forced to produce something monochromatic—gives us the Ramsey numbers, which close the chapter.

Notation.

- $[k]$ will denote $\{1, \dots, k\}$.
- $\sim$ will denote the adjacency relation on graph vertices.
- K_n will denote the complete graph on n vertices.


1.1 Colorings

Definition 1.1.1 (Coloring). Given a graph $G = (V, E)$, a k -coloring of G is a map $c : V \rightarrow [k]$ such that $u \sim v \implies c(u) \neq c(v)$.

1.1.1 2-Colorings of Graphs

Let's start with a motivating question: which graphs can be 2-colored?

Any odd cycle graph can't be 2-colored. That is, any graph that looks like



Let's study when odd cycles exist.

Lemma 1.1.2. *If G has an odd closed walk, then G has an odd cycle.*

Recall that a closed walk need not be a cycle: walks can contain repetitions. So even line graphs can have closed walks (start from one vertex, go to its neighbor, return).

Proof. Suppose G has an odd closed walk. Let W be such a walk of minimal length. Write $W = (x_1, \dots, x_i, \dots, x_j, \dots, x_\ell)$, with $x_\ell = x_1$. Suppose W is not a cycle. Then there must be some x_i, x_j such that $x_i = x_j$ even though $i \neq j$, ie, there must be some repetition. Then, $(x_1, \dots, x_i, x_{j+1}, \dots, x_\ell)$ and $(x_i, x_{i+1}, \dots, x_j)$ are both shorter walks than W . Moreover, one of them must be of odd length. This contradicts minimality of the length of W . So W must be a cycle. □

Here's a cool result.

Theorem 1.1.3. *A graph $G = (V, E)$ can be 2-colored if and only if G has no odd cycle.*

Proof.

($\implies$) We've already seen that odd cycles can't be 2-colored. So if G can be 2-colored, it can't have an odd cycle. We've essentially seen this from the examples above.

($\impliedby$) We may assume G is connected, since otherwise we can color each component separately.

Fix some $v_0 \in V$. Define $c : V \rightarrow \{1, 2\}$ by

$$c(u) = \begin{cases} 2 & \text{if } \text{dist}(v_0, u) \text{ is odd} \\ 1 & \text{if } \text{dist}(v_0, u) \text{ is even} \end{cases}$$

where $\text{dist}(\cdot, \cdot)$ denotes the shortest number of edges from one vertex to another.

We claim that this c is a valid 2-coloring.

Indeed, suppose it is not. Then, there must be some $u, v \in V$ such that $u \sim v$ and $c(u) = c(v)$. Now let's consider the two possible cases on the color.

Suppose $c(u) = c(v) = 1$ for adjacent u, v . Then, there is a path of even length between v and v_0 and another path of even length between u and v_0 . Concatenating the path from v to v_0 with the one from v_0 to u , and closing up with the edge uv , we get a closed walk of length $\text{even} + \text{even} + 1$. This is odd, so by Theorem 1.1.2, G has an odd cycle.

The case $c(u) = c(v) = 2$ is identical: the two paths are now of odd length, so the closed walk they form with the edge uv has length $\text{odd} + \text{odd} + 1$, which is odd again.

Either way, G has an odd cycle, contrary to hypothesis. So c must be a valid 2-coloring.

□

1.1.2 Colorings of Hypergraphs

Coloring is not really a question about graphs. All the definition above asks of G is that we know which pairs of vertices may not share a color, and nothing stops us forbidding larger sets instead.

Definition 1.1.4 (Hypergraph). Given a vertex set V , a **hypergraph** is a collection H of subsets of V . The sets in H are called **hyperedges**.

Definition 1.1.5 (Uniformity). A hypergraph H is k -uniform if $|f| = k$ for all $f \in H$.

The following is thus obvious.

Proposition 1.1.6. A graph is a 2-uniform hypergraph.

Definition 1.1.7. A proper coloring of a hypergraph H is an assignment $c : V \rightarrow [r]$ such that $\forall f \in H, |c(f)| \geq 2$.

We say “no edge is monochromatic”.

We can ask ourselves the following question: “Which 3-uniform hypergraphs are 2-colorable?”

For each k , define $m(k)$ to be the smallest m such that $\exists$ some k -uniform hypergraph with m edges that is not 2-colorable. We can show that $m(2) = 3$. But what is $m(3)$? That is, how many 3-edges do I need to prevent 2-colorability?

Let H be a 3-uniform hypergraph that's not 2-colorable. Assume, further, that H has at most 6 edges (each being a subset of size 3).

First, we'll show that we can reduce, without loss of generality, to the case where H has exactly 6 vertices. Indeed, if H has vertices x, y that are not in any common edge, then I can define H' by “merging” x and y . So H' has one fewer vertex than H . This H' cannot be 2-colorable: if it were, then that coloring would extend to H by using the color of the merged vertex at both x and y , which would make H 2-colorable. And when H has more than 6 vertices, such a pair always exists.

An edge has 3 vertices, so it covers the $\binom{3}{2} = 3$ pairs among them, and H has at most 6 edges, so at most

$$6 \cdot \binom{3}{2} = 18$$

pairs are covered in total. But if H has $n \geq 7$ vertices, there are $\binom{n}{2} \geq \binom{7}{2} = 21$ pairs to cover. So some pair is left uncovered, and merging it brings us one vertex closer to 6.

So now assume that H has exactly 6 edges and 6 vertices and is not 2-colorable.


There is no loss in the vertex count in the other direction either: a hypergraph with fewer than 6 vertices can be handed as many extra vertices as it likes, and vertices lying in no edge change nothing about which colorings are proper.

Now count colorings. Among all the ways of coloring 6 vertices with two colors, consider only the balanced ones, which split V into two sets of three; there are $\frac{1}{2}\binom{6}{3} = 10$ of these. Such a coloring fails exactly when some edge is monochromatic, and an edge of H has three vertices, so it is monochromatic under a balanced coloring precisely when it is one of the two color classes. Each edge f therefore rules out exactly one of the ten balanced colorings, namely the one splitting V into f and $V \setminus f$. Six edges rule out at most six of the ten, so at least four survive, and any one of them is a proper 2-coloring of H . This contradicts our assumption, so no such H exists and

$$m(3) \geq 7$$

That bound is attained, and the hypergraph attaining it is a familiar object.

Example 1.1.8 (The Fano Plane). The **Fano plane** is the 3-uniform hypergraph on the seven vertices below, whose seven edges—by convention, its **lines**—are the three sides of the triangle, the three medians, and the circle through the three edge midpoints.



Every vertex lies on exactly three lines, any two vertices lie on exactly one line together, and any two lines meet in exactly one vertex. That last property is what does the work below.

Proposition 1.1.9. *The Fano plane is not 2-colorable. Hence $m(3) = 7$.*

Proof. Of the $\binom{7}{4} = 35$ sets of four vertices, exactly 28 contain a line: a line together with any one of the four vertices off it gives such a set, which is $7 \cdot 4 = 28$ sets in all, and no two of them coincide because four vertices cannot contain two lines—two lines meet in a vertex, so between

them they cover five. The seven sets left over are the complements of the seven lines.

Now suppose we had a proper 2-coloring, with color classes A and B , and say $|A| \leq 3$. If $|A| = 3$, then A is not a line, since A is monochromatic; so $B = V \setminus A$ is a set of four vertices that is not the complement of a line, and hence contains one. If $|A| \leq 2$, then the vertices outside B lie on at most $3 + 3 - 1 = 5$ lines between them, so at least two lines avoid them both and lie inside B . Either way some line is monochromatic, which is a contradiction.

So the Fano plane is a 3-uniform hypergraph with seven edges that is not 2-colorable, giving $m(3) \leq 7$. $\square$

1.1.3 Bounds on $m(k)$

Pinning down $m(3)$ took a page of case analysis, and doing the same for $m(4)$ is out of the question. What we can do for every k at once is bound it, and the lower bound falls out of an idea that will keep coming back: color at random, and show that a bad coloring is improbable.

Theorem 1.1.10. $m(k) \geq 2^{k-1}$. In other words, any k -uniform hypergraph with $< 2^{k-1}$ edges is 2-colorable.

Proof. Given such a hypergraph, randomly color the vertices red and blue, with equal probability ($1/2$) for each. For each edge f , let $\mathcal{E}_f$ be the event that f is monochromatic. Then,

$$\Pr(\mathcal{E}_f) = \frac{2}{2^k}$$

Taking a simple union bound,

$$\Pr\left(\bigcup_{f \in H} \mathcal{E}_f\right) \leq \frac{|H|}{2^{k-1}} < 1$$

where we interpret $\bigcup_{f \in H} \mathcal{E}_f$ as the event that there is a monochromatic edge. As the probability is less than 1, there must be at least one coloring with no monochromatic edge. $\square$

Note that the above argument can also be written combinatorially instead of probabilistically. In this instance, the probabilistic language essentially conveys the same information as combinatorial language would. But this may not always be true, so it's worth being careful about this.

In the other direction, the truth is only a factor of about k^2 away.

Exercise 1.1.11. Prove that $m(k) = O(k^2 \cdot 2^k)$

1.1.4 Maker-Breaker Games

Theorem 1.1.10 says that a hypergraph with few enough edges has a proper 2-coloring somewhere among all the colorings of its vertices. It turns out that the very same threshold survives a far more demanding requirement: that one of two players be able to build such a coloring one vertex at a time, against an opponent trying to stop them.

Here's a fun fact: a "maker-breaker" game is a game where the "maker" wins if they are able to achieve an objective and the "breaker" wins if they are able to stop the maker from winning. So if you set up a "maker-breaker" version of tic-tac-toe, where only the crosses can win by making three in a row and the noughts can only win by stopping the crosses, the crosses will always win (assuming the crosses play first). This is, of course, different from tic-tac-toe, which, as our own studies from when we were a kid have shown, can always be drawn.

The threshold at which breaker can hold out is exactly the one we have just been looking at.

Theorem 1.1.12 (Erdős-Selfridge). *If H is a k -uniform hypergraph with m edges, and $m < 2^{k-1}$, then breaker wins (if breaker goes second).*

In the language of hypergraphs, the game is 2-coloring; maker winning is equivalent to the existence of a coloring with a monochromatic edge; and breaker winning is equivalent to the inexistence of such edge.

Proof. Let Δ be the maximum degree of H , that is,

$$\Delta = \max_{v \in V} |\{f \in H \mid v \in f\}|$$

We'll prove that if $m + \Delta < 2^k$, then breaker will win. Then, since $\Delta \leq m$, it will follow that if $m < 2^{k-1}$, then $m + \Delta \leq 2m < 2^k$.

At the start of move i , we will conceptualize the danger of edge $f \in H$ by

$$\phi_i(f) = \begin{cases} 2^{\text{number of maker moves in } f} & \text{if there are no breaker moves in } f \\ 0 & \text{if there is a breaker move in } f \end{cases}$$

$$\Phi_i = \sum_{f \in H} \phi_i(f)$$

Consider the strategy where, once maker has played y on move i , breaker plays the unclaimed vertex carrying the most danger, that is, the x maximizing

$$\sum_{f \ni x} \phi(f)$$

with ϕ measured after maker's move. Two things move Φ on move i . Maker's play doubles the danger of every edge through y that breaker has not already killed, so it raises Φ by $a_i = \sum_{f \ni y} \phi_i(f)$; breaker's play sends the danger of every surviving edge through x to 0, so it lowers Φ by the quantity b_i that breaker has just maximized. Hence

$$\Phi_{i+1} = \Phi_i + a_i - b_i$$

The point of breaker's rule is that $a_i \leq b_{i-1}$ for every $i \geq 2$. Indeed, y is unclaimed right up until maker takes it on move i , so it was one of the vertices breaker was choosing between on move $i - 1$, and breaker chose one carrying at least as much danger as it. Between then and the start of move i nothing happens except breaker's own play, which only ever sends dangers to 0, so the danger through y at the start of move i is no greater than it was when breaker passed y over.

Summing $\Phi_{i+1} = \Phi_i + a_i - b_i$ over the first n moves and canceling each a_i against b_{i-1} , everything past the first two terms disappears and we are left with

$$\Phi_{n+1} \leq \Phi_1 + a_1$$

Stopping the same sum one term earlier gives the same bound halfway through a move, after maker has played and before breaker has replied. Now $\Phi_1 = m$, since at the start every edge contains a move of neither player and so contributes $2^0 = 1$, and a_1 is the number of edges through maker's opening vertex, which is at most Δ . So the danger never exceeds $m + \Delta$ at any point in the game.

But if maker ever completes an edge f , then at that moment f contains all k of its vertices from

maker and none from breaker, so $\phi(f) = 2^k$ and hence $\Phi \geq 2^k$. Since $m + \Delta < 2^k$, this never happens, and breaker wins. $\square$

Let's consider higher-dimensional maker-breaker tic-tac-toe.

Example 1.1.13. Turns out that $8 \times 8 \times 8$ maker-breaker tic-tac-toe is breaker-winnable: we can show that $m = \frac{10^3 - 8^3}{2} = 244$ and $\Delta = 7$, meaning $m + \Delta = 251 < 2^k = 256$.

Both numbers come out of the geometry. For m , pad the board with one extra cell in every direction, giving a $10 \times 10 \times 10$ cube, and run each winning line one step past each of its two ends. The two cells we land on lie in the shell we just added, and every cell of the shell is the end of exactly one line: a coordinate of the cell equal to 0 tells the line to increase along that axis, one equal to 9 tells it to decrease, and the rest stay fixed. So the lines are in bijection with the pairs of shell cells, of which there are $\frac{10^3 - 8^3}{2}$. For Δ , a line through a given cell is determined by its direction, and a corner cell has the 3 lines along the axes, the 3 face diagonals and the one long diagonal through it, which is 7; no cell of an $8 \times 8 \times 8$ board does better. And it is the $m + \Delta < 2^k$ form of the criterion doing the work here, since 244 is a long way above $2^{k-1} = 128$.


We won't say any more about tic-tac-toe in this course, but it was a fun aside.

1.2 Ramsey Numbers

Everything so far has asked when a coloring exists. The opposite question is at least as natural: how large does a structure have to be before every coloring of it is forced to contain something monochromatic? Let's step back into "graph land" from "hypergraph land" and ask it of the complete graphs, coloring edges this time rather than vertices.

Definition 1.2.1 (Ramsey Number). The k th Ramsey number, denoted $R(k)$, is the smallest $n \in \mathbb{N}$ such that any 2-coloring of edges of K_n gives a monochromatic K_k .

One immediate result we can see is that $R(3) \geq 6$, which we can see simply by drawing a picture of K_5 and studying what happens if we move our attention to K_6 .



Each color carries a 5-cycle—the outer pentagon in red, the inner pentagram in blue—and a 5-cycle contains no triangle at all, so neither color gives us a monochromatic K_3 . Nothing forces a monochromatic triangle at $n = 5$, and so $R(3) \geq 6$. The picture also shows how little room there is to spare: every vertex of K_5 meets exactly two edges of each color, and the moment we pass to K_6 each vertex has five edges to distribute between two colors, so one of the colors is forced to take three of them.

We can actually prove that $R(3) = 6$, but this is rather nontrivial.

Other known Ramsey numbers are $R(2) = 2$ and $R(4) = 18$. The rest are unknown, though we do have bounds for them. The best (or close to the best) known bounds are

$$2^{\frac{k}{2}} \leq R(k) \leq 3.8^k$$

Bibliography

These lecture notes are based heavily on the following references:

- [1] Nicolas Bourbaki. *Groupes et Algèbres de Lie. Chapitre 1*. 1st ed. Éléments de Mathématique. Édition originale publiée par Hermann, Paris, 1972. Springer Berlin, Heidelberg, 2007. ISBN: 978-3-540-35337-9. DOI: <https://doi.org/10.1007/978-3-540-35337-9>.

Visit <https://thefundamentaltheor3m.github.io/DiscreteMathNotes/main.pdf> for the latest version of these notes. If you have any suggestions or corrections, please feel free to fork and make a pull request to the [associated GitHub repository](#).