

MATH70062: Lie Algebras

Lecturer: Ambrus Pal

Scribe: Sidharth Hariharan

Imperial College London - Autumn 2024

Contents

1	An Introduction to the Theory of Lie Algebras	2
1.1	Important Definitions and First Examples	2
1.1.1	Algebras	2
1.1.2	Subalgebras and Homomorphisms	5
1.1.3	Ideals and Quotients	8
1.1.4	Adjoint and Derivations	9

Chapter 1

An Introduction to the Theory of Lie Algebras

While many of the definitions and constructions we shall see in this course can easily be adapted to any field, we will work over \mathbb{C} for simplicity, unless otherwise stated.

1.1 Important Definitions and First Examples

We will begin by defining the fundamental objects of study in this course. We will then provide some examples of these objects and discuss means of constructing them.

1.1.1 Algebras

We begin by recalling the notion of a bilinear map.

Definition 1.1.1 (Bilinear Map). Let V and W be vector spaces. We say that a map $f : V \times W \rightarrow \mathbb{C}$ is **bilinear** if it is linear in each argument. That is, for all $v, v' \in V$, $w, w' \in W$ and $\lambda \in \mathbb{C}$, we have

$$f(v + v', w) = f(v, w) + f(v', w)$$

$$f(v, w + w') = f(v, w) + f(v, w')$$

$$f(\lambda v, w) = \lambda f(v, w) = f(v, \lambda w)$$

We will be particularly interested in bilinear maps from a vector space to itself.

Definition 1.1.2 (Algebra). An **algebra** is a vector space A equipped with a bilinear map $\cdot : A \times A \rightarrow A$.

Convention. Given any algebra A , we will often refer to the corresponding bilinear map \cdot as the **multiplication** map of A , and denote $\cdot(x, y)$ as simply $x \cdot y$ or even xy (where the definition of \cdot is clear from the context) for any $x, y \in A$.

There are many different kinds of algebras. We will be particularly interested in Lie algebras and associative algebras.

Definition 1.1.3 (Associative Algebras). We say that an algebra A is **associative** if the multiplication map \cdot is associative. That is, for all $x, y, z \in A$, we have

$$(x \cdot y) \cdot z = x \cdot (y \cdot z)$$

We have all seen associative algebras before.

Example 1.1.4 (The Matrix Algebra). The set $M_n(\mathbb{C})$ of $n \times n$ matrices over \mathbb{C} forms an associative algebra under matrix multiplication, known as the Matrix Algebra.

We will come back to associative algebras soon enough. We will now define the main object of study in this module.

Definition 1.1.5 (Lie Algebras). A **Lie algebra** is an algebra L whose bilinear map $[\cdot, \cdot] : L \times L \rightarrow L$ satisfies the following properties:

1. For all $x \in L$, we have $[x, x] = 0$.
2. For all $x, y, z \in L$, we have

$$[x, [y, z]] + [y, [z, x]] + [z, [x, y]] = 0 \quad (1.1.1)$$

Such a bilinear map $[\cdot, \cdot]$ is known as a **Lie Bracket**, and (??) is known as the **Jacobi**

Identity.

Remark. We immediately notice that the first condition (over not just \mathbb{C} but any field) implies the fact that

$$[x, y] = -[y, x] \quad (1.1.2)$$

One simply needs to apply bilinearity and the first condition to evaluate $[x + y, x + y]$. This argument reverses nicely as well, but only over fields of characteristic $\neq 2$.

One may recall that the $[\cdot, \cdot]$ notation is often used in group theory to denote the **commutator** of two elements. The reason why the same notation is used for the Lie bracket is the following.

Lemma 1.1.6. *Let A be an associative algebra. Then, the commutator map $[x, y] = xy - yx$ is a Lie bracket on A .*

Proof. Clearly, $[x, x] = xx - xx = 0$ for all $x \in A$. We now show that $[\cdot, \cdot]$ satisfies (1.1.1): for all $x, y, z \in A$, we have

$$\begin{aligned} [x, [y, z]] + [y, [z, x]] + [z, [x, y]] &= [x, yz - zy] + [y, zx - xz] + [z, xy - yx] \\ &= 6xyz - 6xyz = 0 \end{aligned}$$

where we skip over some of the intermediate computations because they are tedious and uninteresting. □

Lemma 1.1.6 gives us a large class of examples of Lie algebras. One of the most important of these is the following.

Example 1.1.7 (General Linear Lie Algebra). For all $n \in \mathbb{N}$, the set of all $n \times n$ matrices forms a Lie algebra under the commutator bracket: this follows immediately from applying Lemma 1.1.6 to Example 1.1.4. We call this the **General Linear Lie Algebra**, denoted $\mathfrak{gl}(n)$.

Convention. We will denote by $M_n(\mathbb{C})$ the set of all $n \times n$ matrices, viewed (interchangeably) as a *set*, a *vector space* or an *associative algebra*. When viewing it as a *Lie algebra under the commutator bracket*, we will adopt the notation $\mathfrak{gl}(n, \mathbb{C})$, where \mathbb{C} can be replaced by any field. We will usually abbreviate this to $\mathfrak{gl}(n)$, because we will primarily work over \mathbb{C} .

Lastly, we will define the notion of an abelian Lie algebra.

Definition 1.1.8 (Abelian Lie Algebra). A Lie algebra A is said to be **abelian** if for all $x \in A$, we have $[x, x] = 0$.

The reason for this terminology is that if A is an associative algebra whose multiplication map is commutative, then its commutator bracket is identically zero, making the corresponding Lie algebra abelian.

Example 1.1.9. Clearly, $\mathfrak{gl}(1)$ is abelian: for all $x, y \in \mathfrak{gl}(1) = \mathbb{C}$, we have $xy - yx = 0$.

We will now define subalgebras and homomorphisms of algebras, which will allow us to construct more examples of algebras (Lie and otherwise).

1.1.2 Subalgebras and Homomorphisms

As with objects in any category, we have subobjects and morphisms. We will define these over general algebras and apply them to get more examples of Lie algebras.

Definition 1.1.10 (Subalgebras). Let A be a vector space. A **subalgebra** of A is a subspace $B \subseteq A$ such that B is closed under the multiplication map of A . That is, for all $x, y \in B$, we have $x \cdot y \in B$.

Convention. Given an algebra A and a subset $B \subseteq A$, we will denote the statement that B is a subalgebra of A by $B \leq A$.

Definition 1.1.11 (Homomorphisms). Let A and B be algebras. A **homomorphism** $\phi : A \rightarrow B$ is a linear map that respects the multiplication maps of A and B . That is, for all $x, y \in A$, we have

$$\phi(x \cdot y) = \phi(x) \cdot \phi(y)$$

Convention. We will define Lie subalgebras to be subalgebras with respect to the algebra structure given by the Lie bracket, and we will define Lie algebra homomorphisms to be homomorphisms that respect the Lie bracket (ie, that are algebra homomorphisms with respect to the algebra structure given by the Lie bracket).

We have the following unsurprising result.

Lemma 1.1.12. *Let A and B be algebras, and let $\phi : A \rightarrow B$ be a homomorphism. Then,*

1. $\text{im}(\phi) \leq B$
2. $\text{ker}(\phi) \leq A$

Proof. These are standard results, but we will prove them for completeness.

1. Fix $x, y \in \text{im}(\phi)$. Then, there exist $a, b \in A$ such that $\phi(a) = x$ and $\phi(b) = y$. Since ϕ is a homomorphism, we have

$$x \cdot y = \phi(a) \cdot \phi(b) = \phi(a \cdot b) \in \text{im}(\phi)$$

so $\text{im}(\phi)$ is closed under the multiplication map of B .

2. Let $x, y \in \text{ker}(\phi)$. Then, we have

$$\phi(x \cdot y) = \phi(x) \cdot \phi(y) = 0 \cdot 0 = 0$$

where the last equality follows from the fact that \cdot is bilinear. Therefore, $x \cdot y \in \text{ker}(\phi)$, and $\text{ker}(\phi)$ is closed under the multiplication map of A .

□

This allows us to construct another matrix Lie algebra.

Example 1.1.13 (The Special Linear Lie Algebra). For all $n \in \mathbb{N}$, consider the trace map $\text{Tr} : \mathfrak{gl}(n) \rightarrow \mathfrak{gl}(1)$. This is a (Lie) algebra homomorphism: for all $A, B \in \mathfrak{gl}(n)$,

$$\text{Tr}([A, B]) = \text{Tr}(AB - BA) = \text{Tr}(AB) - \text{Tr}(BA) = 0 = [\text{Tr}(A), \text{Tr}(B)]$$

because the Lie algebra $\mathfrak{gl}(1)$ is abelian (see Example 1.1.9). By Lemma 1.1.12, its kernel, the set of all $n \times n$ matrices of trace zero, is a Lie subalgebra of $\mathfrak{gl}(n)$. We call this the **Special Linear Lie Algebra**, denoted $\mathfrak{sl}(n)$.

Remark. In Example 1.1.13, we have indirectly shown that

$$\text{im}([\cdot, \cdot]) = [\mathfrak{gl}(n), \mathfrak{gl}(n)] \subseteq \mathfrak{sl}(n)$$

because of the unique property of the trace that $\text{Tr}(AB) = \text{Tr}(BA)$ for any $A, B \in \mathfrak{gl}(n)$.

The very natural relationship between associative and Lie algebra structures given by Lemma 1.1.6 gives us an elegant criterion for proving that a subspace is a subalgebra of a Lie algebra whose Lie bracket is the commutator of an associative bilinear map.

Proposition 1.1.14. *Let (A, \cdot_A) be an associative algebra and let (B, \cdot_B) be a subalgebra of A . Denoting by $(A, [\cdot, \cdot]_A)$ the Lie algebra whose Lie bracket is the commutator of the multiplication map of A and by $(B, [\cdot, \cdot]_B)$ the Lie algebra whose Lie bracket is the commutator of the multiplication map of B , we have $B' \leq A'$. In other words, the following diagram commutes:*

$$\begin{array}{ccc} (A, \cdot_A) & \longrightarrow & (A, [\cdot, \cdot]_A) \\ \uparrow \text{Associative Subalgebra} & & \uparrow \text{Lie Subalgebra} \\ (B, \cdot_B) & \longrightarrow & (B, [\cdot, \cdot]_B) \end{array} \quad (1.1.3)$$

Proof. First, observe that $[\cdot, \cdot]_B = [\cdot, \cdot]_A|_B$ (ie, the Lie bracket obtained from \cdot_B agrees with the

one obtained from \cdot_A on B): for all $T_1, T_2 \in B$,

$$[T_1, T_2]_B = T_1 \cdot_B T_2 - T_2 \cdot_B T_1 = T_1 \cdot_A T_2 - T_2 \cdot_A T_1 = [T_1, T_2]_A$$

Therefore, since B is closed under $[\cdot, \cdot]_B$ (which, by definition, is a map from $B \times B$ to B), B must be closed under $[\cdot, \cdot]_A$. \square

This allows us to construct more examples still.

Example 1.1.15 (The Upper-Triangular Lie Algebra). For $n \in \mathbb{N}$, we define the **Upper-Triangular Lie Algebra** to be the set of all $n \times n$ upper-triangular matrices (with respect to some predetermined basis), denoted $\mathfrak{t}(n)$. Given that the product of upper-triangular matrices is upper-triangular, $\mathfrak{t}(n)$ forms an associative subalgebra of $M_n(\mathbb{C})$, and therefore, a Lie subalgebra of $\mathfrak{gl}(n)$.

1.1.3 Ideals and Quotients

Throughout this subsection, we will denote by L an arbitrary Lie algebra.

Definition 1.1.16 (Ideal). We say that $I \subseteq L$ is an **ideal** of L , denoted $I \trianglelefteq L$, if I is a linear subspace of L and $[x, y] \in I$ for all $x \in L$ and $y \in I$.

Convention. We will use the notation $[I, L]$ to denote the subspace of $\mathfrak{gl}(n)$ spanned by all elements of the form $[i, \ell]$ for $i \in I$ and $\ell \in L$.

Remark. We could equivalently require that $[I, L] \leq L$ in the definition of an ideal instead of requiring that $[x, y] \in I$ for all $x \in L$ and $y \in I$. Similarly, we can observe that it doesn't matter whether we require $[x, y] \in I$ or $[y, x] \in I$ because of (1.1.2) and bilinearity.

Example 1.1.17 (Trivial Ideals). Given any Lie algebra L , both $\{0\}$ and L are ideals of L .

Lemma 1.1.18. Given a Lie algebra homomorphism $\phi : L \rightarrow K$, $\ker(\phi) \trianglelefteq L$.

Proof. sorry □

Definition 1.1.19 (The Centre of a Lie Algebra). We define the **centre** of L to be

$$Z(L) := \{x \in L \mid \forall y \in L, [x, y] = 0\}$$

Lemma 1.1.20. $Z(L)$ is an ideal of L .

Proof. sorry □

Example 1.1.21. For all $n \in \mathbb{N}$,

$$Z(\mathfrak{gl}(n)) = \{A \in \mathfrak{gl}(n) \mid \exists \lambda \in \mathbb{C} \text{ s.t. } A = \lambda I\}$$

Proof. sorry □

For the remainder of this subsection, let I be an arbitrary ideal of L .

Proposition 1.1.22. Consider the vector space L/I . The map $[\cdot, \cdot] : L/I \times L/I \rightarrow L/I$ given by

$$[x + I, y + I] := [x, y] + I$$

for all $x, y \in L$ is a Lie bracket on L/I .

Proof. We primarily need to show that the Lie bracket on L/I is well-defined. It would then suffice to prove the much easier fact that the quotient map $x \mapsto x + I : L \twoheadrightarrow L/I$ is a Lie algebra homomorphism. sorry □

Definition 1.1.23 (Quotient Algebra).

1.1.4 Adjoints and Derivations

Throughout this subsection, V will refer to a finite-dimensional vector space.

We begin with a general Lie algebra construction.

Definition 1.1.24 (General Linear Lie Algebra over an Arbitrary Vector Space). We define the **General Linear Lie Algebra over V** to be the set of all linear maps from V to V , viewed as a Lie algebra under the commutator bracket. We denote it $\mathfrak{gl}(V)$.

That this is, indeed, a Lie algebra should come as no surprise. Given that this construction is well-defined over *any* vector space, we can, in particular, apply it to Lie algebras.

Let L denote an arbitrary Lie algebra. There exists a rather nice map that relates L with $\mathfrak{gl}(L)$: the adjoint.

Definition 1.1.25 (Adjoint Map). To every $x \in L$, we can associate the linear map

$$\mathrm{ad}(x) : L \rightarrow L : y \mapsto [x, y]$$

We call this map the **adjoint map** associated to x .