

21-603: Model Theory I

Lecturer: Rami Grossberg

Scribes: Sidharth Hariharan and Teresa Pollard

Carnegie Mellon University - Fall 2025

Contents

1	An Introduction to Model Theory	2
1.1	A Crash Course on First-Order Logic	2
1.1.1	Languages and Structures	2
1.1.2	Syntax: Terms, Formulae, Sentences and Theories	4
1.1.3	Semantics: Satisfaction	6
1.1.4	Elementary Equivalence	7
1.1.5	Deduction and Proof	10
1.2	Cardinality and Categoricity	11
1.2.1	The Spectrum Problem	11
1.2.2	Cardinal Arithmetic	12
1.3	A Word on Submodels	14
1.3.1	Submodel Existence	14
1.3.2	Elementary Submodels	15
1.3.3	Chains of Substructures	15
1.3.4	The Löwenheim-Skolem Theorems	16
1.4	Models of Peano Arithmetic	16
1.4.1	Restrictions and Expansions	16
1.4.2	The Language and Theory of Peano Arithmetic	17
1.4.3	Existence of Non-Standard Models of Peano Arithmetic	17
1.4.4	Countable Models of Peano Arithmetic	18
1.4.5	Impossible Deductions	18
2	Another Chapter	20

2.1	Introducing the Main Object of Study in this Chapter	20
2.2	Another Section	20

Bibliography		23
---------------------	--	-----------

Chapter 1

An Introduction to Model Theory

Model Theory is one of the two main branches of mathematical logic, alongside Set Theory. We can view Model Theory as a translation of algebra to the world of set theory, or as a *generalisation* of algebra, with many applications to algebra. Indeed, it is possible to view model theory as a more specialised version of category theory, at least in its power to deal with generality. We can prove tremendously deep theorems about algebra, or other fields of maths, purely using model theory.

Before we can talk about model theory in any detail, we need to recall a few notions from the world of logic.

1.1 A Crash Course on First-Order Logic

1.1.1 Languages and Structures

We begin by recalling the notion of a first-order language.

Definition 1.1.1 (Language). A **language** is a disjoint union

$$\mathbf{L} = \mathbf{F} \cup \mathbf{R} \cup \mathbf{C}$$

of countable sets of symbols, where

- \mathbf{F} is a set of function symbols

- **R** is a set of relation symbols
- **C** is a set of constant symbols

Next, we recall the notion of a structure in a language.

Definition 1.1.2 (Structure). Let **L** be a structure. An **L-structure** is a tuple

$$M = \langle U; F, R, C \rangle$$

consisting of a non-empty set U and functions, relations, and constants that live in **F**, **R** and **C** respectively. U known as the **universe** of M , and is denoted by $|M|$. The function, relation and constant symbols of M are denoted F^M , R^M and C^M respectively.

Any function or relation in a structure has an **arity**, which is informally the number of arguments it takes. An important fact to note is that arities are not a feature of functions and relations themselves, but of their corresponding *symbols*. In other words, **arity is a syntactic notion**. Semantically speaking, when we seek an interpretation of a function symbol of some arity n , we are forced to limit our search to the set of functions from U^n to U .

We now describe the notion of structure-preserving bijections, known as isomorphisms.

Definition 1.1.3 (Isomorphism). Let **L** be a language and let M, N be **L**-structures. We say that a function $g : |M| \rightarrow |N|$ is an **isomorphism** if

1. g is a bijection
2. “ g commutes with functions”
3. “ g commutes with relations”
4. “ g agrees on constants”

where the double-quotes for the second and third point above refer to the fact that we implicitly require an equality of arities condition before we can talk about composing isomorphisms with multi-ary functions.

We are now ready to define submodels.

Definition 1.1.4 (Submodel). Let M, N be \mathbf{L} -structures. We say that M is a submodel of N , denoted $M \subseteq N$, if

1. $|M| \subseteq |N|$.
2. For all function symbols $F(x_1, \dots, x_n)$, the interpretation in M agrees with the interpretation in N .
3. For all relation symbols $R(a_1, \dots, a_n)$, the interpretation in M agrees with the interpretation in N .
4. For all constant symbols C , the interpretation in M agrees with the interpretation in N .

In particular, a submodel of a model is also a model. For instance, if G is a group, ie, a model of the group axioms, then any submodel of G is, in fact, a subgroup of G , and a group in its own right (in that it again models the group axioms).

We now talk about more syntactic elements of a language.

1.1.2 Syntax: Terms, Formulae, Sentences and Theories

Definition 1.1.5 (Terms). Let \mathbf{L} be a language. **Term**(\mathbf{L}) is defined to be the minimal set of finite sequences of symbols^a from

$$\{ (,), [,] \} \cup \mathbf{C} \cup \mathbf{F} \cup \{x_1, x_2, x_3, \dots\}$$

satisfying the following rules:

1. Every constant symbol is a term.
2. Every variable is a term.
3. For all n -ary functions f and n terms t_1, \dots, t_n , $f(t_1, \dots, t_n)$ is a term.
4. Every term arises in this way.

^aHere, the set of variable symbols is countable, but we might, on occasion, need uncountably many variable symbols

Remark. In other words, the elements of **Term**(\mathbf{L}) are exactly the constants, variables, and functions of such (constants and variables).

Recall, from Definition 1.1.3, that isomorphisms are, in particular, bijections that agree on con-

stants. It is possible to show that they also agree on the interpretations of terms in models. We will show this later.

In similar fashion, we can define the formulae in a language.

Definition 1.1.6 (Formulae). Let \mathbf{L} be a language. $\mathbf{Fml}(\mathbf{L})$ is defined to be the minimal set of finite sequences of symbols from

$$\mathbf{Term}(\mathbf{L}) \cup \{\wedge, \vee, \rightarrow, \neg, \dots\} \cup \{=\} \cup \{\forall, \exists\}$$

satisfying the following rules:

1. For all $\tau_1, \tau_2 \in \mathbf{Term}(\mathbf{L})$, $\tau_1 = \tau_2$ is a formula.
2. For all n -ary relations R and n terms t_1, \dots, t_n , $R(t_1, \dots, t_n)$ is a formula.
3. For all connectives \star and formulae Φ and Ψ , $\Phi \star \Psi$ is a formula.
4. For a quantifier Q , variable x and formula $\varphi(x)$, $Qx(\varphi(x))$ is a formula.
5. Every formula arises in this way.

Formulae consisting only of a single relation symbol (including formulae that only consist of an equality) are called **atomic formulae**. The atomic formulae of \mathbf{L} are denoted $\mathbf{AFml}(\mathbf{L})$.

For a formula φ , denote by $\mathbf{FV}(\varphi)$ the set of free variables of φ . It is sometimes useful to distinguish those formulae in a language that contain no free variables.

Definition 1.1.7. Define the set of **sentences** of a language \mathbf{L} to be

$$\mathbf{Sent}(\mathbf{L}) := \{\varphi \in \mathbf{Fml}(\mathbf{L}) \mid \mathbf{FV}(\varphi) = \emptyset\}$$

Essentially, a formula with no free variables is called a sentence. A theory is simply a set of sentences.

Definition 1.1.8 (Theory). Let \mathbf{L} be a language. An \mathbf{L} -theory is any subset $T \subseteq \mathbf{Sent}(\mathbf{L})$.

There are many well-known theories in mathematics. The most familiar examples come from algebra.

Example 1.1.9 (The Theory of Fields). The theory of fields is a first-order theory in the language of fields. This is a language with function symbols $+$, \times , $^{-1}$, relation symbol $=$, and constant symbols 0 and 1 . It also has other first-order symbols, such as quantifiers, connectives and punctuation, but we ignore these (indeed, we will always take for granted that these exist). The theory of fields consists of the following sentences in this language:

1. $\forall x \forall y \forall z [(x + y) + z = x + (y + z)]$
2. $\forall x \forall y [x + y = y + x]$
3. $\forall x \exists y [x + y = 0] \wedge \forall x [x + 0 = x]$
4. $\forall x [x \neq 0 \rightarrow \exists y [xy = 1]]$, where \neq is the obvious shorthand
5. $\forall x [x \times 1 = x]$
6. $\forall x \forall y \forall z [x \times (y + z) = x \times y + x \times z]$

Collectively, these sentences are known as the **theory of fields**.

There are numerous well-known examples of fields. There is the question of how we can formally describe what it means for some structure, such as the rational numbers, to *satisfy* the above sentences. To that end, we introduce the semantics of interpretation, namely, the notion of **satisfaction**.

1.1.3 Semantics: Satisfaction

We begin with the most important definition of this entire section.

Definition 1.1.10 (Satisfaction - Sentences). Let \mathbf{L} be a language and M an \mathbf{L} -structure.

For any $\varphi \in \mathbf{Fml}(\mathbf{L})$, we say that M **models** φ , denoted $M \models \varphi$, if **sorry**

Remark. We note that, (a) $M \models \forall x \varphi(x) \iff N \models \neg \exists x \neg \varphi(x)$ and (b) $M \models \exists x \varphi(x) \iff M \models \neg \forall x \neg \varphi(x)$. This is worthy of note as for proofs, we will often need to induct on (the number of symbols in) formulas - just as we only need the logical connectives \neg, \vee to get the rest, we only need to check satisfaction for a single quantifier and \neg .

We can define satisfaction for theories in the obvious way.

Definition 1.1.11 (Satisfaction - Theories). Let \mathbf{L} be a language and M an \mathbf{L} -structure. Given an \mathbf{L} -theory T , we say that $M \models T$ if $M \models \psi$ for all $\psi \in T$.

We are now ready to state a simple-sounding but rather non-trivial result.

Lemma 1.1.12. *Suppose M and N are both \mathbf{L} -structures. If $M \subseteq N$, then for all $\tau \in \mathbf{Term}(\mathbf{L})$, $\tau^M[a] = \tau^N[a]$, where $a \in |M| \times \cdots \times |M|$ and $\tau^M[a]$ and $\tau^N[a]$ denote interpretations of τ in M and N with the variables all being interpreted as the components of a .*

We do not prove this result. It is not difficult.

Next is a less trivial result.

Lemma 1.1.13. *If $M \subseteq N$, then $M \models \varphi$ if and only if $N \models \varphi$ for all quantifier-free formulae φ .*

Going back to Example 1.1.9, we can now say the following.

Example 1.1.14 (Models of the Theory of Fields). Recall the theory of fields, seen in Example 1.1.9. It can be shown that the following structures in the language of fields satisfy the theory of fields:

$$\mathbb{Q}, \mathbb{R}, \mathbb{C}, \mathbb{Z}/7\mathbb{Z}$$

A model of the theory of fields is known simply as a **field**. We all know that this is an incredibly rich theory. It might have been even richer if Évariste Galois had had better control over his faculties...

1.1.4 Elementary Equivalence

Recall the definition of an isomorphism of structures (Definition 1.1.3). We can regard isomorphism as a *syntactic* notion of equivalence of structures. In this subsection, we explore a *semantic* notion of equivalence of models.

Definition 1.1.15 (Elementary Equivalence). Let \mathbf{L} be a language and let M and N be \mathbf{L} -structures. We say M is **elementarily equivalent** to N if for every sentence $\varphi \in \mathbf{Sent}(\mathbf{L})$, we have that M satisfies φ if and only if N satisfies φ . We denote this

$$M \equiv N$$

We can now relate isomorphisms to equivalence in the following manner.

Theorem 1.1.16. *Let \mathbf{L} be a language and let M and N be \mathbf{L} -structures. If $M \cong N$, then for every $\varphi(x_1, \dots, x_n) \in \mathbf{Fml}(\mathbf{L})$ and every $a_1, \dots, a_n \in |M|$, we have that*

$$M \models \varphi[a_1, \dots, a_n] \iff N \models \varphi[f(a_1), \dots, f(a_n)]$$

Proof. Fix a formula $\varphi(x_1, \dots, x_n) \in \mathbf{Fml}(\mathbf{L})$ with n free variables. We prove the result by performing induction¹ on φ .

Suppose φ is atomic. Then, there are two cases.

- $\varphi(x_1, \dots, x_n)$ is of the form $\tau_1(x_1, \dots, x_n) = \tau_2(x_1, \dots, x_n)$ for terms $\tau_1, \tau_2 \in \mathbf{Term}(\mathbf{L})$. In this case, it is possible to show, by cases on what τ_1 and τ_2 can be (see Definition 1.1.5) that f is compatible with φ .
- φ is of the form $R(x_1, \dots, x_n)$ for some $R \in \mathbf{R}(\mathbf{L})$. This is true by definition of an isomorphism (see Definition 1.1.3).

Suppose, now, that φ is not atomic. It is enough to show that the result shows if φ is of the form $\psi_1(x_1, \dots, x_n) \wedge \psi_2(x_1, \dots, x_n)$, $\psi_1(x_1, \dots, x_n) \vee \psi_2(x_1, \dots, x_n)$, $\neg\psi(x_1, \dots, x_n)$, and $\forall x_1 \forall x_2 \dots \forall x_n \psi(x_1, \dots, x_n)$, as these would be adequate.

- φ is of the form $\psi_1(x_1, \dots, x_n) \wedge \psi_2(x_1, \dots, x_n)$. This is immediate from the definition of satisfaction: for all $a_1, \dots, a_n \in |M|$, we have that

$$M \models \varphi[a_1, \dots, a_n] \iff M \models \psi_1[a_1, \dots, a_n] \text{ and } M \models \psi_2[a_1, \dots, a_n]$$

¹That is, cases

$$\Longleftrightarrow N \models \psi_1[f(a_1), \dots, f(a_n)] \text{ and } N \models \psi_2[f(a_1), \dots, f(a_n)]$$

$$\Longleftrightarrow N \models \varphi[f(a_1), \dots, f(a_n)]$$

as required.

- φ is of the form $\psi_1(x_1, \dots, x_n) \vee \psi_2(x_1, \dots, x_n)$. Similar.
- φ is of the form $\neg\psi(x_1, \dots, x_n)$. **sorry**
- φ is of the form $\exists x_1, \psi(x_1, \dots, x_n)$.² Fix $a_1, \dots, a_n \in |M|$. Then,

$$M \models \varphi[a_1, \dots, a_n] \Longleftrightarrow M \models \exists x \varphi[x, a_2, \dots, a_n]$$

$$\Longleftrightarrow \text{There is some } a \in |M| \text{ such that } M \models \varphi[b, a_2, \dots, a_n]$$

$$\Longleftrightarrow \text{There is some } a \in |M| \text{ such that } N \models \varphi[f(b), f(a_2), \dots, f(a_n)]$$

$$\Longleftrightarrow \text{There is some } b \in |N| \text{ such that } N \models \varphi[b, f(a_2), \dots, f(a_n)]$$

$$\Longleftrightarrow N \models \exists y \varphi[y, f(a_2), \dots, f(a_n)]$$

where we note that the ' \Rightarrow ' direction of the fourth \Longleftrightarrow comes from taking $b = f(a)$ and the ' \Leftarrow ' direction comes from the fact that f is surjective, meaning that we can take a to be any element of $|M|$ such that $f(a) = b$.

We can conclude by noting that the above cases are adequate. See **sorry**. □

Corollary 1.1.17. *Let \mathbf{L} be a language and let M and N be \mathbf{L} -structures. If $M \cong N$, then $M \equiv N$.*

Proof. Let $f : M \xrightarrow{\sim} N$ be an isomorphism from M to N . **sorry** □

We warn the reader that this implication is strict.

Warning. The notion of isomorphism is finer than elementary equivalence!

²This is enough because you can induct on the number of free variables, with exactly this being the inductive step.

The reason for this is that in elementary equivalence, we do not insist that the interpretations in M and N are the same! That is, M could be a sub-structure of N in which the interpretations used for elementary equivalence *differ*. In the case of isomorphism, we insist that the interpretations are the same (or rather, bijective with f).

We end our discourse on first-order logic by briefly discussing the theory of deduction and proof.

1.1.5 Deduction and Proof

Let \mathbf{L} be a language. Recall that $\mathbf{Sent}(\mathbf{L})$ is the set of *sentences* in \mathbf{L} . Throughout this subsection, fix a theory $T \subseteq \mathbf{Sent}(\mathbf{L})$.

Definition 1.1.18 (Provability). We say a sentence $\varphi \in \mathbf{Sent}(\mathbf{L})$ is **provable from T** , denoted $T \vdash \varphi$, if there exists a sequence $\langle \varphi_1, \dots, \varphi_n \rangle$ of \mathbf{L} -sentences such that $\varphi_n = \varphi$ and for all $i < n$, either $\varphi_i \in T$ or φ_i is obtained from $\langle \varphi_1, \dots, \varphi_{i-1} \rangle$ via the standard deduction rules of first-order logic, namely, Modus Ponens and Generalisation.

We can say something about what makes T a “sensible” set from which to deduce things.

Definition 1.1.19 (Consistency). We say that T is **consistent** if there is no $\varphi \in \mathbf{Sent}(\mathbf{L})$ such that $T \vdash \varphi$ and $T \vdash \neg\varphi$.

Consistency is equivalent to model existence.

Theorem 1.1.20 (Gödel-Henkin). *T is consistent if and only if there is an \mathbf{L} -structure M such that $M \models T$.*

We do not prove this theorem here, but we will make extensive use of it.

We end by recalling the compactness theorem for first-order logic.

Theorem 1.1.21 (Compactness, Gödel-Malcev). *If for any finite $T_0 \subseteq T$, there is a*

We now discuss the *size* of a model and a theory.

1.2 Cardinality and Categoricity

Throughout this section, let \mathbf{L} be a language.

Definition 1.2.1 (Cardinality of a Structure). Let M be an \mathbf{L} -structure. The **cardinality** of M , denoted $\|M\|$, is the cardinality of its universe $|M|$.

We can also talk about the size of a theory.

Definition 1.2.2 (Categoricity of a Theory). Let T be an \mathbf{L} -theory. Suppose $\lambda \geq |L|$ is a cardinal. We say that T is **λ -Categorical**, or that T is **categorical in λ** , if for all \mathbf{L} -structures M and N such that $M, N \models T$ and $\|M\| = \|N\| = \lambda$, we have that $M \cong N$.

Categoricity brings up interesting questions, such as the so-called *spectrum problem*.

1.2.1 The Spectrum Problem

The spectrum of a theory with respect to a cardinal is defined as follows.

Definition 1.2.3 (Spectrum). Let T be an \mathbf{L} -theory and let λ be a cardinal. We define the **spectrum of T with respect to λ** to be

$$I(\lambda, T) := |\{M/\cong \mid M \models T \text{ and } \|M\| = \lambda\}|$$

ie, $I(\lambda, T)$ denotes the number of isomorphism classes of models of T of cardinality λ .

It is obvious, from Definition 1.2.2, that a theory T is λ -categorical if and only if $I(\lambda, T) = 1$. However, if T is not λ -categorical, then it is, in general, quite difficult to compute $I(\lambda, T)$. In fact, for most theories and cardinals, computing the spectrum is an *open problem*, referred to as the **spectrum problem**.

There has been some progress on this problem. Steinitz made the following determinations.

Theorem 1.2.4 (Steinitz). Let \mathbf{L} be the language of fields, and let T be the theory of algebraically closed fields of characteristic p (obtained by adding the appropriate sentences

to the Theory of Fields encountered in Example 1.1.9). Then,

1. $I(\aleph_0, T) = \aleph_0$.
2. For all $\lambda > \aleph_0$, $I(\lambda, T) = 1$.

The spectrum problem has been worked on by some of the most eminent logicians of our time, including Rami's advisor, Saharon Shelah, who proved a famous conjecture by Morley (1965). More on the Spectrum Problem can be found on the associated [Wikipedia page](#), and while this is not the most authoritative source, its contents are nonetheless interesting.

Morley also proved a famous conjecture by Łos from the 1950s, which since became known as Morley's Categoricity Theorem.

Theorem 1.2.5 (Morley's Categoricity Theorem, Morley 1965). *Let T be a theory in a language \mathbf{L} . Assume that $|\mathbf{L}| \leq \aleph_0$. If $\exists \lambda > \aleph_0$ such that T is λ -categorical, then $\forall \lambda > \aleph_0$, T is λ -categorical.*

One of our objectives in this course is to prove Morley's Categoricity Theorem.

As a side note, Morley was initially a PhD student of Saunders MacLane's at the University of Chicago. Morley didn't initially finish his PhD, to the point of losing his stipend at Chicago, but somehow landed a job at Berkeley, where he proved this famous theorem. MacLane, a staunch category theorist, didn't believe Morley's work was quite enough to merit a PhD; nevertheless, after being persuaded by the then-nascent (and very excited) model theory community, he eventually relented and awarded Morley his degree.

Here, we end our discussion on the spectrum problem. Before proceeding further, we recall the basics of cardinal arithmetic.

1.2.2 Cardinal Arithmetic

We begin by introducing notation.

Notation. We denote by

- **ZF** the Zermelo-Fraenkel Axioms of Set Theory
- **AC** the Axiom of Choice
- **ZFC** the Zermelo-Fraenkel Axioms with the Axiom of Choice

We denote cardinality of a set A by $|A|$ or $\text{card}(A)$ and write $|A| = |B|$ if and only if there is a bijection from A to B . Informally, a **cardinal** is a measure of cardinality. That is, a set λ is a cardinal if $\lambda = |A|$ for some set A . We denote by \aleph_0 the cardinal of the natural numbers, which we will denote ω in any cardinal- or ordinal-theoretic context.

There are more precise ways in which we can define the notions of ordinals and cardinals. We do not do this here, but we mention that there is an appendix in Rami's book and several sections in my undergrad logic lecture notes [that discuss this](#).

Add
refer-
ences

Definition 1.2.6 (Cardinal Arithmetic). Let λ, μ be cardinals, with $\lambda = |A|$ and $\mu = |B|$.

We denote

$$\lambda + \mu := \text{sorry}$$

$$\lambda \cdot \mu := |A \times B|$$

The following is a famed theorem of Tarski, a direct consequence of which is precisely the fundamental theorem of cardinal arithmetic.

Theorem 1.2.7 (Tarski). *We can make the following deduction:*

$$\mathbf{ZF} \vdash (\mathbf{AC} \leftrightarrow \forall A, |A| \geq \aleph_0 \rightarrow |A \times A| = |A|)$$

Equivalently,

$$\mathbf{ZF} \vdash (\mathbf{AC} \leftrightarrow \forall A, \lambda \geq \aleph_0 \rightarrow \lambda \cdot \lambda = \lambda)$$

The fundamental theorem of cardinal arithmetic, which states that $|\omega \times \omega| = |\omega|$, is clearly just the specialisation of the above result to the case where $\lambda = \aleph_0$.

There is another fact that will be important for our purposes.

Theorem 1.2.8. *For infinite cardinals $\lambda, \mu \geq \aleph_0$, we have*

$$\lambda \cdot \mu = \max(\lambda, \mu) = \lambda + \mu$$

The reason for discussing cardinal arithmetic is that we can exploit it to prove the existence of submodels of specific cardinalities.

1.3 A Word on Submodels

Fix a language \mathbf{L} .

1.3.1 Submodel Existence

We begin by defining the cardinality of a structure.

Definition 1.3.1 (Cardinality of a Structure). Let M be a \mathbf{L} -structure. We define $\text{card}(M)$ to be the cardinality of the union of

We begin with the famed submodel theorem.

Theorem 1.3.2 (The Submodel Theorem [MOAB]). *Let M be a \mathbf{L} -structure. Define $\lambda := |\mathbf{L}| + \aleph_0$. If $A \leq |M|$, then there exists a substructure $N \leq M$ such that*

- (a) $\text{card}(N) \geq A$
- (b) $\text{card}(|N|) \leq |A| + \lambda$

Proof. By recursion on $n < \omega$, define sets $\{B_n \subseteq |M| \mid n < \omega\}$ such that

1. $B_0 = \{c \in \mathbf{C}^M \mid c \text{ is a constant symbol of } c\} \cup A$
2. If $n < \omega$, then $|B_n| \leq |A| + \lambda$
3. For all $n < \omega$, define $B_{n+1} := \{F^M(\bar{a}) \mid \bar{a} \in B_n\} \cup B_n$

This is enough: if we have such a sequence of B_n , then we could take $B := \bigcup_{n < \omega} B_n$ and define $N := \langle B, F^M, R^M, C^M \rangle$. We can show that this satisfies the desired conditions.

(a) **sorry**(b) **sorry**

Given these, all that remains now is to show that this is possible. **sorry** □

 Finish
using
text-
book
proof

1.3.2 Elementary Submodels

Recall the definition of elementary substructures (**sorry**). In this subsection, we define an analogous notion for models.

Definition 1.3.3 (Elementary Submodels). Let M, N be \mathbf{L} -structures. We say that M is an **elementary submodel** of N , denoted $M \preceq N$, if

1. $M \subseteq N$
2. $M \models \varphi[a_1, \dots, a_n]$ iff $N \models \varphi[a_1, \dots, a_n]$ for every $\varphi \in \mathbf{Fml}(\mathbf{L})$ and $a_1, \dots, a_n \in |M|$.

We can relate this to the notion of elementary substructures in the following manner.

Theorem 1.3.4 (Tarski-Vaught 1956). *Let M, N be \mathbf{L} -structures with $M \subseteq N$. If $M \preceq N$, then $M \equiv N$.*

The converse is not true.

Counterexample 1.3.5. **sorry**

1.3.3 Chains of Substructures

Throughout this subsection, fix a linear order (\mathcal{I}, \leq) .

Definition 1.3.6 (Chain). Let $\{M_i \mid i \in \mathcal{I}\}$ be \mathbf{L} -structures. We say they form a **chain** if for all $i_1, i_2 \in \mathcal{I}$, if $i_1 < i_2$ then $M_{i_1} \subseteq M_{i_2}$.

For the remainder of this subsection, fix a chain $\{M_i \mid i \in \mathcal{I}\}$. It is easy to show that

$$N := \bigcup_{i \in \mathcal{I}} M_i$$

is an \mathbf{L} -structure as well. Moreover, $M_i \subseteq N$ for all $i \in \mathcal{I}$.

We can ask ourselves a natural question: suppose T is a theory in \mathbf{L} and that $\forall i \in \mathcal{I}, M_i \models T$. Is it necessarily true that $N \models T$ as well?

The answer turns out to be no when $|\mathcal{I}| \geq \aleph_0$.

Counterexample 1.3.7. Take $I = \omega$ and

1.3.4 The Löwenheim-Skolem Theorems

1.4 Models of Peano Arithmetic

In this section, we build many models of Peano arithmetic, the most standard and universal of which is ω , the natural numbers.

1.4.1 Restrictions and Expansions

Before going any further, we will need to define a central tool: restrictions and expansions. Throughout this subsection, fix a language L and an L -structure M .

Definition 1.4.1 (Restriction/Expansion). Let $L_1 \subseteq L$, so that L contains relations, functions, and constants $\mathbf{R}(L_1)$, $\mathbf{F}(L_1)$, and $\mathbf{C}(L_1)$. The **restriction of M to L** is the L_1 -structure

$$M|_{L_1} := \langle |M|, \mathbf{R}^M(L_1), \mathbf{F}^M(L_1), \mathbf{C}^M(L_1) \rangle$$

Dually, we say that M is the **expansion of $M|_{L_1}$ to L** .

A good example of this is to model-theoretically encode the fact that every field is also an abelian group (additively).

Example 1.4.2 (Restriction: Fields to Abelian Groups). Let L be the language of fields

and let L_1 be the language of (additively expressed) abelian groups. Then, if

$$M = \langle \mathbb{Q}, +, \cdot, 0, 1 \rangle$$

then its restriction to L_1 is

$$M|_{L_1} = \langle \mathbb{Q}, +, 0 \rangle$$

1.4.2 The Language and Theory of Peano Arithmetic

In this subsection, we set up our study of Peano arithmetic by defining the language in which we will work and the theory we will seek to model. We begin with some notation.

Notation. Let

1.4.3 Existence of Non-Standard Models of Peano Arithmetic

We know that there is a standard model of Peano arithmetic, denoted N . This consists intuitively of the natural numbers—that is, ordinals less than ω —with the usual addition, multiplication, additive and multiplicative identities. It turns out there are also other models, whose existence we will see in this subsection.

Before that, we mention that any model of Peano arithmetic admits a linear order.

Theorem 1.4.3. *The following is a valid deduction.*

$$\mathbf{PA} \vdash \forall x \forall y \exists z (y = x + z)$$

Theorem 1.4.4. *There exists some $M \models \mathbf{PA}$ such that M is not isomorphic to the standard model N .*

Proof. Augment the language L to the $L_1 = L_{\mathbf{PA}} \cup \{c\}$ by adding a new constant symbol c . Let

$$T_1 := \mathbf{PA} \cup \{c \neq 0\} \cup \left\{ c \neq \underbrace{1 + \cdots + 1}_{n \text{ times}} \mid n < \omega \right\}$$

We show, using the Compactness Theorem (**sorry**), that there is a model M of T_1 . Given $T_0 \subseteq T_1$ finite, let $n_0 < \omega$ be the largest natural number such that

$$c \neq \underbrace{1 + \cdots + 1}_{n \text{ times}}$$

lies in T_0 . Define the L_1 -structure

$$M_0 := \langle \omega, +, \cdot, 0, 1, a \rangle$$

be the expansion of N to L_1 , with $c^{M_0} = a$. Then, $M_0 \models T_0$. Hence, since every finite subset of T_1 has a model, so does T_1 . Call this model M_1 . Denote by M the restriction of M_1 to $L_{\mathbf{PA}}$.

Now that we have established the existence of M , all that remains is to show that M is not isomorphic to N . Let $b = c^M$. Suppose $M \cong N$ via an isomorphism f . Then, $f(b) \in \omega$, so there is some $n < \omega$ such that

$$f(b) = \underbrace{1 + \cdots + 1}_{n \text{ times}}$$

□

1.4.4 Countable Models of Peano Arithmetic

Recall the definition of the *spectrum of a theory* (Definition 1.2.3). It turns out, we can use the idea of a spectrum to say something rather sophisticated about countable models of Peano Arithmetic.

Theorem 1.4.5. $I(\aleph_0, \mathbf{PA}) = 2^{\aleph_0}$.

1.4.5 Impossible Deductions

We know that \mathbf{PA} admits a model (indeed, a standard model). Therefore, we know that \mathbf{PA} is a consistent theory. However, Gödel famously showed that it is not possible to prove consistency of \mathbf{PA} using \mathbf{PA} alone. That is, he proved the following.

Theorem 1.4.6 (Gödel's Second Incompleteness Theorem). $\mathbf{PA} \nvdash \mathbf{PA}$ is consistent

Remark. Note that no number theorist would accept this as an example of a statement of interest in number theory which is not provable in \mathbf{PA} (so says the model theorist).

Another interesting impossibility result involves Ramsey theorem.

Theorem 1.4.7 (Paris-Harrington 1976). $\mathbf{PA} \nvdash$ A special case of Ramsey's Theorem.

Chapter 2

Another Chapter

You get the idea.

2.1 Introducing the Main Object of Study in this Chapter

Woah. Very cool.

2.2 Another Section

Yup, \lipsum time. Boy do I love L^AT_EX!

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

Nulla malesuada porttitor diam. Donec felis erat, congue non, volutpat at, tincidunt tristique, libero. Vivamus viverra fermentum felis. Donec nonummy pellentesque ante. Phasellus adipiscing semper elit. Proin fermentum massa ac quam. Sed diam turpis, molestie vitae, placerat a, molestie nec, leo. Maecenas lacinia. Nam ipsum ligula, eleifend at, accumsan nec, suscipit a, ipsum. Morbi blandit ligula feugiat magna. Nunc eleifend consequat lorem. Sed lacinia nulla vitae enim. Pellentesque tincidunt purus vel magna. Integer non enim. Praesent euismod nunc eu purus. Donec bibendum quam in tellus. Nullam cursus pulvinar lectus. Donec et mi. Nam vulputate metus eu enim. Vestibulum pellentesque felis eu massa.

Quisque ullamcorper placerat ipsum. Cras nibh. Morbi vel justo vitae lacus tincidunt ultrices. Lorem ipsum dolor sit amet, consectetur adipiscing elit. In hac habitasse platea dictumst. Integer tempus convallis augue. Etiam facilisis. Nunc elementum fermentum wisi. Aenean placerat. Ut imperdiet, enim sed gravida sollicitudin, felis odio placerat quam, ac pulvinar elit purus eget enim. Nunc vitae tortor. Proin tempus nibh sit amet nisl. Vivamus quis tortor vitae risus porta vehicula.

Fusce mauris. Vestibulum luctus nibh at lectus. Sed bibendum, nulla a faucibus semper, leo velit ultricies tellus, ac venenatis arcu wisi vel nisl. Vestibulum diam. Aliquam pellentesque, augue quis sagittis posuere, turpis lacus congue quam, in hendrerit risus eros eget felis. Maecenas eget erat in sapien mattis porttitor. Vestibulum porttitor. Nulla facilisi. Sed a turpis eu lacus commodo facilisis. Morbi fringilla, wisi in dignissim interdum, justo lectus sagittis dui, et vehicula libero dui cursus dui. Mauris tempor ligula sed lacus. Duis cursus enim ut augue. Cras ac magna. Cras nulla. Nulla egestas. Curabitur a leo. Quisque egestas wisi eget nunc. Nam feugiat lacus vel est. Curabitur consectetur.

Suspendisse vel felis. Ut lorem lorem, interdum eu, tincidunt sit amet, laoreet vitae, arcu. Aenean faucibus pede eu ante. Praesent enim elit, rutrum at, molestie non, nonummy vel, nisl. Ut

lectus eros, malesuada sit amet, fermentum eu, sodales cursus, magna. Donec eu purus. Quisque vehicula, urna sed ultricies auctor, pede lorem egestas dui, et convallis elit erat sed nulla. Donec luctus. Curabitur et nunc. Aliquam dolor odio, commodo pretium, ultricies non, pharetra in, velit. Integer arcu est, nonummy in, fermentum faucibus, egestas vel, odio.

Sed commodo posuere pede. Mauris ut est. Ut quis purus. Sed ac odio. Sed vehicula hendrerit sem. Duis non odio. Morbi ut dui. Sed accumsan risus eget odio. In hac habitasse platea dictumst. Pellentesque non elit. Fusce sed justo eu urna porta tincidunt. Mauris felis odio, sollicitudin sed, volutpat a, ornare ac, erat. Morbi quis dolor. Donec pellentesque, erat ac sagittis semper, nunc dui lobortis purus, quis congue purus metus ultricies tellus. Proin et quam. Class aptent taciti sociosqu ad litora torquent per conubia nostra, per inceptos hymenaeos. Praesent sapien turpis, fermentum vel, eleifend faucibus, vehicula eu, lacus.

Visit <https://thefundamentaltheorem.github.io/ModelTheoryNotes/main.pdf> for the latest version of these notes. If you have any suggestions or corrections, please feel free to fork and make a pull request to my repository.