≮Back to Week 1 **XLessons** Prev Next

Matrix-Matrix Multiplication

We multiply two matrices by breaking it into several vector multiplications and concatenating the result.

$$egin{bmatrix} a & b \ c & d \ e & f \end{bmatrix} * egin{bmatrix} w & x \ y & z \end{bmatrix} = egin{bmatrix} a*w+b*y & a*x+b*z \ c*w+d*y & c*x+d*z \ e*w+f*y & e*x+f*z \end{bmatrix}$$

An **m x n matrix** multiplied by an **n x o matrix** results in an **m x o** matrix. In the above example, a 3 x 2 matrix times a 2 x 2 matrix resulted in a 3 x 2 matrix.

To multiply two matrices, the number of **columns** of the first matrix must equal the number of **rows** of the second matrix.

For example:

```
% Initialize a 3 by 2 matrix
 A = [1, 2; 3, 4; 5, 6]
 4 % Initialize a 2 by 1 matrix
 5 B = [1; 2]
7 % We expect a resulting matrix of (3 \text{ by } 2)*(2 \text{ by } 1) = (3 \text{ by } 1)
                                                                               Run
8 mult_AB = A*B
                                                                               Reset
10 % Make sure you understand why we got that result
```

✓ Complete







2/9/2018, 12:15 PM