

Sequence measures protein identity π vs dN/dS
ρ = −0.76 to −0.89
Count evidence S_g vs protein identity π
genes with count evidence vs dN/dS
|ρ| ≤ 0.14
|ρ| ≤ 0.17
Δα vs protein identity π
all genes vs dN/dS
|ρ| ≤ 0.18
|ρ| ≤ 0.19
Liver
Testis
Bone marrow
Kidney
Islet
Eye
Cortex
Lung
−1 −0.8 −0.6 −0.4 −0.2 0 +0.2
Spearman correlation across genes, ρ (one mark per tissue)
(a)
Liver
Testis
Bone marrow
Kidney
Islet
Eye
Cortex
Lung
Δα (centered)
+0.25
0
−0.25
ρ = +0.09
ρ = −0.10
ρ = +0.04
ρ = +0.15
ρ = +0.18
ρ = +0.04
ρ = +0.12
ρ = −0.01
0.4 0.7 1
protein identity π
ρ = −0.13
ρ = +0.12
ρ = −0.02
ρ = −0.14
ρ = −0.19
ρ = −0.06
ρ = −0.13
ρ = −0.03
−2 −1 0
log10 dN/dS
count evidence
no count evidence (imputed from features)
median trend
ρ Spearman correlation of Δα with the x axis
(b)