

Sequence measures
protein identity π vs dN/dS
ρ = −0.76 to −0.89
Count evidence S_g
vs protein identity π
|ρ| ≤ 0.14
genes with count evidence
vs dN/dS
|ρ| ≤ 0.17
Δα
vs protein identity π
|ρ| ≤ 0.18
all genes
vs dN/dS
|ρ| ≤ 0.19
Liver
Testis
Bone marrow
Kidney
Islet
Eye
Cortex
Lung
−1
−0.8
−0.6
−0.4
−0.2
0
+0.2
Spearman correlation across genes, ρ (one mark per tissue)
(a)
Liver
Testis
Bone marrow
Kidney
Islet
Eye
Cortex
Lung
Δα (centered)
0.25
0.00
−0.25
ρ = +0.09
ρ = −0.10
ρ = +0.04
ρ = +0.15
ρ = +0.18
ρ = +0.04
ρ = +0.12
ρ = −0.01
0.4
0.7
1.0
protein identity π
ρ = −0.13
ρ = +0.12
ρ = −0.02
ρ = −0.14
ρ = −0.19
ρ = −0.06
ρ = −0.13
ρ = −0.03
−2
−1
0
dN/dS (log10)
count evidence
no count evidence (imputed from features)
median trend
ρ Spearman correlation of Δα with the x axis
(b)