

ECM: other (n = 55)
Homeobox TFs (n = 58)
Signaling kinases (n = 105)
Forkhead and other TFs (n = 112)
Microtubules (n = 56)
DNA replication and repair (n = 112)
Cell cycle (n = 100)
Zinc-finger TFs (n = 114)
Nuclear receptors (n = 121)
Splicing factors (n = 70)
OXPHOS, nuclear-encoded (n = 66)
Mitochondrial ribosome (n = 58)
2
1
2
1
3
8
1
1
−0.2
−0.1
0
+0.1
Δα (liver 4 species)
(a)
shift
amplitude
+0.51★
−0.20
+0.19★
−0.52
−0.23
+0.18
−0.15
+0.14
−0.49★
+0.13
−0.53★
+0.18
−0.40★
+0.17
−0.53★
+0.06
−0.49★
−0.16★
−0.60★
+0.14
−0.78★
−0.06★
−1.38★
−0.76★
1
0.5
0
−0.5
−1
effect size
Cohen's d of Δα and of |Δα| (signed)
(b)
urea cycle / gluconeogenesis
mitochondrially encoded OXPHOS
enzymes passing every check
supporting tissue
conserved controls
† no margin support
F3†
PDGFRB†
CR2†
VCAM1†
KIT†
C1QC†
DDX4
EEF2
NDUFA1
COX4I1
HNF4A†
RPL3
CYP7A1
ARG2
HAL
PCK1
MT-CO1
ASL
MT-ND2
MT-ND4
MT-CYB
CPS1†
ARG1†
MT-CO3
MT-ATP6
PAH†
MT-CO2
0.2
0.1
0.0
−0.1
−0.2
−0.3
−0.4
−0.5
Δα (liver 4 species)
0.2
0.4
0.6
0.8
1.0
protein prior π
(c)