

Kidney: AKI programs in mouse IRI
label-null p
human · mouse
PT
TAL
EC
FIB
PC
myeloid
DCT
IC
lymphoid
0.005 · 0.010
0.025 · 0.045
0.015 · 0.005
0.030 · 0.025
0.040 · 0.050
0.025 · 0.075
0.33 · 0.040
0.27 · 0.010
1.00 · 1.00
0
50
100
150
200
programs significant in human AKI
transfers to mouse
does not transfer
opposite direction
(a)
IRI 2 d vs 4–20 wk
no 2 d fit
no stage fit
no 2 d fit
no stage fit
0
100
200
transferring programs
0.0
0.4
0.8
program NES, Spearman
mouse IRI 2 d
mouse IRI 4–20 wk
(b)
Healthy-tissue test: does Δα predict transfer?
kidney: AKI vs IRI
PT
TAL
EC
FIB
PC
myeloid
DCT
IC
lymphoid
liver: MASH vs Western diet
hepatocyte
islet: T2D vs db/db, mSTZ
β
α
δ
ductal
endothelial
stellate
−0.08
−0.04
0
0.04
0.08
Δα vs gene-level transfer, ρ_p (95% CI)
(c)
Program transfer vs label null
observed, above null 95%
observed, within null
human-label null, median–95%
0
50
100
150
200
transferring programs
(d)
Cippà 2018 injury genes
acute lists
(primary)
late lists
0.51
0.51
Chimera Δα
profile correlation
0.70
0.72
0.4
0.5
0.6
0.7
0.8
stratified AUROC,
shared vs human-specific
(e)