
median
curated M class
34 genes
1.57
curated F class
20 genes
1.67
curated reference class
16 genes
2.13
0
2
4
6
per-gene mean |attribution|

(e)


genes
δ > 0
per-gene
p
curated M class
17,790 residues
68%
0.010 *
curated F class
12,095 residues
55%
0.927 ns
curated reference class
6,947 residues
62%
0.433 ns
0.00
0.04
0.08
Cliff's δ (hydrophobic vs polar)
per-gene p < 0.05
p ≥ 0.05

(f)


pooled Mann-Whitney over all residues
per-gene paired Wilcoxon
p = 0.05
−log10 p (square-root scale)
60
30
10
5
2
1
0
9.6×10⁻⁵⁷ ***
0.622 ns
0.004 **
0.013 *
0.240 ns
0.197 ns
1.4×10⁻⁴ ***
0.302 ns
2.4×10⁻¹¹ ***
0.404 ns
1.8×10⁻¹⁴ ***
0.174 ns
disorder:
IDR vs ordered
(δ = +0.16)
contact density:
low vs high
(δ = +0.05)
special residues:
aromatic vs rest
(δ = +0.04)
special residues:
glycine vs rest
(δ = +0.05)
special residues:
proline vs rest
(δ = +0.09)
conservation:
entropy high vs low
(δ = +0.06)

(g)