
residue feature
contrast
pooled p
per-gene p
n_g
pLDDT
very high (>90) vs very low (<50)
8.7×10⁻⁶² ***
0.701 ns
63
pLDDT
high (70–90) vs low (50–70)
8.2×10⁻⁹ ***
0.922 ns
68
pLDDT
very high (>90) vs low (50–70)
1.9×10⁻¹⁴ ***
0.494 ns
68
side-chain chemistry
positive vs negative
0.008 **
0.097 ns
70
secondary structure
strand vs coil
0.002 **
0.518 ns
70
solvent accessibility
intermediate vs buried
0.474 ns
0.030 *
70
secondary structure
helix vs coil
0.002 **
0.956 ns
70
secondary structure
helix vs strand
0.852 ns
0.723 ns
70
side-chain chemistry
hydrophobic vs polar
6.0×10⁻⁹ ***
0.043 *
70
special residues
aromatic vs rest
0.240 ns
0.197 ns
70
special residues
glycine vs rest
1.4×10⁻⁴ ***
0.302 ns
70
contact density
low vs high
0.004 **
0.013 *
59
conservation
entropy high vs low
1.8×10⁻¹⁴ ***
0.174 ns
35
contact density
low vs medium
9.5×10⁻¹² ***
0.0496 *
70
side-chain chemistry
hydrophobic vs positive
2.7×10⁻²⁶ ***
2.3×10⁻⁴ ***
70
side-chain chemistry
hydrophobic vs negative
1.2×10⁻¹² ***
0.036 *
70
special residues
proline vs rest
2.4×10⁻¹¹ ***
0.404 ns
70
solvent accessibility
exposed vs buried
2.4×10⁻³⁸ ***
0.090 ns
70
solvent accessibility
exposed vs intermediate
2.7×10⁻³⁶ ***
0.671 ns
70
disorder
IDR vs ordered
9.6×10⁻⁵⁷ ***
0.622 ns
63
−0.1
0.0
0.1
0.2
Cliff's δ (positive: first level has larger |attribution|)
per-gene $p < 0.05$ (within-protein test)
per-gene $p \geq 0.05$