
residue feature
contrast
pooled p
per-gene p
n_g
pLDDT
very high (>90) vs very low (<50)
8.7×10⁻⁶² *** 0.701 ns 63
pLDDT
high (70–90) vs low (50–70)
8.2×10⁻⁹ *** 0.922 ns 68
pLDDT
very high (>90) vs low (50–70)
1.9×10⁻¹⁴ *** 0.494 ns 68
side-chain chemistry
positive vs negative
0.008 ** 0.097 ns 70
secondary structure
strand vs coil
0.002 ** 0.518 ns 70
solvent accessibility
intermediate vs buried
0.474 ns 0.030 * 70
secondary structure
helix vs coil
0.002 ** 0.956 ns 70
secondary structure
helix vs strand
0.852 ns 0.723 ns 70
side-chain chemistry
hydrophobic vs polar
6.0×10⁻⁹ *** 0.043 * 70
special residues
aromatic vs rest
0.240 ns 0.197 ns 70
special residues
glycine vs rest
1.4×10⁻⁴ *** 0.302 ns 70
contact density
low vs high
0.004 ** 0.013 * 59
conservation
entropy high vs low
1.8×10⁻¹⁴ *** 0.174 ns 35
contact density
low vs medium
9.5×10⁻¹² *** 0.0496 * 70
side-chain chemistry
hydrophobic vs positive
2.7×10⁻²⁶ *** 2.3×10⁻⁴ *** 70
side-chain chemistry
hydrophobic vs negative
1.2×10⁻¹² *** 0.036 * 70
special residues
proline vs rest
2.4×10⁻¹¹ *** 0.404 ns 70
solvent accessibility
exposed vs buried
2.4×10⁻³⁸ *** 0.090 ns 70
solvent accessibility
exposed vs intermediate
2.7×10⁻³⁶ *** 0.671 ns 70
disorder
IDR vs ordered
9.6×10⁻⁵⁷ *** 0.622 ns 63
−0.1
0.0
0.1
0.2
Cliff's δ (positive: first level has larger |attribution|)
per-gene p < 0.05 (within-protein test)
per-gene p ≥ 0.05