

Δα, imputed
Forward margin
Sdir, direct
Ridge on protein, target S
Liver H+Mouse, pattern
Liver H+Monkey, pattern
Liver H+Zebrafish, pattern
Liver 4 species, pattern
Testis H+Mouse, pattern
Testis H+Macaque, pattern
Testis 3 species, pattern
Kidney, pattern
Islet, pattern
Liver H+Mouse, level
Liver 4 species, level
Testis H+Mouse, level
Testis 3 species, level
Kidney, level
−0.2
−0.1
0.0
0.1
0.2
ρp with independent gold, masked genes
(a)
Chimera gate, protein only
π
Ridge on protein, target S
seed against other seeds
Nearest proteins, their Δα
Liver H+Mouse
Testis H+Mouse
Liver 4-sp
ρp on held-out genes, given C
1.0
0.8
0.6
0.4
0.2
0.0
−0.2
−0.4
measured Δα
measured S
measured Sdir
pattern gold
(b)
Δα, imputed
Sdir, direct
Forward margin
Ridge on protein, target S
B1, where defined
Δα, anchor scaled by depth
Liver H+Mouse
Liver 4-sp
Testis H+Mouse
ρp, pattern gold, newly sparse
0.25
0.20
0.15
0.10
0.05
0.00
−0.05
−0.10
25%
10%
5%
(c)