


canonical gate
per-gene gate
Δα effect vs matched genes (SD)
−0.6
0.0
0.6
Liver
Testis
Bone marrow
Kidney
Islet
Eye
Cortex
Lung
π z
dN/dS z
B1 z
Glycoprotein [PTM] -19 +15 +24
Secreted [CC] -27 +23 +19
external encapsulating structure (CC) -9 +7 +18
molecular transducer activity (MF) -8 +3 +19
Integrin cell surface interactions (R) -8 +10 +12
KRAS Signaling Up -7 +4 +9
plasma membrane (CC) +9 -17 +30
signaling (BP) +15 -21 +23
Calcium [Ligand] +4 -9 +12
Signal Transduction (R) +35 -41 +18
Hemostasis (R) +9 -10 +15
Cytoskeleton & ECM (category) +8 -9 +13
nucleoplasm (CC) +29 -26 -15
RNA binding (MF) +14 -12 -9
Metabolism of RNA (R) +24 -25 -12
RNA-binding [MF] +12 -10 -7
mRNA processing [BP] +19 -21 -7
Chromatin & DNA processing (category) +12 -12 -7
Mitochondrion [CC] -23 +26 -12
DNA repair (BP) -5 +9 -12
DNA damage [BP] -7 +9 -12
catalytic activity, acting on DNA (MF) -4 +9 -6
Diseases of DNA repair (R) -7 +9 -4
fast protein, above prior
conserved protein, above prior
conserved protein, below prior
fast protein, below prior

(a)


programme pole, mean effect (SD units, matched genes)
above-prior programmes
below-prior programmes
0.6
0.4
0.2
0.0
−0.2
−0.4
6
29
90
429
divergence from human (My)
agreement with other tissues' programme order (ρ)
1.0
0.9
0.8
0.7
0.6
0.5
0.4
0.3
0.2
cortex
liver
canonical
eye
testis
per-gene

(b)