
residue feature
contrast
pooled p
per-gene p
n_g
pLDDT very high (>90) vs very low (<50) 8.7e-62 *** 0.701 ns 63
pLDDT high (70–90) vs low (50–70) 8.2e-09 *** 0.922 ns 68
pLDDT very high (>90) vs low (50–70) 1.9e-14 *** 0.494 ns 68
chemistry positive vs negative 0.008 ** 0.097 ns 70
secondary structure sheet vs coil 0.002 ** 0.518 ns 70
solvent accessibility intermediate vs buried 0.474 ns 0.030 * 70
secondary structure helix vs coil 0.002 ** 0.956 ns 70
secondary structure helix vs sheet 0.852 ns 0.723 ns 70
chemistry hydrophobic vs polar 6.0e-09 *** 0.043 * 70
special residues aromatic vs rest 0.240 ns 0.197 ns 70
special residues glycine vs rest 1.4e-04 *** 0.302 ns 70
contact density low vs high 0.004 ** 0.013 * 59
conservation entropy high vs low 1.8e-14 *** 0.174 ns 35
contact density low vs medium 9.5e-12 *** 0.050 * 70
chemistry hydrophobic vs positive 2.7e-26 *** 2.3e-04 *** 70
chemistry hydrophobic vs negative 1.2e-12 *** 0.036 * 70
special residues proline vs rest 2.4e-11 *** 0.404 ns 70
solvent accessibility exposed vs buried 2.4e-38 *** 0.090 ns 70
solvent accessibility exposed vs intermediate 2.7e-36 *** 0.671 ns 70
disorder IDR vs ordered 9.6e-57 *** 0.622 ns 63
−0.1
0.0
0.1
0.2
Cliff's δ (positive: first level has larger |attribution|)
per-gene $p < 0.05$ (within-protein test)
per-gene $p \geq 0.05$