
residue feature
contrast
pooled p
per-gene p
n_g
pLDDT
very high (>90) vs very low (<50)
8.7e-62 ***
0.701 ns
63
pLDDT
high (70–90) vs low (50–70)
8.2e-09 ***
0.922 ns
68
pLDDT
very high (>90) vs low (50–70)
1.9e-14 ***
0.494 ns
68
chemistry
positive vs negative
0.008 **
0.097 ns
70
secondary structure
sheet vs coil
0.002 **
0.518 ns
70
solvent accessibility
intermediate vs buried
0.474 ns
0.030 *
70
secondary structure
helix vs coil
0.002 **
0.956 ns
70
secondary structure
helix vs sheet
0.852 ns
0.723 ns
70
chemistry
hydrophobic vs polar
6.0e-09 ***
0.043 *
70
special residues
aromatic vs rest
0.240 ns
0.197 ns
70
special residues
glycine vs rest
1.4e-04 ***
0.302 ns
70
contact density
low vs high
0.004 **
0.013 *
59
conservation
entropy high vs low
1.8e-14 ***
0.174 ns
35
contact density
low vs medium
9.5e-12 ***
0.050 *
70
chemistry
hydrophobic vs positive
2.7e-26 ***
2.3e-04 ***
70
chemistry
hydrophobic vs negative
1.2e-12 ***
0.036 *
70
special residues
proline vs rest
2.4e-11 ***
0.404 ns
70
solvent accessibility
exposed vs buried
2.4e-38 ***
0.090 ns
70
solvent accessibility
exposed vs intermediate
2.7e-36 ***
0.671 ns
70
disorder
IDR vs ordered
9.6e-57 ***
0.622 ns
63
−0.1
0.0
0.1
0.2
Cliff's δ (positive: first level has larger |attribution|)
per-gene $p < 0.05$ (within-protein test)
per-gene $p \geq 0.05$