

mean scaled rank within task
0 best
0.5
1 worst
not applicable or not run
5/15 runs finished / runs tried
level change
pattern change
loss
null: placebo sets
null: sham move
null: both species thinned
unperturbed: reference
unperturbed: agreement
label quality: reference
label quality: agreement
composition: reference
composition: agreement
sampling: reference
sampling: agreement
technical: reference
technical: agreement
pattern
pattern: held-out primates
pattern: sorted lineages
pattern: second atlas
pattern: second atlas, nephron
pattern: no acinar cells
pattern: second atlas, no acinar
level: bulk tissue
level: atlases
named gene lists
run time
run time, fewer cells
run time, full pipeline
peak memory
number of tasks
18 18 18 9 12 12
7 3 21 9 14 6 14 6 14 6
99 33 10 22 11 11 11 88 66 62
5 3 3 5
A ground truth
B robustness
C independent data
D cost
overall scaled rank
0 best; bar = 1 − rank
Chimera readouts
profile correlation, clusters of Chimera's latent
Δα, per-gene gate
count evidence S_g
Δα, protein-feature gate
5/15 2/6
direct baselines
specificity similarity
rank-matched level per cell type
profile correlation
profile correlation (residual)
labeled direct margin, no model
fold change
gene features alone
other tools
SAMap
CAMEX gene-embedding cosine
GeneSpectra
CACIMAR
SATURN gene score
sequence measures
LOEUF
protein identity π
dN/dS
0.45
0.47
0.56
0.68
0.29
0.32
0.32
0.36
0.39
0.43
0.50
0.27
0.57
0.61
0.65
0.74
0.60
0.62
0.68