
§L.M.2 Every §L contrast ranked · bar colour = within-gene paired Wilcoxon p
(70 genes, 36832 residues)
§IDR IDR vs ordered
§SASA exposed vs intermediate
§SASA exposed vs buried
§AA proline vs rest
§Chem hydrophobic vs negative
§Chem hydrophobic vs positive
§Contact low vs medium
§Cons entropy high vs low
§Contact low vs high
§AA glycine vs rest
§AA aromatic vs rest
§Chem hydrophobic vs polar
§SS helix vs sheet
§SS helix vs coil
§SASA intermediate vs buried
§SS sheet vs coil
§Chem positive vs negative
§pLDDT very high (>90) vs low (50–70)
§pLDDT high (70–90) vs low (50–70)
§pLDDT very high (>90) vs very low (<50)
ns per-gene p=0.622 · pooled p=9.6e-57
ns per-gene p=0.671 · pooled p=2.7e-36
ns per-gene p=0.090 · pooled p=2.4e-38
ns per-gene p=0.404 · pooled p=2.4e-11
* per-gene p=0.036 · pooled p=1.2e-12
*** per-gene p=2.3e-04 · pooled p=2.7e-26
* per-gene p=0.050 · pooled p=9.5e-12
ns per-gene p=0.174 · pooled p=1.8e-14
* per-gene p=0.013 · pooled p=0.004
ns per-gene p=0.302 · pooled p=1.4e-04
ns per-gene p=0.197 · pooled p=0.240
* per-gene p=0.043 · pooled p=6.0e-09
ns per-gene p=0.723 · pooled p=0.852
ns per-gene p=0.956 · pooled p=0.002
* per-gene p=0.030 · pooled p=0.474
ns per-gene p=0.518 · pooled p=0.002
ns per-gene p=0.097 · pooled p=0.008
ns per-gene p=0.494 · pooled p=1.9e-14
ns per-gene p=0.922 · pooled p=8.2e-09
ns per-gene p=0.701 · pooled p=8.7e-62
paired p < 0.001 (***)
paired p < 0.01 (**)
paired p < 0.05 (*)
paired p ≥ 0.05 (NS)
−0.15
−0.10
−0.05
0.00
0.05
0.10
0.15
0.20
Cliff's δ (positive = first group has larger |attn|) · error bar = 95 % bootstrap CI