

---

# **SIMULATION DES WA-TOR RÄUBER-BEUTE MODELLS UND ERWEITERUNGEN**

---

12. April 2017

Sebastian von der Thannen  
Thomas Heitzinger

TECHNISCHE UNIVERSITÄT WIEN

# Inhaltsverzeichnis

<b>1</b>	<b>Das klassische Wa-Tor Programm</b>	<b>3</b>
<b>2</b>	<b>Das stetige Wa-Tor Programm</b>	<b>3</b>
2.1	Schwarmverhalten . . . . .	3
<b>3</b>	<b>Animationen</b>	<b>3</b>
<b>4</b>	<b>Helmholtz Equation</b>	<b>3</b>
4.1	General . . . . .	3
	<b>Literatur</b>	<b>5</b>

# 1 Das klassische Wa-Tor Programm

## 2 Das stetige Wa-Tor Programm

### 2.1 Schwarmverhalten

## 3 Animationen

## 4 Helmholtz Equation

### 4.1 General

The Helmholtz equation is a partial differential equation of elliptic type.

$$\begin{aligned}\nabla^2 u + \kappa^2 u &= 0 \\ \text{or } \Delta u + \kappa^2 u &= 0\end{aligned}\tag{4.1}$$

where  $\Delta$  is the Laplace Operator with  $\Delta u := \sum_{i=1}^d \frac{\partial^2 u}{\partial x_i^2}$ .

We are interested in radial symmetric solutions to the Helmholtz equation. In order to find them it is helpful to transform equation (4.1) into polar coordinates for the two dimensional case or spherical coordinates for the three dimensional case.

**Lemma 4.1.** *The Laplace operator in  $\mathbb{R}^2$  is given by  $\Delta u = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2}$ . The Cartesian coordinates can be represented by polar coordinates as follows:*

$$\begin{cases} x = r \cos \theta \\ y = r \sin \theta \end{cases}$$

Using these substitutions it can be shown that:

$$\Delta u = \frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2}$$

Similarly, in  $\mathbb{R}^3$  we transform to a spherical coordinate system:

$$\begin{cases} x = r \sin \theta \cos \varphi \\ y = r \sin \theta \sin \varphi \\ z = r \cos \theta \end{cases}$$

which yields the form:

$$\Delta u = \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial u}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial u}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 u}{\partial \varphi^2}\tag{4.2}$$

Helmholtz' equation is obtained by adding the term  $\kappa^2 u$ . Since we are solely interested in radial symmetric solutions we would like to eliminate the angular variables  $\theta$  and  $\varphi$  from the equation. This leads us to:

## Abbildungsverzeichnis



## Literatur

- [1] Milton Abramowitz und Irene A. Stegun. *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*. ninth Dover printing, tenth GPO printing. New York: Dover, 1964, S. 364.
- [2] Jean-Pierre Berenger. “A Perfectly Matched Layer for the Absorption of Electromagnetic Waves”. In: *J. Comput. Phys.* 114.2 (Okt. 1994), S. 185–200. ISSN: 0021-9991. DOI: 10.1006/jcph.1994.1159. URL: <http://dx.doi.org/10.1006/jcph.1994.1159>.
- [3] Andreas Kirsch und Frank Hettlich. *The Mathematical Theory of Maxwell’s Equations*. <http://www.math.kit.edu/ianmip/lehre/sobraum2014s/media/main.pdf>. Accessed: 2016-04-15.
- [4] Eric W. Weisstein. *Helmholtz Differential Equation – Polar Coordinates*. <http://mathworld.wolfram.com/HelmholtzDifferentialEquationPolarCoordinates.html>. Accessed: 2016-04-08.
- [5] Peter Young. *Helmholtz’s and Laplace’s Equations in Spherical Polar Coordinates: Spherical Harmonics and Spherical Bessel Functions*. [http://physics.ucsc.edu/~peter/116C/helm\\_sp.pdf](http://physics.ucsc.edu/~peter/116C/helm_sp.pdf). Accessed: 2016-01-18.